University of Groningen

Real-life evidence in ERS clinical practice guidelines

Fally, Markus; Nagavci, Blin; Tonia, Thomy; Berge, Maarten van den; Bush, Andrew; Brightling, Christopher; Roche, Nicolas

Published in:
European Respiratory Journal

DOI:
[10.1183/13993003.01718-2021](https://doi.org/10.1183/13993003.01718-2021)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Fally, M., Nagavci, B., Tonia, T., Berge, M. V. D., Bush, A., Brightling, C., & Roche, N. (2021). Real-life evidence in ERS clinical practice guidelines: from foes to friends. *European Respiratory Journal*, 58(2), [2101718]. <https://doi.org/10.1183/13993003.01718-2021>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Real-life evidence in ERS clinical practice guidelines: from foes to friends

Markus Fally^{1,2}, Blin Nagavci^{2,3}, Thomy Tonia^{2,4}, Maarten van den Berge⁵, Andrew Bush^{2,6}, Christopher Brightling⁷ and Nicolas Roche^{2,8}

¹Dept of Respiratory Medicine, Copenhagen University Hospital – Bispebjerg and Frederiksberg, Copenhagen, Denmark. ²European Respiratory Society Guidelines Working Group, Switzerland. ³Institute for Evidence in Medicine, Medical Center – University of Freiburg, Freiburg, Germany. ⁴Institute of Social and Preventive Medicine, University of Bern, Bern, Switzerland. ⁵Groningen Research Institute for Asthma and COPD (GRIAC), University of Groningen, University Medical Center Groningen, Groningen, The Netherlands. ⁶Dept of Paediatric Respiratory Medicine, Imperial College and Royal Brompton Hospital, London, UK. ⁷Dept of Respiratory Sciences, University of Leicester, Leicester, UK. ⁸Dept of Pulmonology, Cochin Hospital, Assistance Publique – Hôpitaux de Paris (AP-HP), University of Paris, Paris, France.

Corresponding author: Nicolas Roche (nicolas.roche@aphp.fr)

Shareable abstract (@ERSpublications)

This editorial discusses 1) why data from real life should be included in clinical practice guidelines, and 2) how to integrate this real-life evidence in the guideline development process
<https://bit.ly/3wdLBDH>

Cite this article as: Fally M, Nagavci B, Tonia T, *et al.* Real-life evidence in ERS clinical practice guidelines: from foes to friends. *Eur Respir J* 2021; 58: 2101718 [DOI: 10.1183/13993003.01718-2021].

Copyright ©The authors 2021. For reproduction rights and permissions contact permissions@ersnet.org

Received: 17 June 2021
Accepted: 22 June 2021

Introduction

Throughout the past decades, European Respiratory Society (ERS) task forces have produced and published clinical practice guidelines (CPGs), statements, technical standards and other documents to synthesise and summarise bodies of evidence for caregivers, and thereby improve healthcare quality in respiratory medicine. Among these various types of documents, only CPGs can propose recommendations for clinical practice. As such, they need to rely on a very strong methodology to limit the risk of recommending suboptimal care.

Traditionally, the highest levels of evidence come from randomised controlled trials (RCTs) [1, 2]. However, this does not mean that other types of research should be excluded from evidence synthesis as part of guidelines development processes. Indeed, they could usefully complement RCTs, provided that their methods are rigorous, and the results are properly analysed and transparently interpreted [3, 4]. In this editorial, we summarise how real-life evidence could and should be integrated in ERS CPGs.

The development of ERS clinical practice guidelines

The first crucial task of an ERS task force developing a CPG is to carefully consider the research questions and outcomes of interest. Only thereafter, a systematic review of the literature, as well as an assessment of the quality of evidence, can be performed (figure 1). For the latter, the ERS uses Grading of Recommendations, Assessment, Development and Evaluation (GRADE), an approach adopted and recommended by many organisations including the National Institute for Health and Care Excellence, the American Thoracic Society and the World Health Organization [5]. This rigorous method considers a number of factors, in addition to risk of bias, for assessing the quality of evidence. Moreover, it ensures a transparent link between evidence and recommendations when applying the Evidence to Decision (EtD) framework for grading of the strength of a recommendation [6].

The GRADE approach can be used for data originating from RCTs as well as observational studies [5]. When following the GRADE approach, the developers of guidelines must evaluate the risk of bias, inconsistency, indirectness and imprecision, as well as publication bias, to assess the certainty of evidence (table 1) [5]. This applies to data from both RCTs and observational studies. For the latter, large effect

FIGURE 1 The European Respiratory Society (ERS) clinical practice guideline development process at a glance. PICO: Population, Intervention, Comparison, Outcome; GRADE: Grading of Recommendations, Assessment, Development and Evaluation.

sizes, dose responses and opposing biases may lead to an upgrading of the quality of evidence. This systematic approach that takes every aspect of a published study into account makes it possible to ultimately adjust and grade the quality of evidence as “very low”, “low”, “moderate” or “high”. In a next step, the GRADE EtD framework allows for additional considerations, such as balance of benefits and harms, values and preferences, feasibility, equity, acceptability and resource use [6].

TABLE 1 Factors that impact the quality of evidence in randomised controlled trials and observational studies

Factors that can reduce the quality of evidence in randomised controlled trials and observational studies	
Risk of bias	Limitations in the study design and execution
Inconsistency	Unexplained heterogeneity of results
Indirectness	Indirect comparisons or differences in study populations, interventions or outcomes
Imprecision	Wide confidence intervals due to few patients and few events
Publication bias	Systematic under- or overestimation of a beneficial or harmful effect due to selective publication of studies
Factors that can increase the quality of evidence in observational studies	
Large magnitude of an effect	Point estimates for relative risks or hazard ratios way below or above 1
Dose–response gradient	The presence of a dose–response gradient may increase the confidence in the results
Plausible residual confounding	The absence of residual confounding would have increased the intervention’s effects

Adapted from the GRADE (Grading of Recommendations, Assessment, Development and Evaluation) handbook [5].

This systematic approach makes it possible for the expert panel to transparently draw their final conclusions while taking various aspects and perspectives into consideration, and make recommendations that are supported by the evidence [7].

Real-life evidence

The historical understanding that RCTs produce the evidence with the highest quality has often been challenged because patient populations in these studies often are selected, not reflecting the patients seen in everyday clinical practice [8]. Hence, clinically important data from real life may be missed and not sufficiently emphasised by healthcare professionals and policymakers. A brilliant example of the disparity between patients included in RCTs and real-life cohorts has recently been published by BROWN *et al.* [9]. When comparing data from 342 patients against trial eligibility criteria from 37 RCTs evaluating biological therapies for severe asthma, less than 10% of patients in their real-life cohort were found to be eligible [9]. Similar concerns have been reported regarding other major lung diseases, for example COPD, lung cancer and bronchiectasis [10–16]. In such cases, the recommendations derived from RCTs might not be applicable to most patients.

Although the ERS applies a very strict methodological approach and considers both randomised and observational studies, the generalisability of our CPGs might be improved further by the inclusion of real-life evidence from other sources, such as administrative databases or healthcare registers [3]. These sources can complement RCTs by: 1) confirming or challenging their generalisability for different populations or settings, 2) exploring clinically relevant outcomes not available in RCTs, 3) providing safety data, and 4) allowing to explore possible determinants of treatment effects to be further confirmed in prospective RCTs [3]. In some cases, when it is not possible to perform an RCT due to, for example, ethical or feasibility reasons, real-life evidence might even be the only way to generate new data.

The ERS promotes an integrative approach in science and guideline development

Several ERS initiatives were implemented recently to close the gap that often exists between evidence generated from RCTs and real life. With the ERS Clinical Research Collaborations (CRCs), the respiratory community has the possibility to build international research networks to conduct pan-European pragmatic trials that are generalisable and sufficiently large to impact clinical practice (<https://www.ersnet.org/science-and-research/clinical-research-collaboration-application-programme/>). Furthermore, the ERS offers investigators the opportunity to promote their research by endorsing pragmatic trials (<https://www.ersnet.org/science-and-research/pragmatic-trials-endorsement/>). Pragmatic trials can be endorsed when they investigate respiratory diseases, meet stringent criteria of quality and are not dependent on a single sponsor from the pharmaceutical industry or another for-profit entity. In addition, CRCs provide an excellent platform to collaboratively develop and use data from healthcare registers [17]. Hence, the ERS promotes every type of evidence that can lead to a better understanding of a certain respiratory condition, as long as the highest quality standards are satisfied [3].

Conclusion: how to integrate real-life evidence in ERS CPGs

The crucial mission for ERS task forces developing CPGs is to appreciate not only data originating from RCTs, but also other sources, to get the best picture of the current evidence and draw solid conclusions [3, 4]. For CPGs, the ERS Guidelines Working Group recommends a thorough process with the selection of a limited number of clinical questions in a PICO (Population, Intervention, Comparison, Outcome) format that can include data both from RCTs and observational studies, and that are assessed *via* a systematic review and the application of the GRADE process [18]. These questions can be complemented with additional non-comparative questions that are addressed *via* a narrative review of the literature. The guideline panel then chooses outcomes that are critical or important for clinical decision making and that are relevant for the patients. For the final recommendations, the EtD framework should be systematically applied both for PICO and narrative questions. Real-life evidence that has not been considered in the systematic or narrative reviews should be considered and taken into account in the EtD framework [4].

With this approach, we ensure that ERS CPGs give recommendations that are transparent, trustworthy and clinically relevant both for clinicians and patients.

Conflict of interest: M. Fally has nothing to disclose. B. Nagavci has nothing to disclose. T. Tonia acts as a methodologist for ERS. M. van den Berge has nothing to disclose. A. Bush has nothing to disclose. C. Brightling has nothing to disclose. N. Roche reports grants and personal fees from Boehringer Ingelheim, Novartis, GSK and Pfizer, personal fees from AstraZeneca, Chiesi, Sanofi and Zambon, outside the submitted work.

Support statement: M. Fally and B. Nagavci were recipients of a European Respiratory Society (ERS) Fellowship in Guidelines Methodology (MTF 2018-01 and MTF 2018-02).

References

- 1 Prazma CM, Wenzel SE, Price R, *et al.* The impact of duration of oral corticosteroid use on co-morbidities in a severe asthma population treated with mepolizumab. *Am J Respir Crit Care Med* 2015; 191: Suppl., A4279.
- 2 Concato J, Shah N, Horwitz RJ. Randomized, controlled trials, observational studies, and the hierarchy of research designs. *N Engl J Med* 2000; 342: 1887–1892.
- 3 Roche N, Anzueto A, Bosnic Anticevich S, *et al.* The importance of real-life research in respiratory medicine: manifesto of the Respiratory Effectiveness Group. *Eur Respir J* 2019; 54: 1901511.
- 4 Gershon AS, Lindenauer PK, Wilson KC, *et al.* Informing healthcare decisions with observational research assessing causal effect. An Official American Thoracic Society Research Statement. *Am J Respir Crit Care Med* 2021; 203: 14–23.
- 5 Schünemann H, Brożek J, Guyatt G, *et al.* GRADE handbook for grading quality of evidence and strength of recommendations. Updated October 2013. The GRADE Working Group, 2013. Available from: www.guidelinedevelopment.org/handbook
- 6 Alonso-Coello P, Schünemann HJ, Moher J, *et al.* GRADE Evidence to Decision (EtD) frameworks: a systematic and transparent approach to making well informed healthcare choices. 1: Introduction. *BMJ* 2016; 353: i2016.
- 7 Li SA, Alexander PE, Reljic T, *et al.* Evidence to Decision framework provides a structured “roadmap” for making GRADE guidelines recommendations. *J Clin Epidemiol* 2018; 104: 103–112.
- 8 Sherman RE, Anderson SA, Dal Pan GJ, *et al.* Real-world evidence – what is it and what can it tell us? *N Engl J Med* 2016; 375: 2293–2297.
- 9 Brown T, Jones T, Gove K, *et al.* Randomised controlled trials in severe asthma: selection by phenotype or stereotype. *Eur Respir J* 2018; 52: 1801444.
- 10 Halpin DMG, Kerkhof M, Soriano JB, *et al.* Eligibility of real-life patients with COPD for inclusion in trials of inhaled long-acting bronchodilator therapy. *Respir Res* 2016; 17: 120.
- 11 Scichilone N, Basile M, Battaglia S, *et al.* What proportion of chronic obstructive pulmonary disease outpatients is eligible for inclusion in randomized clinical trials? *Respiration* 2014; 87: 11–17.
- 12 Kruis AL, Ställberg B, Jones RCM, *et al.* Primary care COPD patients compared with large pharmaceutically-sponsored COPD studies: an UNLOCK validation study. *PLoS One* 2014; 9: e90145.
- 13 Travers J, Marsh S, Caldwell B, *et al.* External validity of randomized controlled trials in COPD. *Respir Med* 2007; 101: 1313–1320.
- 14 Chalmers JD, McDonnell MJ, Rutherford R, *et al.* The generalizability of bronchiectasis randomized controlled trials: a multicentre cohort study. *Respir Med* 2016; 112: 51–58.
- 15 Heuvels ME, Wisnivesky J, Stricker BH, *et al.* Generalizability of results from the National Lung Screening Trial. *Eur J Epidemiol* 2012; 27: 669–672.
- 16 Pahus L, Burgel P-R, Roche N, *et al.* Randomized controlled trials of pharmacological treatments to prevent COPD exacerbations: applicability to real-life patients. *BMC Pulm Med* 2019; 19: 127.
- 17 Brightling C, Genton C, Bill W, *et al.* ERS clinical research collaborations: underpinning research excellence. *Eur Respir J* 2018; 52: 1801534.
- 18 Huang X, Lin J, Demner-Fushman D. Evaluation of PICO as a knowledge representation for clinical questions. *AMIA Annu Symp Proc* 2006; 2006: 359–363.