University of Groningen

Synthesis of Optically Active beta- or gamma-Alkyl-Substituted Alcohols through Copper-Catalyzed Asymmetric Allylic Alkylation with Organolithium Reagents

Guduguntla, Sureshbabu; Fañanás-Mastral, Martín; Feringa, Bernard

Published in:
Journal of Organic Chemistry

DOI:
[10.1021/jo401536u](https://doi.org/10.1021/jo401536u)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2013

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Guduguntla, S., Fañanás-Mastral, M., & Feringa, B. (2013). Synthesis of Optically Active beta- or gamma-Alkyl-Substituted Alcohols through Copper-Catalyzed Asymmetric Allylic Alkylation with Organolithium Reagents. *Journal of Organic Chemistry*, 78(17), 8274-8280. <https://doi.org/10.1021/jo401536u>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Synthesis of Optically Active β - or γ -Alkyl-Substituted Alcohols through Copper-Catalyzed Asymmetric Allylic Alkylation with Organolithium Reagents

Sureshbabu Guduguntla, Martín Fañanás-Mastral,* and Ben L. Feringa*

Stratingh Institute for Chemistry, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands

S Supporting Information

ABSTRACT: An efficient one-pot synthesis of optically active β -alkyl-substituted alcohols through a tandem copper-catalyzed asymmetric allylic alkylation (AAA) with organolithium reagents and reductive ozonolysis is presented. Furthermore, hydroboration–oxidation following the Cu-catalyzed AAA leads to the corresponding homochiral γ -alkyl-substituted alcohols.

■ INTRODUCTION

Chiral nonracemic alcohols (and derivatives) are very important building blocks in the synthesis of numerous biologically active compounds. In particular, optically active primary alcohols bearing alkyl substitution at the β - or γ -positions are key intermediates in the total synthesis of several natural products including arundic acid,¹ Lyrica,² bongkreic acids,³ gynnastatin A,⁴ and vitamins E and K.⁵ There are a number of methods available for the synthesis of this type of alcohol based on chiral auxiliaries⁶ and enzyme-catalyzed kinetic resolution of racemic compounds.⁷ In 1995, Negishi reported a Zr-catalyzed asymmetric carboalumination of alkenes followed by a lipase catalyzed resolution method to access these building blocks in good yields with excellent enantiomeric excess.⁸ The development of alternative catalytic enantioselective protocols remains an important challenge in view of the potential of these highly versatile building blocks.

Cu-catalyzed AAA is among the most powerful enantioselective C–C bond-forming reactions.⁹ In sharp contrast with the well-known Pd-catalyzed asymmetric allylic alkylation reaction,¹⁰ which is characterized by the use of soft and stabilized nucleophiles, Cu-catalyzed asymmetric allylic alkylation is characterized by the formation of C–C bonds with organometallic reagents, resulting in a complementary method. The reaction usually proceeds with high S_N2' selectivity and provides access to a carbon stereocenter next to a terminal olefin which can readily be further functionalized. Pioneered by Bäckvall and van Koten,¹¹ Cu-catalyzed AAA has been widely studied and its synthetic utility has been shown in the total synthesis of several natural products and biologically active compounds.¹² Recently, our group reported for the first time the use of highly reactive organolithium reagents in copper-catalyzed asymmetric allylic alkylation of allyl bromides with excellent regio- and enantioselectivity using Taniaphos as a chiral ligand.¹³ We also implemented this methodology for both allyl bromides and chlorides in the enantioselective

synthesis of tertiary and quaternary stereocenters using phosphoramidite ligands.¹⁴

Herein we present a highly enantioselective one pot synthesis of β -alkyl-substituted alcohols through Cu-catalyzed AAA of allyl bromides with various organolithium reagents followed by a reductive ozonolysis reaction. The direct use of organolithium reagents is also extended to the synthesis of γ -alkyl-substituted alcohols through Cu-catalyzed AAA of allyl bromides with RLi reagents followed by a hydroboration oxidation reaction (Scheme 1).¹⁵

Scheme 1. Synthesis of Optically Active Alcohols through Copper-Catalyzed Asymmetric Allylic Alkylation with Organolithium Reagents **■ RESULTS AND DISCUSSION**

Our strategy is based on a tandem Cu-catalyzed AAA/reductive ozonolysis to achieve highly enantioenriched β -alkyl-substituted alkyl alcohols in a chemo-, regio-, and enantioselective one-pot operation with no racemization. Using the well-established conditions for the Cu-catalyzed AAA with organolithium reagents,^{13,14} we optimized the conditions for the synthesis of highly enantioenriched β -alkyl-substituted alcohols in a one-pot protocol (Table 1). We started our study with commercially

Received: July 15, 2013

Published: August 21, 2013

Table 1. Optimization Conditions for the One-Pot Cu-Catalyzed Asymmetric Allylic Alkylation Followed by Reductive Ozonolysis

entry	NaBH ₄ (x equiv)	4a:5 (%) ^a
1	2.5	70:30
2	5	90:10
3 ^b	10	100:0

^aThe ratio was determined by ¹H NMR and GC–MS. ^b10 equiv of water was added to the reaction mixture. L1 = (+)-(R,R_p)-Taniaphos (see Table 2).

Table 2. One-Pot Synthesis of β-Alkyl-Substituted Alcohols through Cu-Catalyzed Asymmetric Allylic Alkylation of Allyl Bromides with Organolithium Reagents Followed by Reductive Ozonolysis

1a R = Ph

1b R = *p*-BrC₆H₄1c R = BrCH₂

1d R =

1e R =

entry ^a	1	R'	L	2:3 ^b (%)	4, yield ^c (%)	4, ee ^d (%)
1	1a	Me	L1	90:10	85	4a, 98
2	1a	<i>n</i> -Bu	L1	90:10	60	4b, 99
3	1a	<i>n</i> -Hex	L1	88:12	70	4c, 99
4	1a	<i>i</i> -Bu	L2	88:12	70	4d, 84
5	1b	Me	L1	90:10	84	4e, 99
6	1b	<i>n</i> -Bu	L1	85:15	75	4f, 97
7	1b	<i>n</i> -Hex	L1	87:13	90	4g, 99
8	1c	<i>n</i> -Hex	L1	100:0	60	4h, 97
9	1c	Me	L1	100:0	30 ^e	4i, 72
10	1d	<i>n</i> -Hex	L1	90:10	65	4j, >99:1 (<i>dr</i>) ^f
11	1d	Me	L1	80:20	50	4k, >99:1 (<i>dr</i>) ^f
12 ^g	1e	<i>n</i> -Bu	L1	100:0	73	4l, 97

^aReactions were run on a 0.2–0.5 mmol scale using 1.2 equiv of R'Li diluted with hexane (1.5 equiv diluted with toluene in the case of MeLi) which was added over 2 h using a syringe pump to a 0.1 M solution of substrate in CH₂Cl₂. ^bRatio of S_N2':S_N2 products was determined by GC–MS and ¹H NMR analysis of a sample taken before ozonolysis. ^cThe corresponding alcohol obtained from the S_N2 product could be separated by column chromatography unless otherwise noted (see the Experimental Section). ^dDetermined by chiral HPLC. ^eThe low yield is due to volatility issues. ^f*dr* determined by ¹H NMR. ^g10% of double 1,2-addition product A was isolated in this case.

available cinnamyl bromide 1a. After Cu-catalyzed AAA of 1a,¹³ the reaction mixture was quenched with EtOH and purged with ozone for 20 min followed by purging with nitrogen. When 2.5

equiv of NaBH₄ was added to reduce the ozonide, a mixture of desired alcohol 4a and aldehyde 5 was obtained in a 70:30 ratio (Table 1, entry 1). Doubling the amount of NaBH₄ did not lead

Scheme 2. One-Pot Synthesis of α -Alkyl-Substituted Aldehyde through Cu-Catalyzed Asymmetric Allylic Alkylation of Allyl Bromides with Organolithium Reagents Followed by Ozonolysis

Table 3. Synthesis of γ -Alkyl-Substituted Alcohols through Cu-Catalyzed Asymmetric Allylic Alkylation of Allyl Bromides with Organolithium Reagents Followed by a Hydroboration/Oxidation

1a R = Ph 1b R = *p*-BrC₆H₄ 1f R = BnOCH₂

entry ^a	1	R'	L	2:3 ^b (%)	(2 + 3), yield (%)	6, yield ^c (%)	6, ee ^d (%)
1	1a	<i>n</i> -Bu	L1	90:10	88	67	6a, 99
2	1a	<i>n</i> -Hex	L1	90:10	86	74	6b, 99
3	1b	Me	L1	90:10	90	70	6c, 99
4	1b	<i>n</i> -Hex	L1	87:13	93	90	6d, 99
5	1f	Me	L1	85:15	81	75	6e, 81
6	1f	<i>n</i> -Bu	L1	85:15	85	93	6f, 88

^aReactions were run on a 0.2 mmol scale using 1.2 equiv of R'Li diluted with hexane (1.5 equiv diluted with toluene in the case of MeLi) which was added over 2 h using a syringe pump to a 0.1 M solution of substrate in CH₂Cl₂. ^bRatio of S_N2':S_N2 products was determined by GC–MS and crude ¹H NMR. ^cCalculated based on 2. ^dDetermined by chiral HPLC.

to full conversion toward the desired alcohol either (Table 1, entry 2), probably due to the formation of the corresponding acetal in the reaction mixture which was hydrolyzed during the workup giving rise to aldehyde 5. In order to achieve full conversion to alcohol 4a, 10 equiv of NaBH₄ and 10 equiv of water were used to hydrolyze the acetal in situ (Table 1, entry 3). Under these conditions, no aldehyde 5 was observed, and the desired alcohol 4a was obtained in good overall yield with very high enantioselectivity (see Table 2, entry 1).¹⁶

Having optimized conditions for the one-pot protocol for the synthesis of β -alkyl substituted alcohols, the scope of the reaction was examined. We employed this tandem consecutive Cu-catalyzed AAA/reductive ozonolysis protocol with organolithium reagents such as MeLi, *n*-BuLi, and *n*-HexLi on cinnamyl bromide 1a, achieving excellent enantioselectivities (98–99%) and good overall yields (60–85%) (Table 2, entries 1–3). More hindered reagents, such as *i*-BuLi, could also be used in this tandem application leading to the desired alcohol 4d in 70% overall yield with a high ee of 84% (Table 2, entry 4). It is important to note that phosphoramidite ligand L2¹⁷ had to be used in this case. To show the functional group tolerance of this protocol, we performed the reaction with *p*-bromo-substituted substrate 1b using different organolithium reagents. The tandem reaction provides the desired β -alkyl-substituted alcohols with excellent enantioselectivities 97–99% and high overall yields 75–90% without any traces of side products (i.e., halogen–lithium exchange) (Table 2, entries 5–

7). The allyl bromide 1c bearing an aliphatic bromide (BrCH₂ substituent) was converted with *n*-HexLi affording alcohol 4h in 60% yield and again with excellent enantioselectivity (97% ee) (Table 2, entry 8). A decrease in the enantioselectivity was observed when 1c was treated with MeLi as an alkylating source (Table 2, entry 9). The allyl bromide 1d bearing an acetal protected chiral 1,2-diol functionality upon the tandem application with MeLi and *n*-HexLi provided excellent diastereoselectivity (*anti/syn* ratios of >99:1) (Table 2, entries 10 and 11). An ester functionality is also tolerated, and the one-pot Cu-catalyzed AAA/reductive ozonolysis of 1e led to an exclusive S_N2' substitution that provided the desired alcohol 4l in 73% yield with 97% ee (Table 2, entry 12).

As the results summarized in Table 2 show, a highly versatile one-pot catalytic protocol to access a range of homochiral β -substituted alcohols is now available using common organolithium reagents and allyl bromides. This one-pot protocol, based on readily available compounds, avoids the isolation of the branched alkenes, which can be volatile (especially methyl-substituted compounds), thus affording better overall yields than the corresponding two step version.

It is important to note that when the Cu-catalyzed AAA followed by ozonolysis was performed of 1d with subsequent Me₂S treatment we were able to isolate the corresponding α -alkyl substituted aldehyde in good yield 70% with very high diastereomeric ratio (*anti/syn* = >99:1) without affecting the integrity of the stereogenic center (Scheme 2).

We also explored a similar strategy for the synthesis of highly enantioenriched γ -alkylated alcohols via Cu-catalyzed AAA followed by a hydroboration/oxidation reaction. First, we performed the Cu-catalyzed AAA on cinnamyl bromide **1a** using *n*-BuLi and *n*-HexLi. The corresponding olefins were formed in good yields of 88% and 86%, respectively, with excellent enantioselectivity. The hydroboration/oxidation reaction of these olefins using commercially available 9-BBN led to the corresponding γ -alkylated alcohols **6a** and **6b** in good yields (67% and 74%) without erosion of the enantiomeric excess (Table 3, entries 1 and 2). Olefins bearing a *p*-bromo substituent, obtained via Cu-catalyzed AAA from **1b**, were converted in a hydroboration/oxidation sequence to the corresponding alcohols **6c** and **6d** in 70% and 90% yield, respectively, being optically pure (99% ee) (Table 3, entries 3 and 4). Allyl bromide **1f**, bearing a benzyl ether functionality, was also subjected to the Cu-catalyzed AAA/hydroboration/oxidation sequence providing the corresponding alcohols **6e** and **6f** in good yields (75% and 93%) albeit with slightly lower enantioselectivities (81% and 88% ee, respectively) (Table 3, entries 5 and 6).

As illustrated in Scheme 3, a β -alkyl-substituted aldehyde can also be readily synthesized. For instance, the oxidation of the

Scheme 3. Oxidation of Primary Alcohols to Aldehydes with Dess–Martin Periodinane

corresponding primary alcohol **6e** with Dess–Martin periodinane (DMP) provided aldehyde **8**, an important intermediate in the total synthesis of danshenspiroketallactone.¹⁸

CONCLUSIONS

In summary, we have developed a highly enantioselective synthesis of β -alkyl-substituted alcohols through a one-pot Cu-catalyzed asymmetric allylic alkylation with organolithium reagents followed by reductive ozonolysis. The synthesis of γ -alkyl-substituted alcohols was also achieved through Cu-catalyzed asymmetric allylic alkylation with organolithium reagents followed by a hydroboration oxidation. These protocols do not compromise the stereochemical integrity and provide ready access to highly valuable chiral building blocks.

EXPERIMENTAL SECTION

General Procedures. Chromatography was performed on silica gel (230–400 mesh). Thin-layer chromatography was performed on silica plates. Compounds were visualized by UV and cerium/molybdenum or potassium permanganate staining. Progress and conversion of the reaction were determined by GC–MS. Mass spectra were recorded on a mass spectrometer using an Orbitrap analyzer. ¹H and ¹³C NMR were recorded on 400 and 100.59 MHz using CDCl₃ as solvent. Chemical shift values are reported in ppm with the solvent resonance as the internal standard (CHCl₃: δ 7.26 for ¹H, δ 77.0 for ¹³C). Data are reported as follows: chemical shifts, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, br = broad, m = multiplet), coupling constants (Hz), and integration. Optical rotations were measured on a polarimeter with a 10 cm cell (*c* given in g/100 mL).

Enantiomeric excesses were determined by chiral HPLC analysis using a diode array detector.

All reactions were carried out under a nitrogen atmosphere using oven-dried glassware and using standard Schlenk techniques. All the reagents, starting materials, and ligand **L1** were purchased from commercial sources and used without further purification. Dichloromethane and toluene were used from the solvent purification system. *n*-Hexane was dried and distilled over sodium. Allyl bromides **1b**,¹⁹ **1d**,²⁰ **1e**,²¹ and **1f**²² were prepared following literature procedures. Phosphoramidite ligand **L2** was prepared as reported in the literature.²³

Racemic products were synthesized by reaction of the allyl bromides **1** and the corresponding organolithium reagent at –78 °C in CH₂Cl₂ in the presence of CuI (10 mol %) and PPh₃ (20 mol %).

General Procedure for the One-Pot Synthesis of β -Alkyl Substituted Alcohols through Cu-Catalyzed Asymmetric Allylic Alkylation of Allyl Bromides with Organolithium Reagents Followed by Reductive Ozonolysis. A Schlenk tube equipped with septum and stirring bar was charged with CuBr·SMe₂ (0.01 mmol, 2.06 mg, 5 mol %) and the appropriate ligand (0.012 mmol, 6 mol %). Dry dichloromethane (2 mL) was added, and the solution was stirred under nitrogen at room temperature for 15 min. Then, allyl bromide **1** (0.2 mmol) was added, and the resulting solution was cooled to –80 °C. In a separate Schlenk tube, the corresponding organolithium reagent (0.24 mmol, 1.2 equiv) was diluted with hexane (toluene in the case of MeLi, combined volume of 1 mL) under nitrogen and added dropwise to the reaction mixture over 2 h using a syringe pump. Once the addition was complete, the mixture was stirred for another 2 h at –80 °C. The reaction was quenched with EtOH (2 mL), and then ozone was bubbled through the solution for 20 min. After being stirred for 20 min (solution stays blue), the reaction mixture was purged with nitrogen to remove excess ozone (disappearance of blue color). Sodium borohydride (75.66 mg, 2 mmol, 10 equiv) was added, followed after 10 min by the addition of H₂O (10 equiv). Subsequently, the reaction mixture was warmed to room temperature and stirred overnight. The mixture was quenched by addition of extra water (5 mL). The layers were separated, and the aqueous layer was extracted with DCM (2 × 10 mL). The combined organic layers were dried with sodium sulfate and concentrated in vacuo. The crude product was purified by flash chromatography on silica gel using different mixtures of *n*-pentane/Et₂O as eluent.

Note: The S_N2':S_N2 ratio was determined by GC–MS analysis on a sample obtained after quenching with EtOH, which was passed through a short plug of silica gel to remove transition-metal residues.

(R)-2-Phenylpropan-1-ol (4a). Purification by column chromatography (SiO₂, 10–30% Et₂O/pentane, gradient) afforded an inseparable mixture of **4a** and benzyl alcohol in the ratio of 90:10 (51 mg, yield = 85%) as a colorless oil: 98% ee, [α]_D²⁰ = +5.0 (*c* = 1 in CHCl₃) [lit.²⁴ (97% ee): [α]_D²³ = +16.2 (*c* = 1 in CHCl₃)]; ¹H NMR (400 MHz, CDCl₃) δ 7.42–7.18 (m, 5H), 4.68 (s, 2H), 3.69 (d, *J* = 6.8 Hz, 2H), 2.9–3.0 (m, 1H), 1.63 (s, 1H), 1.28 (d, *J* = 7.1 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 143.7, 140.9, 128.6, 128.5, 127.6, 127.5, 127.0, 126.7, 68.7, 65.3, 42.4, 17.6; HRMS (APCI+, *m/z*) calcd for C₉H₁₁ [M – H₂O]⁺ 119.08553, found 119.08549. Enantiomeric excess was determined by chiral HPLC analysis, Chiralcel OB-H column, *n*-heptane/*i*-PrOH 90:10, 40 °C, 217 nm, retention times (min): 10.40 (major) and 11.11 (minor).

(R)-2-Phenylhexan-1-ol (4b). Purification by column chromatography (SiO₂, 5–20% Et₂O/pentane, gradient) afforded **4b** (25 mg, yield = 60%) as a colorless oil. 99% ee, [α]_D²⁰ = –11.0 (*c* = 1 in CHCl₃); [lit.²⁵ [α]_D²⁰ = –18.0 (*c* = 3.73 in CH₂Cl₂)]; ¹H NMR (400 MHz, CDCl₃) δ 7.38–7.18 (m, 5H), 3.79–3.68 (m, 2H), 2.82–2.72 (m, 1H), 1.75–1.65 (m, 1H), 1.62–1.51 (m, 1H), 1.49–1.11 (m, 4H), 0.84 (t, *J* = 7.3 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 142.5, 128.6, 128.1, 126.7, 67.6, 48.7, 31.8, 29.5, 22.7, 14.0; HRMS (APCI+, *m/z*) calcd for C₁₂H₁₇ [M – H₂O]⁺ 161.13248, found 161.13245. Enantiomeric excess was determined by chiral HPLC analysis, Chiralcel OJ-H column, *n*-heptane/*i*-PrOH 95:5, 40 °C, 220 nm, retention times (min): 13.51 (major) and 14.17 (minor).

(*R*)-2-Phenyl-octan-1-ol (**4c**). Purification by column chromatography (SiO₂, 5–20% Et₂O/pentane, gradient) afforded **4c** (33 mg, yield = 70%) as a colorless oil: 99% ee, $[\alpha]_D^{20} = -14.2$ ($c = 1$ in CHCl₃); [lit.²⁶ (*S*)-enantiomer (92% ee): $[\alpha]_D^{21} = +15.1$ ($c = 0.99$ in CHCl₃)]; ¹H NMR (400 MHz, CDCl₃) δ 7.37–7.18 (m, 5H), 3.81–3.66 (m, 2H), 2.82–2.72 (m, 1H), 1.77–1.63 (m, 1H), 1.62–1.50 (m, 1H), 1.34–1.11 (m, 8H), 0.85 (t, $J = 6.9$ Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 142.5, 128.6, 128.0, 126.6, 67.6, 48.7, 32.0, 31.7, 29.3, 27.3, 22.6, 14.0; HRMS (APCI+, m/z) calcd for C₁₄H₂₁ [M – H₂O]⁺ 189.16378, found 189.16372. Enantiomeric excess was determined by chiral HPLC analysis, Chiralcel OJ-H column, *n*-heptane/*i*-PrOH 95:5, 40 °C, 220 nm, retention times (min): 10.63 (major) and 11.33 (minor).

(*R*)-4-Methyl-2-phenylpentan-1-ol (**4d**). Purification by column chromatography (SiO₂, 5–20% Et₂O/pentane, gradient) afforded **4d** (30 mg, yield = 70%) as a colorless oil: 84% ee, $[\alpha]_D^{20} = -10.2$ ($c = 1$ in CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 7.38–7.19 (m, 5H), 3.76–3.64 (m, 2H), 2.95–2.84 (m, 1H), 1.64–1.51 (m, 1H), 1.49–1.36 (m, 3H), 0.86 (t, $J = 6.7$ Hz, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 142.4, 128.6, 128.1, 126.7, 68.0, 46.4, 41.1, 25.2, 23.5, 21.8; HRMS (APCI+, m/z) calcd for C₁₂H₁₇ [M – H₂O]⁺ 161.13248, found 161.13238. Enantiomeric excess was determined by chiral HPLC analysis, Chiralcel AD-H column, *n*-heptane/*i*-PrOH 95:5, 40 °C, 212 nm, retention times (min): 12.26 (major) and 13.04 (minor).

(*R*)-2-(4-Bromophenyl)propan-1-ol (**4e**). Purification by column chromatography (SiO₂, 10–20% Et₂O/pentane, gradient) afforded an inseparable mixture of **4e** and *p*-bromobenzyl alcohol in a ratio of 90:10 (27 mg, yield = 84%) as a colorless oil: 99% ee, $[\alpha]_D^{20} = +8.6$ ($c = 1$ in CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 7.48 (d, $J = 8.4$ Hz, 2H), 7.44 (d, $J = 8.5$ Hz, 2H), 7.23 (d, $J = 8.4$ Hz, 2H), 7.11 (d, $J = 8.5$ Hz, 2H), 4.64 (s, 2H), 3.72–3.62 (m, 2H), 2.96–2.86 (m, 1H), 1.25 (d, $J = 7.0$ Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 142.7, 139.8, 131.6, 131.6, 129.2, 128.6, 121.4, 120.4, 68.4, 64.5, 41.9, 17.5; HRMS (APCI+, m/z) calcd for C₉H₁₀Br [M – H₂O]⁺ 196.99604, found 196.99617. Enantiomeric excess was determined by chiral HPLC analysis, Chiralcel OB-H column, *n*-heptane/*i*-PrOH 95:5, 40 °C, 224 nm, retention times (min): 14.37 (minor) and 14.9 (major).

(*R*)-2-(4-Bromophenyl)hexan-1-ol (**4f**). Purification by column chromatography (SiO₂, 10–20% Et₂O/pentane, gradient) afforded **4f** (34 mg, yield = 75%) as a colorless oil: 97% ee, $[\alpha]_D^{20} = -13.8$ ($c = 1$ in CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 7.45 (d, $J = 8.4$ Hz, 2H), 7.09 (d, $J = 8.4$ Hz, 2H), 3.77–3.64 (m, 2H), 2.78–2.68 (m, 1H), 1.80–1.60 (m, 1H), 1.58–1.45 (m, 1H), 1.38–1.04 (m, 2H), 0.84 (t, $J = 7.1$ Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 141.6, 131.7, 129.8, 120.4, 67.4, 48.2, 31.6, 29.4, 22.7, 13.9; HRMS (APCI+, m/z) calcd for C₁₂H₁₆Br [M – H₂O]⁺ 239.04299, found 239.04261. Enantiomeric excess was determined by chiral HPLC analysis, Chiralcel AD-H column, *n*-heptane/*i*-PrOH 95:5, 40 °C, 230 nm, retention times (min): 15.04 (major) and 16.01 (minor).

(*R*)-2-(4-Bromophenyl)octan-1-ol (**4g**). Purification by column chromatography (SiO₂, 10–20% Et₂O/pentane, gradient) afforded **4g** (38 mg, yield = 90%) as a colorless oil: 99% ee, $[\alpha]_D^{20} = -13.2$ ($c = 1$ in CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 7.44 (d, $J = 8.4$ Hz, 2H), 7.08 (d, $J = 8.4$ Hz, 2H), 3.77–3.60 (m, 2H), 2.78–2.68 (m, 1H), 1.74–1.61 (m, 1H), 1.60–1.47 (m, 1H), 1.38–1.07 (m, 8H), 0.85 (t, $J = 6.6$ Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 141.7, 131.7, 129.8, 120.4, 67.4, 48.2, 31.9, 31.7, 29.3, 27.2, 22.6, 14.0; HRMS (APCI+, m/z) calcd for C₁₄H₂₀Br [M – H₂O]⁺ 267.07429, found 267.07391. Enantiomeric excess was determined by chiral HPLC analysis, Chiralcel AD-H column, *n*-heptane/*i*-PrOH 95:5, 40 °C, 226 nm, retention times (min): 14.20 (major) and 14.90 (minor).

(*S*)-2-(Bromomethyl)octan-1-ol (**4h**). Purification by column chromatography (SiO₂, 10–30% Et₂O/pentane, gradient) afforded **4h** (15 mg, yield = 60%) as a colorless oil: 97% ee, $[\alpha]_D^{20} = -9.4$ ($c = 1$ in CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 3.50 (dd, $J = 10.5$, 5.8 Hz, 2H), 3.41 (dd, $J = 10.5$, 6.5 Hz, 2H), 1.68–1.53 (m, 1H), 1.39–1.34 (m, 1H), 1.33–1.19 (m, 8H), 1.16–1.02 (m, 1H), 0.88 (t, $J = 6.6$ Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 68.4, 35.7, 33.1, 31.8, 29.6, 26.9, 22.7, 16.7, 14.1; HRMS (ESI+, m/z) calcd for C₉H₂₀BrO [M + H]⁺ 223.09752, found 223.09707. The enantiomeric excess was

determined for the benzoate ester of the alcohol. Enantiomeric excess was determined by chiral HPLC analysis, Chiralcel OD-H column, *n*-heptane/*i*-PrOH 98:2, 40 °C, 232 nm, retention times (min): 8.43 (minor) and 9.06 (major).

(*S*)-3-Bromo-2-methylpropan-1-ol (**4i**). Purification by column chromatography (SiO₂, 10–30% Et₂O/pentane, gradient) afforded **4i** (22.5 mg, yield = 30%) as a colorless oil: 72% ee, $[\alpha]_D^{20} = +1.6$ ($c = 1$ in CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 3.67–3.56 (m, 2H), 3.54–3.45 (m, 2H), 2.08–1.97 (m, 1H), 1.43 (s, 1H), 1.03 (d, $J = 6.8$ Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 65.4, 37.6, 37.3, 15.5. The enantiomeric excess was determined for the benzoate ester of the alcohol. Enantiomeric excess was determined by chiral HPLC analysis, Chiralcel OB-H column, *n*-heptane/*i*-PrOH 100:0, 40 °C, 226 nm, retention times (min): 27.92 (major) and 31.4 (minor).

(*S*)-2-((*S*)-2,2-Dimethyl-1,3-dioxolan-4-yl)octan-1-ol (**4j**). Purification by column chromatography (SiO₂, 10–40% Et₂O/pentane, gradient) afforded **4j** (25 mg, yield = 65%) as a colorless oil: *anti:syn* = >99:1, $[\alpha]_D^{20} = +22.0$ ($c = 1$ in CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 4.10 (dd, $J = 7.9$, 6.1 Hz, 1H), 4.02 (dd, $J = 14.0$, 7.7 Hz, 1H), 3.75 (dd, $J = 11.2$, 3.1 Hz, 1H), 3.69–3.60 (m, 3H), 1.71–1.61 (m, 1H), 1.41 (s, 3H), 1.35 (s, 3H), 1.33–1.18 (m, 10H), 0.87 (t, $J = 6.5$ Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 109.1, 80.0, 68.8, 64.6, 44.0, 31.7, 29.5, 28.1, 27.0, 26.6, 25.5, 22.6, 14.0; HRMS (ESI+, m/z) calcd for C₁₃H₂₆O₃Na [M + Na]⁺ 253.17742, found 253.17761. The diastereomeric ratio was determined by ¹H NMR.

(*S*)-2-((*S*)-2,2-Dimethyl-1,3-dioxolan-4-yl)propan-1-ol (**4k**). Purification by column chromatography (SiO₂, 10–40% Et₂O/pentane, gradient) afforded **4k** (23 mg, yield = 50%) as a colorless oil: *anti:syn* = >99:1, $[\alpha]_D^{20} = +15.2$ ($c = 1$ in CHCl₃) [lit.²⁷ (*anti:syn* = >20:1): $[\alpha]_D^{25} = +15.8$ ($c = 0.88$ in CHCl₃)]; ¹H NMR (400 MHz, CDCl₃) δ 4.10 (dd, $J = 7.9$, 6.1 Hz, 1H), 3.98–3.90 (m, 1H), 3.71–3.56 (m, 3H), 2.77 (br s, 1H), 1.91–1.78 (m, 1H), 1.41 (s, 3H), 1.35 (s, 3H), 0.82 (d, $J = 7.0$ Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 109.4, 80.9, 68.8, 67.5, 39.2, 26.6, 25.7, 13.1; HRMS (ESI+, m/z) calcd for C₈H₁₆O₃Na [M + Na]⁺ 183.09917, found 183.09885. The diastereomeric ratio was determined by ¹H NMR.

(*S*)-1-Hydroxyhexan-2-yl Benzoate (**4l**). Purification by column chromatography (SiO₂, 10–30% Et₂O/pentane, gradient) afforded **4l** (51 mg, yield = 53%) as a colorless oil: 97% ee, $[\alpha]_D^{20} = -23.0$ ($c = 1$ in CHCl₃); [lit.²⁸ (94% ee): $[\alpha]_D^{23} = -24$ ($c = 0.4$ in CHCl₃)]; ¹H NMR (400 MHz, CDCl₃) δ 8.05 (d, $J = 7.3$ Hz, 2H), 7.56 (t, $J = 7.5$ Hz, 1H), 7.43 (t, $J = 7.9$ Hz, 2H), 5.20–5.12 (m, 1H), 3.85–3.71 (m, 2H), 2.29 (br s, 1H), 1.82–1.62 (m, 2H), 1.47–1.29 (m, 4H), 0.90 (t, $J = 7.0$ Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 166.9, 133.1, 130.2, 129.7, 128.4, 76.4, 64.9, 30.4, 27.5, 22.6, 13.9; HRMS (ESI+, m/z) calcd for C₁₃H₁₈O₃Na [M + Na]⁺ 245.11482, found 245.11461. Enantiomeric excess was determined by chiral HPLC analysis, Chiralcel OB-H column, *n*-heptane/*i*-PrOH 95:5, 40 °C, 225 nm, retention times (min): 15.08 (major) and 15.91 (minor).

General Procedure for the Synthesis of γ -Alkyl-Substituted Alcohols through Cu-Catalyzed Asymmetric Allylic Alkylation of Allyl Bromides with Organolithium Reagents Followed by a Hydroboration/Oxidation. To a solution of the alkene (0.2 mmol) in dry THF (1 mL) was added 9-BBN in THF (0.5 M, 0.6 mmol, 3 equiv), and the mixture was stirred at room temperature for 2 h. Then the mixture was cooled to 0 °C, ethanol (2.0 mL), an aq. solution of NaOH (6.0 M, 1.2 mL), and H₂O₂ (30% in water, 4 mL) were added, and the mixture was warmed to room temperature while being stirred overnight. The reaction was quenched with an aqueous solution of Na₂S₂O₃ (5 mL) and the mixture extracted with dichloromethane (3 $\times$ 5 mL). The organic layer was dried with MgSO₄ and filtered and the solvent evaporated in vacuo. The crude product was purified by flash chromatography on silica gel using various mixtures of *n*-pentane/Et₂O as eluent.

(*S*)-3-Phenylheptan-1-ol (**6a**). Purification by column chromatography (SiO₂, 5–20% Et₂O/pentane, gradient) afforded **6a** (29 mg, yield = 67%) as a colorless oil: 99% ee, $[\alpha]_D^{20} = -1.6$ ($c = 1$ in CHCl₃); [lit.²⁹ (94% ee): $[\alpha]_D^{20} = -1.28$ ($c = 1.02$ in CHCl₃)]; ¹H NMR (400 MHz, CDCl₃) δ 7.30–7.05 (m, 5H), 3.52–3.37 (m, 2H), 2.68–2.58 (m, 1H), 1.97–1.85 (m, 1H), 1.83–1.70 (m, 1H), 1.69–

1.41 (m, 4H), 1.29–0.98 (m, 4H), 0.79 (t, $J = 7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 145.2, 128.4, 127.6, 126.1, 61.2, 42.4, 39.6, 36.7, 34.7, 29.7, 27.4, 22.7, 22.6, 14.0; HRMS (APCI+, m/z) calcd for $\text{C}_{13}\text{H}_{19}$ [$\text{M} - \text{H}_2\text{O}$] $^+$ 175.14813, found 175.14825. Enantiomeric excess was determined by chiral HPLC analysis, Chiralcel AS-H column, n -heptane/ i -PrOH 95:5, 40 °C, 210 nm, retention times (min): 8.43 (major) and 8.87 (minor).

(*S*)-3-Phenylnonan-1-ol (**6b**). Purification by column chromatography (SiO_2 , 5–20% Et_2O /pentane, gradient) afforded **6b** (38 mg, yield = 74%) as a colorless oil: 99% ee, $[\alpha]_{\text{D}}^{20} = -4.8$ ($c = 1$ in CHCl_3); [lit.³⁰ (*R*)-enantiomer (68% ee): $[\alpha]_{\text{D}}^{20} = +3.7$ ($c = 0.84$ in CHCl_3)]; ^1H NMR (400 MHz, CDCl_3) δ 7.33–7.08 (m, 5H), 3.55–3.40 (m, 2H), 2.72–2.62 (m, 1H), 1.99–1.87 (m, 1H), 1.87–1.73 (m, 1H), 1.73–1.49 (m, 4H), 1.33–1.02 (m, 10H), 0.79 (t, $J = 7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 145.3, 128.4, 127.6, 126.1, 61.2, 42.5, 39.6, 37.0, 34.7, 31.7, 29.3, 27.4, 25.2, 22.6, 14.0; HRMS (APCI+, m/z) calcd for $\text{C}_{15}\text{H}_{23}$ [$\text{M} - \text{H}_2\text{O}$] $^+$ 203.17943, found: 203.17960. Enantiomeric excess was determined by chiral HPLC analysis, Chiralcel AS-H column, n -heptane/ i -PrOH 95:5, 40 °C, 210 nm, retention times (min): 8.20 (major) and 8.75 (minor).

(*S*)-3-(4-Bromophenyl)butan-1-ol (**6c**). Purification by column chromatography (SiO_2 , 10–20% Et_2O /pentane, gradient) afforded **6c** (28 mg, yield = 70%) as a colorless oil: 99% ee, $[\alpha]_{\text{D}}^{20} = +21.2$ ($c = 1$ in CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.41 (d, $J = 8.4$ Hz, 2H), 7.07 (d, $J = 8.5$ Hz, 2H), 3.65–3.41 (m, 2H), 2.97–2.76 (m, 1H), 1.95–1.68 (m, 2H), 1.24 (d, $J = 7.0$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 145.8, 131.5, 128.7, 119.7, 60.9, 40.7, 35.8, 22.2; HRMS (APCI+, m/z): calculated for $\text{C}_{10}\text{H}_{12}\text{Br}$ [$\text{M} - \text{H}_2\text{O}$] $^+$ 211.01169, found 211.01177. Enantiomeric excess was determined by chiral HPLC analysis, Chiralcel OJ-H column, n -heptane/ i -PrOH 95:5, 40 °C, 224 nm, retention times (min): 15.19 (major) and 15.96 (minor).

(*S*)-3-(4-Bromophenyl)nonan-1-ol (**6d**). Purification by column chromatography (SiO_2 , 10–20% Et_2O /pentane, gradient) afforded **6d** (48 mg, yield = 90%) as a colorless oil: 99% ee, $[\alpha]_{\text{D}}^{20} = -4.6$ ($c = 1$ in CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.40 (d, $J = 8.4$ Hz, 2H), 7.03 (d, $J = 8.4$ Hz, 2H), 3.54–3.46 (m, 1H), 3.45–3.37 (m, 1H), 2.70–2.60 (m, 1H), 1.96–1.86 (m, 1H), 1.78–1.66 (m, 1H), 1.66–1.48 (m, 2H), 1.43 (br s, 1H), 1.34–0.98 (m, 10H), 0.84 (t, $J = 6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 144.3, 131.4, 129.4, 119.7, 60.9, 41.8, 39.5, 36.8, 34.7, 31.7, 29.3, 27.4, 22.6, 15.3, 14.1; HRMS (APCI+, m/z) calcd for $\text{C}_{15}\text{H}_{24}\text{BrO}$ [$\text{M} + \text{H}$] $^+$ 299.1005, found 299.0996. Enantiomeric excess was determined by chiral HPLC analysis, Chiralcel OD-H column, n -heptane/ i -PrOH 95:5, 40 °C, 226 nm, retention times (min): 12.56 (major) and 13.08 (minor).

(*S*)-4-(Benzyloxy)-3-methylbutan-1-ol (**6e**). Purification by column chromatography (SiO_2 , 5–20% Et_2O /pentane, gradient) afforded **6e** (26 mg, yield = 74%) as a colorless oil: 81% ee, $[\alpha]_{\text{D}}^{20} = -3.2$ ($c = 1$ in CHCl_3); [lit.¹⁵ (92% ee): $[\alpha]_{\text{D}}^{20} = -5.5$ ($c = 2.7$ in CHCl_3)]; ^1H NMR (400 MHz, CDCl_3) δ 7.39–7.25 (m, 5H), 4.52 (s, 2H), 3.75–3.59 (m, 2H), 3.34 (ddd, $J = 16.4$, 9.1, 6.2 Hz, 2H), 2.38 (br s, 1H), 2.01–1.88 (m, 1H), 1.69–1.48 (m, 2H), 0.94 (d, $J = 6.9$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 138.0, 128.4, 127.7, 127.7, 76.1, 73.2, 61.1, 38.0, 31.4, 17.7; HRMS (ESI+, m/z) calcd for $\text{C}_{12}\text{H}_{18}\text{O}_2\text{Na}$ [$\text{M} + \text{Na}$] $^+$ 217.11990, found 217.12006. Enantiomeric excess was determined by chiral HPLC analysis, Chiralcel OD-H column, n -heptane/ i -PrOH 99:1, 40 °C, 213 nm, retention times (min): 55.99 (major) and 62.79 (minor).

(*S*)-3-((Benzyloxy)methyl)heptan-1-ol (**6f**). Purification by column chromatography (SiO_2 , 5–30% Et_2O /pentane, gradient) afforded **6f** (26 mg, yield = 93%) as a colorless oil: 88% ee, $[\alpha]_{\text{D}}^{20} = -7.6$ ($c = 1$ in CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.39–7.22 (m, 5H), 4.52 (s, 2H), 3.76–3.55 (m, 2H), 3.47 (dd, $J = 9.1$, 3.9 Hz, 1H), 3.34 (dd, $J = 9.0$, 7.5 Hz, 1H), 2.47 (br s, 1H), 1.83–1.64 (m, 2H), 1.63–1.49 (m, 1H), 1.40–1.12 (m, 6H), 0.88 (t, $J = 6.9$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 137.9, 128.4, 127.7, 74.3, 73.3, 61.2, 36.6, 36.4, 31.8, 29.2, 22.9, 14.1; HRMS (ESI+, m/z) calcd for $\text{C}_{15}\text{H}_{24}\text{O}_2\text{Na}$ [$\text{M} + \text{Na}$] $^+$ 259.16685, found 259.16709. Enantiomeric excess was determined by chiral HPLC analysis, Chiralcel OD-H column, n -heptane/ i -PrOH 99:1, 40 °C, 212 nm, retention times (min): 41.12 (major) and 44.90 (minor).

General procedure for the One-Pot Synthesis of α -Alkyl-Substituted Aldehydes through Cu-Catalyzed Asymmetric Allylic Alkylation of Allyl Bromides with Organolithium Reagents Followed by Ozonolysis. A Schlenk tube equipped with septum and stirring bar was charged with $\text{CuBr}\cdot\text{SMe}_2$ (0.01 mmol, 2.06 mg, 5 mol %) and the appropriate ligand (0.012 mmol, 6 mol %). Dry dichloromethane (2 mL) was added, and the solution was stirred under nitrogen at room temperature for 15 min. Then, allyl bromide **1** (0.2 mmol) was added, and the resulting solution was cooled to -80 °C. In a separate Schlenk flask, the corresponding organolithium reagent (0.24 mmol, 1.2 equiv) was diluted with hexane (toluene in the case of MeLi , combined volume of 1 mL) under nitrogen, and the solution was added dropwise to the reaction mixture over 2 h using a syringe pump. Once the addition was complete, the mixture was stirred for another 2 h at -80 °C. The reaction was quenched with EtOH (2 mL) and ozone was bubbled for 20 min through the reaction mixture. After being stirred for 20 min (solution stays blue), the reaction mixture was purged with nitrogen (disappearance of blue color). Dimethyl sulfide (0.08 mL, 1 mmol, 5 equiv) was added and the mixture was allowed to warm to room temperature while stirred overnight. The reaction mixture was concentrated in vacuo. The crude product was purified by flash chromatography on silica gel using different mixtures of n -pentane/ Et_2O as eluent.

Note: The $\text{S}_{\text{N}}2$: $\text{S}_{\text{N}}2'$ ratio was determined by GC–MS analysis on a sample obtained after quenching with EtOH, which was passed through a short plug of silica gel to remove transition metal residues.

(*R*)-2-((*S*)-2,2-Dimethyl-1,3-dioxolan-4-yl)octanal (**7**). Purification by column chromatography (SiO_2 , 5–20% Et_2O /pentane, gradient) afforded **7** (28 mg, yield = 70%) as a colorless oil: *anti:syn* = >99:1, $[\alpha]_{\text{D}}^{20} = -3.4$ ($c = 1$ in CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 9.70 (d, $J = 3.5$ Hz, 1H), 4.28 (dd, $J = 13.5$, 6.6 Hz, 1H), 4.10 (dd, $J = 8.4$, 6.2 Hz, 1H), 3.69 (dd, $J = 8.3$, 6.8 Hz, 1H), 2.46–2.36 (m, 1H), 1.80–1.64 (m, 2H), 1.40 (s, 3H), 1.34 (s, 3H), 1.32–1.20 (m, 8H), 0.87 (t, $J = 6.6$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 203.4, 109.5, 75.5, 67.7, 55.3, 31.5, 29.3, 27.1, 26.5, 26.3, 25.3, 22.5, 14.0; HRMS (ESI+, m/z) calcd for $\text{C}_{13}\text{H}_{24}\text{O}_3\text{Na}$ [$\text{M} + \text{Na}$] $^+$ 251.16177, found 251.16153. The diastereomeric ratio was determined by ^1H NMR.

General Procedure for the Synthesis of β -Alkyl-Substituted Aldehydes through the Oxidation of γ -Alkyl-Substituted Primary Alcohols with Dess–Martin Periodinane. To a stirred suspension of Dess–Martin periodinane (99 mg, 0.24 mmol, 1.8 equiv) and NaHCO_3 (33 mg, 0.39 mmol, 3 equiv) in dichloromethane (2 mL) at 0 °C was added dropwise a solution of the alcohol (25 mg, 0.13 mmol, 1 equiv) in 1 mL of dichloromethane. The reaction mixture was stirred at 0 °C for 2 h, silica gel was added, and the solvents were removed in vacuo. The crude mixture was purified by flash chromatography on silica gel using different mixtures of n -pentane/ Et_2O as eluent.

(*S*)-4-(Benzyloxy)-3-methylbutanal (**8**). Purification by column chromatography (SiO_2 , 10–40% Et_2O /pentane, gradient) afforded **8** (18 mg, yield = 75%) as a colorless oil: 81% ee, $[\alpha]_{\text{D}}^{20} = -10.6$ ($c = 1$ in CHCl_3); [lit.³¹ (100% ee): $[\alpha]_{\text{D}}^{20} = -11.2$ ($c = 1.7$ in CHCl_3)]; ^1H NMR (400 MHz, CDCl_3) δ 9.76 (t, $J = 2.2$ Hz, 1H), 7.50–7.12 (m, 5H), 4.50 (s, 2H), 3.42 (dd, $J = 9.1$, 5.2 Hz, 1H), 3.25 (dd, $J = 9.1$, 7.7 Hz, 1H), 2.56 (ddd, $J = 16.1$, 6.3, 2.3 Hz, 1H), 2.50–2.35 (m, 1H), 2.28 (ddd, $J = 16.1$, 6.9, 2.1 Hz, 1H), 0.99 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 138.0, 128.4, 127.7, 127.7, 76.1, 73.2, 61.1, 38.0, 31.4, 17.7; HRMS (ESI+, m/z) calcd for $\text{C}_{12}\text{H}_{16}\text{O}_2\text{Na}$ [$\text{M} + \text{Na}$] $^+$ 215.10425, found 215.10405.

■ ASSOCIATED CONTENT

Supporting Information

General procedure for the synthesis of benzoate ester of the alcohols **4h** and **4i**. ^1H and ^{13}C NMR spectra and HPLC analysis for new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: b.l.feringa@rug.nl, m.fananas.mastral@rug.nl.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Financial support from The Netherlands Organization for Scientific Research (NWO-CW, NSF program), the National Research School of Catalysis (NRSC-C), and the Ministry of Education, Culture and Science (Gravity Program 024.001.035) is gratefully acknowledged. We thank M. Smith (GC and HPLC) and T. D. Tiemersma-Wegman (HRMS) (University of Groningen) for technical assistance.

■ REFERENCES

- (1) Pérez, M.; Fañanás-Mastral, M.; Hornillos, V.; Rudolph, A.; Bos, P. H.; Harutyunyan, S. R.; Feringa, B. L. *Chem.—Eur. J.* **2012**, *18*, 11883.
- (2) Burk, M. J.; Ramsden, J. A. *J. Org. Chem.* **2003**, *68*, 5731.
- (3) Francais, A.; Leyva, A.; Etxebarria-Jardi, G.; Ley, S. V. *Org. Lett.* **2010**, *12*, 340.
- (4) Gualandi, A.; Emer, E.; Capdevila, M. G.; Cozzi, P. G. *Angew. Chem., Int. Ed.* **2011**, *50*, 7842.
- (5) Huo, S.; Negishi, E. *Org. Lett.* **2001**, *3*, 3253.
- (6) Nicolaou, K. C.; Ninkovic, S.; Sarabia, F.; Vourloumis, D.; He, Y.; Vallberg, H.; Finlay, M. R. V.; Yang, Z. *J. Am. Chem. Soc.* **1997**, *119*, 7974.
- (7) (a) Imogai, H.; LarchevêQue, M. *Tetrahedron: Asymmetry* **1997**, *8*, 965. (b) Naemura, K.; Murata, M.; Tanaka, R.; Yano, M.; Hirose, K.; Tobe, Y. *Tetrahedron: Asymmetry* **1996**, *7*, 3285.
- (8) (a) Kondakov, D. Y.; Negishi, E. *J. Am. Chem. Soc.* **1995**, *117*, 10771. (b) Kondakov, D. Y.; Negishi, E. *J. Am. Chem. Soc.* **1996**, *118*, 1577. (c) Liang, B.; Novak, T.; Tan, Z.; Negishi, E. *J. Am. Chem. Soc.* **2006**, *128*, 2770.
- (9) For some recent reviews on Cu-catalyzed AAA, see: (a) Harutyunyan, S. R.; den Hartog, T.; Geurts, K.; Minnaard, A. J.; Feringa, B. L. *Chem. Rev.* **2008**, *108*, 2824. (b) Alexakis, A.; Bäckvall, J. E.; Krause, N.; Pàmies, O.; Diéguez, M. *Chem. Rev.* **2008**, *108*, 2796. (c) Langlois, J.-B.; Alexakis, A. In *Transition Metal Catalyzed Allylic Substitution in Organic Synthesis*; Kazmaier, U., Ed.; Springer-Verlag: Berlin, 2012; pp 235–268.
- (10) (a) Tsuji, J. In *Palladium Reagents and Catalysts*; John Wiley & Sons: New York, 1995; p 290. (b) Trost, B. M.; Crawley, M. L. *Chem. Rev.* **2003**, *103*, 2921. (c) Trost, B. M. *J. Org. Chem.* **2004**, *69*, 5813. (d) Pfaltz, A.; Drury, W. J. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 5723. (e) Trost, B. M. *Org. Process Res. Dev.* **2012**, *16*, 185.
- (11) van Klaveren, M.; Persson, E. S. M.; del Villar, A.; Grove, D. M.; Bäckvall, J.-E.; van Koten, G. *Tetrahedron Lett.* **1995**, *36*, 3059.
- (12) For selected examples, see: (a) Tissot-Croset, K.; Alexakis, A. *Tetrahedron Lett.* **2004**, *45*, 7375. (b) Gillingham, D.; Hoveyda, A. H. *Angew. Chem., Int. Ed.* **2007**, *46*, 3860. (c) Mao, B.; Geurts, K.; Fañanás-Mastral, M.; van Zijl, A. W.; Fletcher, S. P.; Minnaard, A. J.; Feringa, B. L. *Org. Lett.* **2011**, *13*, 948. (d) Geerdink, D.; ter Horst, B.; Lepore, M.; Mori, L.; Puzo, G.; Hirsch, A. K. H.; Gilleron, M.; de Libero, G.; Minnaard, A. J. *Chem. Sci.* **2013**, *4*, 709.
- (13) Pérez, M.; Fañanás-Mastral, M.; Bos, P. H.; Rudolph, A.; Harutyunyan, S. R.; Feringa, B. L. *Nat. Chem.* **2011**, *3*, 377.
- (14) Fañanás-Mastral, M.; Pérez, M.; Bos, P. H.; Rudolph, A.; Harutyunyan, S. R.; Feringa, B. L. *Angew. Chem., Int. Ed.* **2012**, *51*, 1922.
- (15) For the construction of optically active bifunctional building blocks through Cu-catalyzed AAA with Grignard reagents, see: van Zijl, A. W.; López, F.; Minnaard, A. J.; Feringa, B. L. *J. Org. Chem.* **2007**, *72*, 2558.
- (16) Compound **4a** was obtained together with 10% of benzyl alcohol derived from the ozonolysis of S_N2 product **3a** as an inseparable mixture.
- (17) Tissot-Croset, K.; Polet, D.; Alexakis, A. *Angew. Chem., Int. Ed.* **2004**, *43*, 2426.
- (18) Chorley, D. F.; Chen, J. L.; Furkert, D. P.; Sperry, J.; Brimble, M. A. *Synlett* **2012**, 23, 128.
- (19) Hammond, M. L.; Zambias, R. A.; Chang, M. A.; Jensen, N. P.; McDonald, J.; Thompson, K.; Boulton, D. A.; Kopka, I. E.; Hand, K. M.; Opas, E. E.; Bach, T.; Davies, P.; MacIntyre, D. E.; Bonney, R. J.; Humes, J. L. *J. Med. Chem.* **1990**, *33*, 908.
- (20) (a) Marshall, J. A.; Trometer, J. D.; Cleary, D. G. *Tetrahedron* **1989**, *45*, 391. (b) Ellwood, A. R.; Mortimer, A. J. P.; Tocher, D. A.; Porter, M. J. *Synlett* **2008**, 2199.
- (21) Lombardi, M.; Morganti, S.; Trombini, C. J. *J. Org. Chem.* **2003**, *68*, 997.
- (22) Ireland, T.; Grossheimann, G.; Wieser-Jeunesse, C.; Knochel, P. *Angew. Chem., Int. Ed.* **1999**, *38*, 3212.
- (23) Tissot-Croset, K.; Polet, D.; Gille, S.; Hawner, C.; Alexakis, A. *Synthesis* **2004**, 2586.
- (24) Huang, Z.; Tan, Z.; Novak, T.; Zhu, G.; Negishi, E. *Adv. Synth. Catal.* **2007**, *349*, 539.
- (25) Imogai, H.; LarchevêQue, M. *Tetrahedron: Asymmetry* **1997**, *8*, 965.
- (26) Allen, A. E.; MacMillan, D. V. C. *J. Am. Chem. Soc.* **2011**, *133*, 4260.
- (27) Shinoyama, M.; Shirokawa, S.; Nakazaki, A.; Kobayashi, S. *Org. Lett.* **2009**, *11*, 1277.
- (28) Kano, T.; Mii, H.; Maruoka, K. *J. Am. Chem. Soc.* **2009**, *131*, 3450.
- (29) Ito, H.; Nagahara, T.; Ishihara, K.; Saito, S.; Yamamoto, H. *Angew. Chem., Int. Ed.* **2004**, *43*, 994.
- (30) Shintani, R.; Takatsu, K.; Takeda, M.; Hayashi, T. *Angew. Chem., Int. Ed.* **2011**, *50*, 8656.
- (31) Wroblewski, A.; Sahasrabudhe, K.; Aubé, J. *J. Am. Chem. Soc.* **2004**, *126*, 5475.