University of Groningen

Access to chiral α -bromo and α -H-substituted tertiary allylic alcohols via copper(I) catalyzed 1,2-addition of Grignard reagents to enones

Madduri , Ashoka V.R.; Minnaard, Adriaan J.; Harutyunyan, Syuzanna R.

Published in:
Organic & Biomolecular Chemistry

DOI:
[10.1002/chin.201234057](https://doi.org/10.1002/chin.201234057)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2012

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Madduri , A. V. R., Minnaard, A. J., & Harutyunyan, S. R. (2012). Access to chiral α -bromo and α -H-substituted tertiary allylic alcohols via copper(I) catalyzed 1,2-addition of Grignard reagents to enones. *Organic & Biomolecular Chemistry*, 10(14), 2878-2884. <https://doi.org/10.1002/chin.201234057>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

SUPPORTING INFORMATION I

Access to Chiral α -Bromo and α -H-Substituted Tertiary Allylic Alcohols via Copper(I) Catalyzed 1,2-Addition of Grignard Reagents to Enones

Ashoka V. R. Madduri, Adriaan J. Minnaard,* Syuzanna R. Harutyunyan*

Stratingh Institute for Chemistry, University of Groningen, Nijenborgh 4, 9747 AG, Groningen, The Netherlands

General Information

Flash chromatography: Merck silica gel type 9385 230-400 mesh, TLC: Merck silica gel 60, 0.25 mm. Components were visualized by UV and Seebach's reagent, a mixture of phosphomolybdic acid (25 g), cerium (IV) sulfate (7.5 g), H₂O (500 mL) and H₂SO₄ (25 mL) or potassium permanganate staining. Progress and conversion of the reaction were determined by GC-MS (GC, HP6890; MS HP5973) with an HP1 or HP5 column (Agilent Technologies, Palo Alto, CA). High resolution mass spectra (HRMS) were recorded on a AEI-MS-902 and FTMS orbitrap (Thermo Fisher Scientific) mass spectrometer. ¹H- and ¹³C-NMR were recorded on a Varian AMX400 (400 and 100.59 MHz, respectively) or a Varian Gemini 200, using CDCl₃ as solvent. Chemical shift values are reported in ppm with the solvent resonance as the internal standard (CHCl₃: δ 7.26 for ¹H, δ 77.0 for ¹³C). Data are reported as follows: chemical shifts, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, br = broad, m = multiplet), coupling constants (Hz), and integration. Carbon assignments are based on APT ¹³C-NMR experiments. Optical rotations were measured on a *Schmidt + Haensch* polarimeter (Polartronic MH8) with a 10 cm cell (*c* given in g/100 mL). Enantiomeric excesses were determined by HPLC analysis using a Shimadzu LC-10ADVP HPLC equipped with a Shimadzu SPD-M10AVP diode array detector or by capillary GC analysis (HP 6890, CP-Chiralsil-Dex-CB column (25 m x 0.25 mm) or ChiralDEX B-PM (30 m x 0.25 mm x 0.25 μ m)) using a flame ionization detector.

All reactions were carried out under a nitrogen atmosphere using oven dried glassware and using standard Schlenk techniques. *t*BuOMe and dichloromethane were dried and distilled from calcium hydride; toluene, THF and *n*-hexane were dried and distilled from sodium. All copper salts were

purchased from Aldrich, and used without further purification. Starting materials were prepared following literature procedures.^{1,2} Grignard reagents were purchased from Aldrich (*i*BuMgBr (2 M in Et₂O), EtMgBr (3 M in Et₂O). Ligands **L1-L6** were purchased from Aldrich. Racemic products were synthesized by reaction of the α,β -unsaturated ketones (**1-3**) and the corresponding Grignard reagent at rt in Et₂O. All Grignard reagents were prepared from the corresponding alkyl bromides and Mg activated with I₂ in Et₂O. Absolute configurations of products were not determined.

Formation of a single copper complex:

The copper based catalyst employed in the reactions was normally prepared in situ by mixing **L5** with CuBr·SMe₂ in 1.1:1 ratio. Formation of a single copper complex was anticipated upon mixing both reagents. This was supported by both ¹H and ³¹P NMR spectroscopy upon mixing **L5** and CuBr·SMe₂ in 1:1 ratio. NMR spectra showed complete disappearance of the signals corresponding to the free ligand **L5** and appearance of a new set of signals corresponding to the copper complex of **L5**.

NMR spectra of all new compounds

¹H NMR

¹ Lu, S-M., Bolm, C. *Angew. Chem., Int. Ed.* **2008**, 47, 8920.

² Moser, R., Boskovic, Z. V., Crowe, C. S., Lipshutz, B. H. *J. Am. Chem Soc.* **2010**, 132, 7852.

^{13}C NMR
APT
HMQC

HMQC (Heteronuclear Multiple-Quantum Coherence) :

a and **b** are the signals observed for the protons attached to that particular carbon.

^1H NMR & ^{31}P NMR spectrum of ligand L1 (CDCl_3 , 500MHz):

^1H NMR & ^{31}P NMR spectrum of $\text{CuBr}\cdot\text{SMe}_2$ complex of L1 (CDCl_3 , 500MHz):