

## University of Groningen

### **Five-year follow-up of oral functioning and quality of life in patients with oral cancer with implant-retained mandibular overdentures**

Korfage, A.; Schoen, P.J.; Raghoobar, G.M.; Bouma, J.; Burlage, F.R.; Roodenburg, J.L.; Vissink, A.; Reintsema, H.

*Published in:*

Head and Neck: Journal of the Sciences and Specialties of the Head and Neck

*DOI:*

[10.1002/hed.21544](https://doi.org/10.1002/hed.21544)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2011

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Korfage, A., Schoen, P. J., Raghoobar, G. M., Bouma, J., Burlage, F. R., Roodenburg, J. L., Vissink, A., & Reintsema, H. (2011). Five-year follow-up of oral functioning and quality of life in patients with oral cancer with implant-retained mandibular overdentures. *Head and Neck: Journal of the Sciences and Specialties of the Head and Neck*, 33(6), 831-839. <https://doi.org/10.1002/hed.21544>

#### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

#### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# FIVE-YEAR FOLLOW-UP OF ORAL FUNCTIONING AND QUALITY OF LIFE IN PATIENTS WITH ORAL CANCER WITH IMPLANT-RETAINED MANDIBULAR OVERDENTURES

Anke Korfage, DDS,<sup>1</sup> Pieter J. Schoen, MD, DDS, PhD,<sup>2</sup> Gerry M. Raghoobar, MD, DDS, PhD,<sup>1</sup> Jelte Bouma, PhD,<sup>3</sup> Fred R. Burlage, MD, PhD,<sup>4</sup> Jan L. N. Roodenburg, DDS, PhD,<sup>1</sup> Arjan Vissink, MD, DDS, PhD,<sup>1</sup> Harry Reintsema, DDS, PhD<sup>1</sup>

<sup>1</sup>Department of Oral and Maxillofacial Surgery and Maxillofacial Prosthetics, University of Groningen and University Medical Center Groningen, Groningen, The Netherlands. E-mail: a.korfage@kchir.umcg.nl

<sup>2</sup>Department of Oral and Maxillofacial Surgery, Admiraal de Ruyter Hospital, Goes, The Netherlands

<sup>3</sup>Graduate School for Health Research (SHARE), University of Groningen and University Medical Center Groningen, Groningen, The Netherlands

<sup>4</sup>Department of Radiotherapy, University of Groningen and University Medical Center Groningen, Groningen, The Netherlands

Accepted 4 June 2010

Published online 9 December 2010 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/hed.21544

**Abstract:** *Background.* The purpose of this prospective study was to assess the quality of life (QOL) and oral functioning of patients with oral cancer up to 5 years after prosthodontic rehabilitation with mandibular implant-retained overdentures.

*Methods.* Fifty patients who had received implants during ablative surgery were evaluated by standardized questionnaires before and after oncological and prosthetic treatment.

*Results.* In 20 of 24 surviving patients, the dentures were functional after 5 years. In these survivors, oral function remained unchanged during this period. In the 6 patients with concurrent comorbidity, global health and QOL had deteriorated, while in the patients without comorbidity, global health and QOL were very high. Five-year survivors had a higher global health and better oral functioning at the 1-year evaluation than nonsurvivors.

*Conclusion.* Oral function and denture satisfaction were high and did not change over time for survivors. Deterioration in overall global health and QOL was associated with concurrent comorbidity. © 2010 Wiley Periodicals, Inc. *Head Neck* 33: 831–839, 2011

**Keywords:** head and neck cancer; edentulous; dental implants; dental prosthesis; quality of life

**P**rosthodontic rehabilitation in patients with oral cancer is challenging as oral functioning is hampered due to the surgical treatment and the subsequent radiotherapy. As a consequence of this combined treatment, wearing a mandibular prosthesis is severely impeded due to the changed anatomic conditions and the intolerance of the denture-bearing mucosa to mechanical loading.<sup>1–4</sup> A solution for this problem might be to provide the patients with an implant-retained mandibular overdenture.

Implant survival in irradiated mandibles, although generally lower than in healthy patients, has still been shown to be relatively high in most articles shown in the literature, and patients have reported an improved level of oral functioning when provided with such a denture.<sup>5–13</sup> Also, assessment of the effect of such a treatment on the patients' functioning and overall quality of life (QOL) is of the utmost importance.<sup>14–20</sup>

In healthy subjects, no clinically relevant changes in oral functioning and patient satisfaction are to be expected after the first year of prosthodontic rehabilitation with an implant-retained overdenture.<sup>21,22</sup> In patients with oral cancer, it is questionable whether this is also applicable, or whether the remaining side effects of the oncological treatment and the impact of having had cancer are more prominent and veil the beneficial effect of an adequate prosthodontic rehabilitation on oral function and QOL. Thus, the purpose of this prospective study was to assess oral functioning and QOL in patients with oral cancer in whom implants had been installed during ablative tumor surgery, up to 5 years after prosthodontic rehabilitation with implant-retained mandibular overdentures.

## MATERIALS AND METHODS

**Patients and Treatment.** All consecutive edentulous patients with oral cancer referred to the Head and Neck Oncology group of the University Medical Center Groningen between May 1998 and April 2002 were screened to be included in this study. Inclusion criteria were edentulous upper and lower jaw, history of prosthetic problems related to the lack of stability, and retention of the lower denture or expected denture-related problems after oncology treatment, first

Correspondence to: A. Korfage  
© 2010 Wiley Periodicals, Inc.

malignancy in head and neck region (squamous cell carcinoma of the tongue, floor of mouth, mandibular gingiva, buccal mucosa, or oropharynx), and the need for primary ablative surgery. The patients were screened by an experienced maxillofacial surgeon (G.M.R.) and prosthodontist (H.R.). It was required that little or no improvement was to be expected from making new dentures after oncological treatment. Patients were offered conventional or implant-based treatment. Fifty-three patients fulfilled the inclusion criteria and 50 patients accepted the option of implant installation during ablative surgery. Two patients refused to have implants installed and 1 patient had never worn a prosthesis. Informed consent was provided from all patients before treatment.

Tumor surgery and implant insertion were performed at the University Medical Center Groningen. All implants (3.75 mm Brånemark screw implants with a machined surface, Nobelbiocare, Gothenburg, Sweden) were inserted during the ablative tumor surgery procedure. All implants were placed in the interforaminal region of the native bone of the mandible in a 2-stage surgical procedure. A 3-month osseointegration period before abutment connection was considered in patients not having radiotherapy after tumor surgery (18 patients). If postoperative radiotherapy was scheduled (32 patients), in general, starting within 6 weeks after surgery, the osseointegration time before abutment connection was increased to 9 months after surgery. All patients were treated by 1 maxillofacial surgeon (G.R.) and 1 prosthodontist (H.R.). Details are described in the article by Schoen et al.<sup>10</sup>

**Functional Assessments and Quality of Life.** Preoperatively, on the day of hospital admission ( $T_0$ ), patients were asked to complete questionnaires regarding oral functioning and QOL. The questionnaires were administered by an investigator not involved in treatment of the patients (P.S.). Similar questionnaires and questionnaires regarding denture satisfaction and the impact of denture-related problems on social activities had to be completed 6 weeks ( $T_1$ ), 1 year ( $T_2$ ), and 5 years ( $T_3$ ) after placing the new dentures.

QOL was assessed using the core questionnaire (Quality of Life Questionnaire-Core 30-questions [QLQ-C30]) and head and neck module (Quality of Life Questionnaire-Core 30 Head and Neck 35-questions [QLQ-H&N35]) of the European Organization for Research and Treatment of Cancer (EORTC).<sup>23</sup> Psychological, physical, and social impact of oral disorders was assessed using the Oral Health Impact Profile (OHIP).<sup>24</sup> General QOL was assessed with the Linear Analogue Self Assessment method (LASA; 1-item version).<sup>25</sup> Denture satisfaction was assessed using a validated questionnaire consisting of 8 separate items focusing on the function of upper and lower dentures, and on specific features such as aesthetics, retention, and functional comfort.<sup>26</sup> Overall denture satisfaction was expressed on a 10-point rating scale (0–10); 0

being completely dissatisfied, to 10 being completely satisfied. Subjective chewing ability was assessed using a 9-item questionnaire on which the patient could rate on a 3-point scale their ability to chew different kinds of food.<sup>27</sup> Impact of denture problems on social activities, such as going out, and contacting and visiting people, was assessed with the Groningen Activity Restriction Scale Dentistry.<sup>28</sup>

**Data Analysis.** The obtained data were evaluated using SPSS (version 16.0 for Windows, SPSS, Chicago, IL). Data are shown as means  $\pm$  SD. Changes were stated as significant if  $p < .05$ . When comparing different groups of patients at the same time, the Mann–Whitney test was used. When comparing results within groups at different times, the Wilcoxon signed-rank test was applied.

## RESULTS

**Patients and Implants.** Patient characteristics are presented in Table 1. In total, 50 patients, 35 men and 15 women (mean age,  $61.5 \pm 11.2$  years; range, 41–81 years) were included at  $T_0$ .

In total, 195 implants were placed in the initial group of 50 patients; of them, 18 patients were treated with surgery only (72 implants) and 32 patients were treated with radiotherapy in addition (123 implants). During the 5-year follow-up, a total of 14 implants were lost; 13 implants in 6 patients who received radiotherapy (implant survival rate 89.4%) and 1 implant in a non-irradiated patient (implant survival rate 98.6%).

At  $T_2$ , 1 year after denture placement, 35 overdentures were in function. Twelve patients had died (48 implants), 7 before abutment connection. Two patients had refused abutment connection (6 implants), because they did not want any additional, nontumor-related, surgery; and 1 patient had already lost 3 implants before abutment connection. The results of  $T_2$  have been published previously.<sup>10</sup> At  $T_3$ , 5 years after denture placement, 26 patients were deceased. Another 4 patients who survived  $T_3$  had to be excluded from follow-up due to removal of the superstructures related to local irritation ( $n = 2$ ), loss of 3 implants ( $n = 1$ ), and the impossibility of making a denture after ablation because of derived anatomic limitations ( $n = 1$ ). Of the remaining 20 patients with functional dentures at  $T_3$ , 9 patients were irradiated (45%).

## Quality of Life and Functional Assessments

*European Organization for Research and Treatment of Cancer Quality of Life Questionnaire-Core 30-Questions and Quality of Life Questionnaire-Core 30 Head and Neck 35-Questions.* The results of the EORTC QLQ-C30 and QLQ-H&N35 questionnaires are presented in Table 2. The results of the evaluations after 1 and 5 years are presented for patients who survived  $T_3$  ( $n = 20$ ), divided into irradiated (RTX;  $n = 9$ ) and non-irradiated patients (non-RTX;  $n = 11$ ). Hardly any differences between and within the groups were found. In the total group, the

**Table 1.** Patient characteristics.

| Age at diagnosis, y | Sex | Primary tumor         | Stage | Total intraforaminal dose, Gy | Status        |
|---------------------|-----|-----------------------|-------|-------------------------------|---------------|
| 57                  | F   | Mandibular gingiva    | T4N1  | —                             | 1 (NTR)       |
| 59                  | M   | Floor of mouth        | T4N2b | —                             | 1 (NTR)       |
| 77                  | F   | Tongue                | T3N2b | 64                            | 1 (TR)        |
| 79                  | M   | Floor of mouth        | T4N0  | 60                            | 1 (TR)        |
| 52                  | F   | Tongue/floor of mouth | T2N1  | 64                            | 1 (TR)        |
| 53                  | M   | Floor of mouth        | T4N0  | 65                            | 1 (TR)        |
| 69                  | M   | Oropharynx            | T2N2b | 64                            | 1 (TR)        |
| 81                  | M   | Oropharynx            | T3N1  | 30                            | 2 (NTR)       |
| 52                  | F   | Tongue                | T2N1  | 58                            | 2 (NTR)       |
| 61                  | M   | Mandibular gingiva    | T2N0  | 64                            | 2 (TR)        |
| 81                  | F   | Tongue/floor of mouth | T2N0  | —                             | 2 (TR)        |
| 50                  | M   | Mandibular gingiva    | T4N2b | 61                            | 2 (TR)        |
| 75                  | M   | Tonsil                | T2N0  | —                             | 3 (NTR)       |
| 64                  | M   | Floor of mouth        | T2N2c | 59                            | 3 (NTR)       |
| 59                  | M   | Tonsil                | T3N0  | 60                            | 3 (NTR)       |
| 68                  | F   | Floor of mouth        | T2N0  | —                             | 3 (NTR)       |
| 65                  | M   | Mandibular gingiva    | T2N0  | —                             | 3 (NTR)       |
| 49                  | F   | Base of tongue        | T3N1  | 58                            | 3 (NTR)       |
| 66                  | M   | Mandibular gingiva    | T4N2b | 67                            | 3 (NTR)       |
| 48                  | M   | Floor of mouth        | T4N1  | 55                            | 3 (NTR)       |
| 78                  | F   | Mandibular gingiva    | T1N0  | —                             | 3 (NTR)       |
| 54                  | M   | Mandibular gingiva    | T4N1  | 62                            | 3 (NTR)       |
| 70                  | M   | Mandibular gingiva    | T4N2b | 50                            | 3, 4 (NTR)    |
| 50                  | M   | Floor of mouth        | T2N1  | 65                            | 3 (TR)        |
| 66                  | M   | Mandibular gingiva    | T4N2b | 64                            | 3 (TR)        |
| 59                  | M   | Oropharynx            | T4N2b | 61                            | 3 (TR)        |
| 49                  | F   | Floor of mouth        | T2N0  | 57                            | 4             |
| 76                  | F   | Mandibular gingiva    | T4N0  | 64                            | 4             |
| 49                  | M   | Floor of mouth        | T2N0  | 50                            | 4 (after 1 y) |
| 71                  | M   | Tonsil                | T3N1  | 67                            | 4 (after 1 y) |
| 43                  | M   | Tongue/floor of mouth | T2N0  | —                             | 5             |
| 65                  | M   | Floor of mouth        | T2N1  | 70                            | 5             |
| 43                  | F   | Tongue                | T1N0  | —                             | 5             |
| 55                  | F   | Tongue                | T2N0  | —                             | 5             |
| 77                  | M   | Tongue                | T1N0  | —                             | 5             |
| 56                  | F   | Floor of mouth        | T1N0  | —                             | 5             |
| 41                  | M   | Base of tongue        | T3N0  | 63                            | —             |
| 54                  | M   | Tongue                | T2N1  | 46                            | —             |
| 51                  | F   | Floor of mouth        | T1N0  | 61                            | —             |
| 64                  | M   | Mandibular gingiva    | T4N0  | 62                            | —             |
| 52                  | M   | Oropharynx            | T3N0  | 12                            | —             |
| 65                  | M   | Floor of mouth        | T2N0  | —                             | —             |
| 63                  | F   | Tongue                | T3N2c | 62                            | —             |
| 46                  | M   | Tongue                | T3N0  | 64                            | —             |
| 54                  | M   | Mandibular gingiva    | T1N0  | —                             | —             |
| 69                  | M   | Tongue                | T2N0  | —                             | —             |
| 71                  | M   | Tongue                | T2N0  | —                             | —             |
| 72                  | M   | Tongue                | T2N0  | —                             | —             |
| 66                  | M   | Tongue                | T3N2b | 66                            | —             |
| 80                  | M   | Tongue                | T2N0  | —                             | —             |

Abbreviations: F, female; M, male; NTR, not tumor-related; TR, tumor-related.

Notes: Patient characteristics regarding age, sex, primary tumor, staging, total interforaminal dose of radiotherapy, and status: 1: died in first year, before prosthesis could be made; 2: died in the first year after delivery of prosthesis; 3: died after first year, but before 5-year evaluation; 4: wears no prosthesis; 5: comorbidity notified at T3.

reported global health and general health after 5 years was lower than after 1 year ( $p < .05$ ) and general QOL tended to decrease ( $p = .070$ ). Weight loss had increased in 4 years. In irradiated patients, the mouth opening was reported more restricted and dry mouth was more severe (only significant after 1 year;  $p < .05$ ).

**Comorbidity.** Based on the data in the patients' medical histories, patients were subdivided into 2 groups

based on the comorbidity noticed at T<sub>3</sub> (Table 3). Six patients were identified with comorbidity, including secondary radiotherapy (after T<sub>2</sub>) in the head and neck region ( $n = 2$ ), an established stroke, lung metastases, severe lung emphysema, and a transient ischemic attack (Table 1). When looking into detail in these patients, global health, physical function, fatigue, and dyspnea were significantly worse in these patients with comorbidity. QOL and global health were very high in patients without comorbidity and remained at the same level

**Table 2.** EORTC QLQ-C30 and EORTC QLQ-H&N 35 questionnaires.

|                                      | After 1 y                  |                                 | After 5 y                  |                                 |
|--------------------------------------|----------------------------|---------------------------------|----------------------------|---------------------------------|
|                                      | Irradiated<br><i>n</i> = 9 | Non-irradiated<br><i>n</i> = 11 | Irradiated<br><i>n</i> = 9 | Non-irradiated<br><i>n</i> = 11 |
| <b>EORTC QLQ-C30</b>                 |                            |                                 |                            |                                 |
| Global health status/quality of life | 93.5 ± 8.1                 | 74.2 ± 24.6*                    | 83.3 ± 12.5                | 64.4 ± 30.5 <sup>†</sup>        |
| Physical functioning                 | 85.9 ± 17.5                | 73.3 ± 23.5                     | 88.9 ± 10.0                | 68.5 ± 33.3                     |
| Role functioning                     | 90.7 ± 14.7                | 77.3 ± 31.0                     | 88.9 ± 18.6                | 72.7 ± 38.2                     |
| Emotional functioning                | 94.4 ± 16.7                | 87.9 ± 22.5                     | 91.7 ± 15.0                | 79.5 ± 28.0                     |
| Cognitive functioning                | 90.7 ± 12.1                | 86.4 ± 19.5                     | 88.9 ± 8.3                 | 75.8 ± 27.2                     |
| Social functioning                   | 94.4 ± 11.8                | 86.4 ± 30.6                     | 88.9 ± 16.7                | 83.3 ± 25.8                     |
| Fatigue                              | 13.6 ± 16.5                | 20.2 ± 30.2                     | 12.3 ± 14.1                | 24.2 ± 26.7                     |
| Nausea and vomiting                  | 0.0 ± 0.0                  | 3.0 ± 6.7                       | 5.6 ± 16.7                 | 1.5 ± 5.0                       |
| Pain                                 | 13.0 ± 16.2                | 10.6 ± 25.0                     | 13.0 ± 23.2                | 9.1 ± 15.6                      |
| Dyspnea                              | 0.0 ± 0.0                  | 24.2 ± 36.8                     | 11.1 ± 23.6                | 27.3 ± 46.7                     |
| Insomnia                             | 3.7 ± 11.1                 | 9.1 ± 15.6                      | 3.7 ± 11.1                 | 9.1 ± 15.6                      |
| Appetite loss                        | 0.0 ± 0.0                  | 9.1 ± 30.2                      | 7.4 ± 14.7                 | 16.7 ± 28.3                     |
| Constipation                         | 3.7 ± 11.1                 | 0.0 ± 0.0                       | 3.7 ± 11.1                 | 3.0 ± 10.1                      |
| Diarrhea                             | 0.0 ± 0.0                  | 6.1 ± 13.5                      | 11.1 ± 23.6                | 6.1 ± 13.5                      |
| Financial difficulties               | 14.8 ± 17.6                | 6.1 ± 20.1                      | 22.2 ± 37.3                | 10.0 ± 16.1                     |
| <b>EORTC QLQ-H&amp;N35</b>           |                            |                                 |                            |                                 |
| Pain                                 | 15.7 ± 22.6                | 6.1 ± 9.9                       | 19.4 ± 11.8                | 9.1 ± 17.3                      |
| Swallowing                           | 19.4 ± 15.6                | 6.8 ± 9.0*                      | 12.7 ± 14.2                | 15.8 ± 23.4                     |
| Sensory problems                     | 18.5 ± 17.6                | 15.2 ± 32.0                     | 13.0 ± 23.2                | 22.7 ± 31.0                     |
| Speech problems                      | 13.6 ± 18.2                | 9.1 ± 14.8                      | 18.5 ± 22.9                | 14.1 ± 21.1                     |
| Trouble with social eating           | 22.2 ± 19.5                | 12.1 ± 25.6                     | 21.3 ± 28.0                | 20.0 ± 28.7                     |
| Trouble with social contact          | 4.4 ± 11.1                 | 5.5 ± 12.9                      | 5.2 ± 15.6                 | 4.2 ± 8.0                       |
| Less sexuality                       | 16.7 ± 28.9                | 14.8 ± 32.7                     | 18.8 ± 30.1                | 25.0 ± 34.5                     |
| Teeth                                | 14.8 ± 33.8                | 9.1 ± 15.6                      | 25.9 ± 32.4                | 6.7 ± 21.1                      |
| Opening mouth                        | 44.4 ± 28.9                | 9.1 ± 21.6 <sup>‡</sup>         | 25.9 ± 32.4                | 6.7 ± 14.1                      |
| Dry mouth                            | 55.6 ± 28.9                | 21.2 ± 22.5*                    | 51.9 ± 29.4                | 26.7 ± 34.4                     |
| Sticky saliva                        | 33.3 ± 28.9                | 12.1 ± 16.8                     | 37.0 ± 35.1                | 30.3 ± 34.8                     |
| Coughing                             | 14.8 ± 17.6                | 27.3 ± 25.0                     | 14.8 ± 17.6                | 15.2 ± 22.9                     |
| Felt ill                             | 3.7 ± 11.1                 | 6.1 ± 20.1                      | 14.8 ± 33.8                | 15.2 ± 22.9                     |
| Pain killers                         | 22.2 ± 44.1                | 45.5 ± 52.2                     | 22.2 ± 44.1                | 18.2 ± 40.5                     |
| Nutritional supplements              | 22.2 ± 44.1                | 9.1 ± 30.2                      | 22.2 ± 44.1                | 18.2 ± 40.5                     |
| Feeding tube                         | 0.0 ± 0.0                  | 0.0 ± 0.0                       | 0.0 ± 0.0                  | 18.2 ± 40.5                     |
| Weight loss                          | 0.0 ± 0.0                  | 9.1 ± 30.2                      | 22.2 ± 44.1                | 36.4 ± 50.5                     |
| Weight gain                          | 11.1 ± 33.3                | 27.3 ± 46.7                     | 0.0 ± 0.0                  | 9.1 ± 30.2                      |

Abbreviations: EORTC QLQ-C30, European Organization for Research and Treatment of Cancer Quality of Life Questionnaire-Core 30-questions; EORTC QLQ-H&N35, European Organization for Research and Treatment of Cancer Quality of Life Questionnaire-Core 30 Head and Neck 35-questions.

\*Significant difference between irradiated and non-irradiated patients at the same point in time  $p < .05$ .

<sup>†</sup>Significant difference between 5 years after placement and one year after placement  $p < .05$ .

<sup>‡</sup>Significant difference between irradiated and non-irradiated patients at the same point in time  $p < .01$ .

Notes: Results of the functional scales, symptom scales and single items of the EORTC QLQ-C30 and multi-item scales and single items of the EORTC QLQ-H&N 35 questionnaires for the 5 years surviving patients with a functional implant-retained overdenture, at 1 and 5 years after placement of the dentures (for irradiated and non-irradiated patients). For the 1-year results ( $n = 35$  patients) see Schoen et al.<sup>10</sup>

between T<sub>2</sub> and T<sub>3</sub>. When comparing the T<sub>1</sub>-data and T<sub>2</sub>-data, there was a progressive decrease in general health, global health, and cognitive function over time in patients with comorbidity. A tendency toward a decrease in cognitive function ( $p = .078$ ) and an increase in weight loss ( $p = .083$ ) with time was also seen in patients with comorbidity.

**Radiotherapy.** The global health status in irradiated patients was higher than the non-irradiated patients. However, 5 patients with comorbidity were among the 11 non-irradiated patients, whereas there was only 1 patient with comorbidity among the 9 irradiated patients. When excluding the patients with comorbidity, the differences in the EORTC QLQ-C30 disappeared.

At T<sub>2</sub>, the irradiated patients reported a dryer mouth, less opening of the mouth, and more difficulties with

swallowing in the QLQ-H&N35 questionnaires (Table 2). At T<sub>3</sub>, the differences between irradiated and non-irradiated patients did not reach significance, although trends were seen toward a dryer mouth ( $p = .095$ ) and more pain ( $p = .056$ ) in irradiated patients. When taking comorbidity into account, we saw several differences in the QLQ-H&N35; the irradiated patients reported a dryer mouth, more pain ( $p < .05$ ), less opening of the mouth, more problems in speech, and more problems related to the dentures ( $p = .059$ ).

When comparing the irradiated patients with the non-irradiated patients, over time, global health and global health-related QOL tended to decrease for the irradiated patients ( $p = .059$  and  $p = .066$ ).

**Survivors versus Nonsurvivors.** When looking retrospectively into the 35 patients with functional



**Table 3.** Comorbidity versus no comorbidity.

| EORTC QLQ-C30                        | Comorbidity<br><i>n</i> = 6 | No comorbidity<br><i>n</i> = 14 |
|--------------------------------------|-----------------------------|---------------------------------|
| Global health status/quality of life | 48.6 ± 27.6                 | 83.3 ± 16.3*                    |
| Physical functioning                 | 50.0 ± 35.5                 | 89.5 ± 9.3*                     |
| Role functioning                     | 55.6 ± 44.3                 | 90.5 ± 16.9                     |
| Emotional functioning                | 69.4 ± 33.6                 | 91.7 ± 14.2                     |
| Cognitive functioning                | 63.9 ± 30.6                 | 89.3 ± 10.5                     |
| Social functioning                   | 69.4 ± 28.7                 | 92.9 ± 14.2                     |
| Fatigue                              | 40.7 ± 26.0                 | 9.5 ± 12.2*                     |
| Nausea and vomiting                  | 2.8 ± 6.8                   | 3.6 ± 13.4                      |
| Pain                                 | 16.7 ± 18.3                 | 8.3 ± 19.3                      |
| Dyspnea                              | 61.1 ± 49.1                 | 2.4 ± 8.9†                      |
| Insomnia                             | 11.1 ± 17.2                 | 4.8 ± 12.1                      |
| Appetite loss                        | 33.3 ± 33.3                 | 4.8 ± 12.1                      |
| Constipation                         | 5.6 ± 13.6                  | 2.4 ± 8.9                       |
| Diarrhea                             | 11.1 ± 17.2                 | 7.1 ± 19.3                      |
| Financial difficulties               | 16.7 ± 18.3                 | 15.4 ± 32.2                     |

Abbreviation: EORTC QLQ-C30, European Organization for Research and Treatment of Cancer Quality of Life Questionnaire-Core 30 questions.

\*Significant difference between patients with and without comorbidity after 5 years  $p < .01$ .

†Significant difference between patients with and without comorbidity after 5 years  $p < .05$ .

Notes: Results of the functional scales, symptom scales, and single items of the EORTC QLQ-C30 for patients with and without comorbidity, 5 years after placement of the dentures.

dentures at T<sub>2</sub>, there were some differences between the 5-year survivors with functional dentures ( $n = 20$ ) and those patients who did not make it to T<sub>3</sub> ( $n = 12$ ; the results are not depicted in a table in this article). At T<sub>2</sub>, the 5-year survivors had reported a higher global health and fewer problems with swallowing ( $p < .05$ ) than the nonsurvivors. Nonsurvivors tended to report more pain and a lower general QOL than the survivors ( $p = .068$ ).

**Oral Health Impact Profile, Functional Assessments, Social Restrictions, and Denture Satisfaction.** The OHIP results are presented in Table 4, and the results of questionnaires regarding oral functioning and denture satisfaction are presented in Table 5. Over time, there were no changes in results for the

total group, neither were differences seen between patients with or without comorbidity.

**Radiotherapy.** A tendency toward more pain was reported in the OHIP in the irradiated group ( $p = .067$ ) between T<sub>2</sub> and T<sub>3</sub>. When excluding patients with comorbidity, more differences were found between irradiated patients and non-irradiated patients: at T<sub>3</sub>, irradiated patients reported more functional limitations and physical pain ( $p < .05$ ) than non-irradiated patients, a tendency was seen toward more physical disability ( $p = .081$ ) and a higher score in the handicap domain ( $p = .081$ ) in irradiated patients. Previously, we reported that overall denture satisfaction was higher in non-irradiated patients than in irradiated patients at T<sub>2</sub>,<sup>10</sup> but in the irradiated patients, denture satisfaction was also rather

**Table 4.** OHIP questionnaire.

|                          | After 1 y                  |                                 | After 5 y                  |                                 |
|--------------------------|----------------------------|---------------------------------|----------------------------|---------------------------------|
|                          | Irradiated<br><i>n</i> = 9 | Non-irradiated<br><i>n</i> = 11 | Irradiated<br><i>n</i> = 9 | Non-irradiated<br><i>n</i> = 11 |
| OHIP14                   | 12.4 ± 10.9                | 6.3 ± 8.9*                      | 12.8 ± 12.1                | 6.7 ± 6.5                       |
| Functional limitation    | 12.0 ± 6.5                 | 6.6 ± 5.4†                      | 11.1 ± 5.9                 | 7.3 ± 4.8                       |
| Physical pain            | 7.0 ± 9.5                  | 4.0 ± 6.3                       | 11.0 ± 9.6                 | 4.3 ± 6.1‡                      |
| Physical disability      | 13.0 ± 10.7                | 5.9 ± 8.1§                      | 10.4 ± 10.5                | 6.2 ± 6.3                       |
| Psychological discomfort | 2.1 ± 5.3                  | 0.8 ± 1.9                       | 3.4 ± 5.8                  | 1.0 ± 2.0                       |
| Psychological disability | 2.0 ± 4.3                  | 0.9 ± 2.4                       | 1.9 ± 3.6                  | 0.9 ± 1.0                       |
| Social disability        | 1.3 ± 2.5                  | 0.8 ± 1.8                       | 1.3 ± 2.7                  | 0.9 ± 1.3                       |

Abbreviations: OHIP, Oral Health Impact Profile.

\*Significant difference between irradiated and non-irradiated patients at the same point in time;  $p < .05$ .

†Tendency toward difference between irradiated and non-irradiated patients at the same point in time;  $p = .056$ .

‡Tendency toward difference between irradiated and non-irradiated patients at the same point in time;  $p = .067$ .

§Tendency toward difference between irradiated and non-irradiated patients at the same point in time;  $p = .056$ .

Notes: Results of the Oral Health Impact Profile (OHIP) questionnaire, at 1 and 5 years after placement of the dentures, for irradiated and non-irradiated 5-year survivors with a functional implant-retained overdenture. For the 1-year results ( $n = 35$  patients) see Schoen et al.<sup>10</sup>

**Table 5.** Oral functioning and denture satisfaction.

|                              | After 1 y                  |                                 | After 5 y                  |                                 |
|------------------------------|----------------------------|---------------------------------|----------------------------|---------------------------------|
|                              | Irradiated<br><i>n</i> = 9 | Non-irradiated<br><i>n</i> = 11 | Irradiated<br><i>n</i> = 9 | Non-irradiated<br><i>n</i> = 11 |
| GARS-D                       | 2.6 ± 4.6                  | 1.9 ± 3.9                       | 3.5 ± 5.0                  | 2.8 ± 5.1                       |
| Denture satisfaction         | 12.9 ± 4.8                 | 11.6 ± 4.4                      | 13.9 ± 4.8                 | 11.8 ± 3.1                      |
| Overall denture satisfaction | 8.4 ± 1.2                  | 8.5 ± 1.4                       | 8.5 ± 1.3                  | 8.9 ± 1.1                       |
| Chewing/eating               | 7.4 ± 7.0                  | 3.8 ± 4.3                       | 6.0 ± 6.7                  | 4.6 ± 4.9                       |
| LASA quality of life         | 81.8 ± 18.5                | 69.3 ± 24.9                     | 87.4 ± 9.5                 | 65.3 ± 28.7*                    |

Abbreviations: GARS-D, Groningen Activity Restriction Scale Dentistry; LASA, Linear Analogue Self Assessment.

\*Tendency toward difference between irradiated and non-irradiated patients at the same point in time  $p = .055$ .

Notes: Results of questionnaires regarding oral functioning and denture satisfaction, at 1 and 5 years after placement of the dentures, for irradiated and non-irradiated patients with a functional implant-retained overdenture. For the 1-year results ( $n = 35$  patients) see Schoen et al.<sup>10</sup>

high. On the other scales of functional assessment, the non-irradiated patients showed better results than the irradiated patients at T<sub>2</sub>.<sup>10</sup> At T<sub>3</sub>, denture satisfaction again scored high, but denture satisfaction and functional assessment showed no differences between irradiated and non-irradiated patients. Overall QOL, as measured with the LASA, showed, as did the EORTC QLQ-C30, a higher QOL for the irradiated patients after 5 years ( $p = .055$ ), but this difference disappeared when taking comorbidity into account.

**Survivors versus Nonsurvivors.** At T<sub>1</sub>, nonsurvivors reported to be more concerned with the future functioning of their dentures than the 5-year survivors ( $p < .05$ ). There tended to be more social restrictions and chewing problems ( $p = .095$  and  $p = .074$ ) for the non-survivors than for the survivors. At T<sub>2</sub>, survivors reported less social restrictions than nonsurvivors ( $p = .059$ ). Also, survivors tended to be more satisfied with their dentures than the nonsurvivors ( $p = .087$ ).

## DISCUSSION

The surviving 20 patients with functional dentures did not report a difference in oral function between 1 year and 5 years after prosthetic rehabilitation. The observed deterioration in overall global health and QOL was strongly associated with concurrent comorbidity in 6 patients. For patients without known comorbidity, general QOL and global health were very high.

No difference in oral function was reported at the 1 year and 5 year follow-ups after placement of the prostheses. This observation is comparable to results of studies in healthy subjects.<sup>21,22</sup> The oral function of the patients in this study was reasonable, but lower than in healthy subjects.<sup>27</sup> Still, the denture satisfaction was very high. However, there was a difference in global health, oral and social functioning, and denture satisfaction between the 5-year survivors and the nonsurvivors, indicating a “natural” selection of patients. This is in agreement with the findings of

other studies,<sup>30,31</sup> where high scores of functioning scales and low scores on symptom items at 1-year follow-up seemed to predict a high survival at 5 years. In our study, survivors reported fewer problems with swallowing and less restrictions in social activities. The nonsurvivors were more concerned with the future functioning of their dentures than the survivors. An explanation can be that the 20 patients with a functional denture had a lower percentage of large tumors compared to the nonsurvivors (Table 1), thus needing less extensive surgery with less morbidity. Also, among the deceased and excluded patients at T<sub>3</sub>, a larger percentage had received radiotherapy in comparison to the survivors, probably giving less favorable oral conditions.

The scores of the EORTC QLQ-C30 and QLQ-H&N35 questionnaires at T<sub>3</sub> are comparable to the results of other QOL studies in patients with head and neck cancer.<sup>30–34</sup> The patients without known comorbidity reported high scores comparable to those of healthy subjects. This observation indicates that even after oncological treatment, patients still can reach “normal” health levels. Furthermore, in previous studies, the question was raised whether patients do value oral rehabilitation as essential in their life after head and neck cancer. In one study reporting on general QOL in patients without an implant-borne overdenture, no difference in general QOL was found between the patients who wore their mandibular dentures and patients who did not.<sup>35</sup> A review relating QOL to functional outcome also showed no difference in QOL between patients with a conventional dental/tissue-supported denture, an implant-retained overdenture, and patients without dentures.<sup>17</sup> Most patients reported satisfactory outcomes regardless of the type or presence of prosthetic rehabilitation. This finding is in agreement with the findings of Murphy et al,<sup>18</sup> as data correlated QOL with functional outcome and symptom burden often fails to demonstrate a consistent relationship. The latter authors suggested that this may be attributed to methodological issues in the study design or the patient’s ability to adapt to functional and symptom control problems.

It is obvious that certain stages of disease and cancer treatment will lead to disastrous anatomic or physiological conditions in which oral rehabilitation cannot be restored to a level comparable to the level before onset of the disease. However, the patients' ability to adapt to functional problems and to accept the loss of some oral functions should not be underestimated. Another conclusion could be that validated sensitive instruments to rate the influences of oral rehabilitation on QOL are still not available for general application. Regarding general health-related QOL, such validated instruments are commonly available.<sup>15</sup> However, these general health-related QOL questionnaires seem to lack the discriminating ability to measure the effects of prosthodontic treatment on QOL in patients with oral cancer. Efforts have been made to develop instruments that might solve this problem, such as Liverpool Oral Rehabilitation Questionnaire, which was developed in 2004 and has been used ever since.<sup>36-39</sup> Also, more specific questionnaires that focus on head and neck function, such as speech and swallowing are currently available.<sup>40</sup> Unfortunately, we were not able to use such questionnaires as these questionnaires were not available at the time of inclusion of our patients into our study.

It seems that other factors such as comorbidity are far more important in determining the patients' QOL being an important caution that has to be considered when interpreting the results of the questionnaires regarding general health. With a closer look, the decrease in QOL we observed appeared to be caused by a small group of patients with severe comorbidity. Most patients with comorbidity were not irradiated. When taking this comorbidity into account, the specific head and neck module reveals differences between the irradiated and non-irradiated patients even after 5 years, which can be related to the late effects of the radiotherapy, such as dry mouth, less opening of the mouth, and problems with swallowing and speech. Terrell et al<sup>41</sup> ranked comorbidity to be the second greatest predictor of decreased QOL in patients with head and neck cancer. In our study, we did not apply standard comorbidity measures as the Adult Comorbidity Evaluation-27, that are currently used in studies to code and quantify comorbidity.<sup>42-44</sup> Nevertheless, we were able to indicate that comorbidity apparently played a larger role in decreased QOL scores than radiotherapy. However, 2 patients received radiotherapy between T<sub>2</sub> and T<sub>3</sub> due to recurrent disease. In the analysis, these patients were scored as non-irradiated (intention-to-treat procedure) and were considered as having comorbidity. This could also explain why differences are only found in the head and neck module when excluding patients with comorbidity.

Implant loss was higher in patients who received radiotherapy post-tumor surgery. This is in agree-

ment with other studies.<sup>5,7,8,11</sup> A review reports that the increase in the risk of implant failure in irradiated patients may be up to 12 times greater; however, the magnitude of this difference should be accepted with caution, because studies making these comparisons are of poor to moderate quality.<sup>12</sup> The failure rate of 10.6% in irradiated bone over a period of 5 years found in our study is considered good. However, 26 patients had died and the percentage of patients who had received postoperative radiotherapy decreased over time among the survivors (73% at baseline vs 54% 5 years after placement of the dentures [ $n = 20$ ] and patients that survived). This could have contributed to the relatively low failure rate of implants in irradiated bone.

The percentage of patients rehabilitated with the help of dental implants placed after ablative surgery and postoperative radiotherapy varies in the literature. Reported percentage are between 22% and 91%,<sup>9,14,45-50</sup> depending largely on the type of patients included, the type of reconstruction, the survival rate of patients and implants, and the length of the follow-up. In our study where the implants were placed during ablative surgery, a relatively large number of the living patients were rehabilitated with dentures (at T<sub>2</sub>, 92%; at T<sub>3</sub>, 83%). No delay or complications in oncological treatment were seen due to the placement of the implants at that time. Still, 2 patients refused abutment connection because of the expected extra burden of abutment connection surgery. Also, from previous data, it was concluded that many patients refrain from further surgery, including implant installation, after they survived head and neck oncology treatment, despite an improvement of oral functioning that was to be expected postsurgery.<sup>35</sup> When placing the implants during ablation, a significant time reduction of (pre)prosthetic rehabilitation can be achieved. Consequently, a large percentage of patients and even patients with a worse general prognosis can benefit for some time from the improvements in aesthetics and oral function. Future studies might identify patients who are less likely to benefit from implant placement per ablation. Our study indicates that implant installation during ablative surgery results in a high percentage of rehabilitated patients, also after 5 years. From a health economics point of view, however, the loss of resources needs further consideration by performing a cost-effectiveness analysis.

Based on this study, we conclude that the overall global health and QOL deteriorated in the total group between 1 and 5 years after placement of the dentures, which was due to concurrent comorbidity in a small number of patients. The global health and QOL for patients without comorbidity was very high. A large number of surviving patients could benefit from an implant-retained mandibular overdenture (83%)



after 5 years. The oral function and denture satisfaction was high and did not change over time for the 5-year survivors.

## REFERENCES

- Buchbinder D, Urken ML, Vickery C, Weinberg H, Sheiner A, Biller H. Functional mandibular reconstruction of patients with oral cancer. *Oral Surg Oral Med Oral Pathol* 1989;68(4 Pt 2):499–503; discussion 503–504.
- Hayter JP, Cawood JI. Oral rehabilitation with endosteal implants and free flaps. *Int J Oral Maxillofac Surg* 1996;25:3–12.
- Marker P, Siemssen SJ, Bastholt L. Osseointegrated implants for prosthetic rehabilitation after treatment of cancer of the oral cavity. *Acta Oncol* 1997;36:37–40.
- Misieki DJ, Chang AK. Implant reconstruction following removal of tumors of the head and neck. *Otolaryngol Clin North Am* 1998;31:689–725.
- Granström G. Radiotherapy, osseointegration and hyperbaric oxygen therapy. *Periodontol* 2000 2003;33:145–162.
- Müller F, Schädler M, Wahlmann U, Newton JP. The use of implant-supported prostheses in the functional and psychosocial rehabilitation of tumor patients. *Int J Prosthodont* 2004;17:512–517.
- Granström G. Osseointegration in irradiated cancer patients: an analysis with respect to implant failures. *J Oral Maxillofac Surg* 2005;63:579–585.
- Yerit KC, Posch M, Seemann M, et al. Implant survival in mandibles of irradiated oral cancer patients. *Clin Oral Implants Res* 2006;17:337–344.
- Schoen PJ, Raghoobar GM, Bouma J, et al. Rehabilitation of oral function in head and neck cancer patients after radiotherapy with implant-retained dentures: effects of hyperbaric oxygen therapy. *Oral Oncol* 2007;43:379–388.
- Schoen PJ, Raghoobar GM, Bouma J, et al. Prosthodontic rehabilitation of oral function in head-neck cancer patients with dental implants placed simultaneously during ablative tumour surgery: an assessment of treatment outcomes and quality of life. *Int J Oral Maxillofac Surg* 2008;37:8–16.
- Colella G, Cannavale R, Pentenero M, Gandolfo S. Oral implants in irradiated patients: a systematic review. *Int J Oral Maxillofac Implants* 2007;22:616–622.
- Inde S, Kopp S, Gundlach K, Konstantinović VS. Effects of radiation therapy on craniofacial and dental implants: a review of the literature. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2009;107:56–65.
- Schoen PJ, Reintsema H, Raghoobar GM, Vissink A, Roodenburg JL. The use of implant retained mandibular prostheses in the oral rehabilitation of head and neck cancer patients. A review and rationale for treatment planning. *Oral Oncol* 2004;40:862–871.
- Rogers SN, Panasar J, Pritchard K, Lowe D, Howell R, Cawood JI. Survey of oral rehabilitation in a consecutive series of 130 patients treated by primary resection for oral and oropharyngeal squamous cell carcinoma. *Br J Oral Maxillofac Surg* 2005;43:23–30.
- Rogers SN, Ahad SA, Murphy AP. A structured review and theme analysis of papers published on 'quality of life' in head and neck cancer: 2000–2005. *Oral Oncol* 2007; 43:843–868.
- Rogers SN. Quality of life for head and neck cancer patients—has treatment planning altered? *Oral Oncol* 2009; 45:435–439.
- Tang JA, Rieger JM, Wolfaardt JF. A review of functional outcomes related to prosthetic treatment after maxillary and mandibular reconstruction in patients with head and neck cancer. *Int J Prosthodont* 2008;21:337–354.
- Murphy BA, Ridner S, Wells N, Dietrich M. Quality of life research in head and neck cancer: a review of the current state of the science. *Crit Rev Oncol Hematol* 2007; 62:251–267.
- Pusic A, Liu JC, Chen CM, et al. A systematic review of patient-reported outcome measures in head and neck cancer surgery. *Otolaryngol Head Neck Surg* 2007;136:525–535.
- Mlynarek AM, Rieger JM, Harris JR, et al. Methods of functional outcomes assessment following treatment of oral and oropharyngeal cancer: review of the literature. *J Otolaryngol Head Neck Surg* 2008;37:2–10.
- Meijer HJ, Raghoobar GM, Batenburg RH, Visser A, Vissink A. Mandibular overdentures supported by two or four endosseous implants: a 10-year clinical trial. *Clin Oral Implants Res* 2009;20:722–728.
- Raghoobar GM, Meijer HJ, van't Hof M, Stegenga B, Vissink A. A randomized prospective clinical trial on the effectiveness of three treatment modalities for patients with lower denture problems. A 10 year follow-up study on patient satisfaction. *Int J Oral Maxillofac Surg* 2003;32:498–503.
- Björdal K, Ahlner-Elmqvist M, Tolleson E, et al. Development of a European Organization for Research and Treatment of Cancer (EORTC) questionnaire module to be used in quality of life assessments in head and neck cancer patients. EORTC Quality of Life Study Group. *Acta Oncol* 1994;33:879–885.
- Slade GD, Spencer AJ. Development and evaluation of the Oral Health Impact Profile. *Community Dent Health* 1994; 11:3–11.
- Andrews FM, Withey SB. Social factors of well-being. New York: Plenum Press; 1976.
- Vervoor JM, Duinkerke AS, Luteijn F, van de Poel AC. Assessment of denture satisfaction. *Community Dent Oral Epidemiol* 1988;16:364–367.
- Stellingsma K, Slagter AP, Stegenga B, Raghoobar GM, Meijer HJ. Masticatory function in patients with an extremely resorbed mandible restored with mandibular implant-retained overdentures: comparison of three types of treatment protocols. *J Oral Rehabil* 2005;32:403–410.
- Bouma J, Boerrigter LM, Van Oort RP, van Sonderen E, Boering G. Psychosocial effects of implant-retained overdentures. *Int J Oral Maxillofac Implants* 1997;12:515–522.
- Stellingsma K, Bouma J, Stegenga B, Meijer HJ, Raghoobar GM. Satisfaction and psychosocial aspects of patients with an extremely resorbed mandible treated with implant-retained overdentures. A prospective, comparative study. *Clin Oral Implants Res* 2003;14:166–172.
- Abendstein H, Nordgren M, Boysen M, et al. Quality of life and head and neck cancer: a 5 year prospective study. *Laryngoscope* 2005;115:2183–2192.
- Nordgren M, Hammerlid E, Björdal K, Ahlner-Elmqvist M, Boysen M, Jannert M. Quality of life in oral carcinoma: a 5-year prospective study. *Head Neck* 2008;30:461–470.
- Klug C, Neuburg J, Glaser C, Schwarz B, Kermer C, Millesi W. Quality of life 2–10 years after combined treatment for advanced oral and oropharyngeal cancer. *Int J Oral Maxillofac Surg* 2002;31:664–669.
- Schliephake H, Jamil MU. Prospective evaluation of quality of life after oncologic surgery for oral cancer. *Int J Oral Maxillofac Surg* 2002;31:427–433.
- Infante-Cossio P, Torres-Carranza E, Cayuela A, Hens-Aumente E, Pastor-Gaitan P, Gutierrez-Perez JL. Impact of treatment on quality of life for oral and oropharyngeal carcinoma. *Int J Oral Maxillofac Surg* 2009;38:1052–1058.
- Schoen PJ, Reintsema H, Bouma J, Roodenburg JL, Vissink A, Raghoobar GM. Quality of life related to oral function in edentulous head and neck cancer patients posttreatment. *Int J Prosthodont* 2007;20:469–477.
- Pace-Balzan A, Cawood JI, Howell R, Lowe D, Rogers SN. The Liverpool Oral Rehabilitation Questionnaire: a pilot study. *J Oral Rehabil* 2004;31:609–617.
- Pace-Balzan A, Cawood JI, Howell R, Butterworth CJ, Lowe D, Rogers SN. The further development and validation of the Liverpool Oral Rehabilitation Questionnaire: a cross-sectional survey of patients attending for oral rehabilitation and general dental practice. *Int J Oral Maxillofac Surg* 2006;35:72–78.
- Pace-Balzan A, Butterworth CJ, Dawson LJ, Lowe D, Rogers SN. The further development and validation of the Liverpool Oral Rehabilitation Questionnaire (LORQ) version 3: a cross-sectional survey of patients referred to a dental hospital for removable prostheses replacement. *J Prosthet Dent* 2008;99: 233–242.
- Pace-Balzan A, Butterworth C, Lowe D, Rogers SN. The responsiveness of the Liverpool Oral Rehabilitation Questionnaire (LORQ): a pilot study. *Int J Prosthodont* 2009;22:456–458.
- Kanats AN, Rogers SN. A guide of the questionnaires used in the measurement of health-related quality of life in head and neck oncology. *Tumori* 2008;94:724–731.
- Terrell JE, Ronis DL, Fowler KE, et al. Clinical predictors of quality of life in patients with head and neck cancer. *Arch Otolaryngol Head Neck Surg* 2004;130:401–408.

42. Johnston AS, Piccirillo JF, Creech CM, Littenberg B, Jeffe D, Spitznagel ELJ. Validation of a comorbidity education program. *J Regist Manage* 2001;28:125–131.
43. Piccirillo JF, Costas I. The impact of comorbidity on outcomes. *ORL J Otorhinolaryngol Relat Spec* 2004;66:180–185.
44. Datema FR, Ferrier MB, van der Schroeff MP, Baatenburg de Jong RJ. Impact of comorbidity on short-term mortality and overall survival of head and neck cancer patients. *Head Neck* 2010;32:728–736.
45. Garrett N, Roumanas ED, Blackwell KE, et al. Efficacy of conventional and implant-supported mandibular resection prostheses: study overview and treatment outcomes. *J Prosthet Dent* 2006;96:13–24.
46. Schepers RH, Slagter AP, Kaanders JH, van den Hoogen FJ, Merks MA. Effect of postoperative radiotherapy on the functional result of implants placed during ablative surgery for oral cancer. *Int J Oral Maxillofac Surg* 2006;35:803–808.
47. Nelson K, Heberer S, Glatzer C. Survival analysis and clinical evaluation of implant-retained prostheses in oral cancer resection patients over a mean follow-up period of 10 years. *J Prosthet Dent* 2007;98:405–410.
48. Hundepool AC, Dumans AG, Hofer SO, et al. Rehabilitation after mandibular reconstruction with fibula free-flap: clinical outcome and quality of life assessment. *Int J Oral Maxillofac Surg* 2008;37:1009–1013.
49. Smolka K, Kraehenbuehl M, Eggensperger N, et al. Fibula free flap reconstruction of the mandible in cancer patients: evaluation of a combined surgical and prosthodontic treatment concept. *Oral Oncol* 2008;44: 571–581.
50. Adell R, Svensson B, Bågenholm T. Dental rehabilitation in 101 primarily reconstructed jaws after segmental resections—possibilities and problems. An 18-year study. *J Craniomaxillofac Surg* 2008;36:395–402.