University of Groningen

Placental haemosiderosis

Khong, T. Y.; Toering, T. J.; Erwich, J. J. H. M.

Published in:
Pathology

DOI:
[10.3109/00313025.2010.494298](https://doi.org/10.3109/00313025.2010.494298)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2010

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Khong, T. Y., Toering, T. J., & Erwich, J. J. H. M. (2010). Placental haemosiderosis: authors' reply. *Pathology*, 42(5), 502-502. <https://doi.org/10.3109/00313025.2010.494298>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Placental haemosiderosis: authors' reply

Sir,

We thank Dr Stanek¹ for his interest in our article.² His summary of our findings and conclusion are correct to a point but misleading. We did not comment on the presence of haemosiderin in the chorionic villi, which forms a large part of his letter, not because of his presumption that the iron stains were negative there but because, specifically, we indicated that we were examining the deposition of iron in the chorionic and/or basal plate and the membranes only.

In his first point, we did not say that an approach of routine histochemistry staining for iron is not useful. We did say, however, that it is not a useful marker for chronic placental separation and for adverse neonatal outcome. His statement that he believes that some amount of stainable iron can be seen in about half of placentas does not seem to be evidence-based, which is what we have rectified in our study.

We will not respond to his third point in detail as we did not examine the chorionic villi for iron staining, as noted earlier. However, we will point out that statements such as 'probably less frequently' and 'I regard ...' are without merit unless they can be substantiated by a reference or a body of experiment.

His conclusion that choriodecidual haemosiderin may not have a substantial clinical significance is correct but not because it is performed routinely. It is precisely that they were not performed routinely but in selected cases that clinical associations with choriodecidual haemosiderosis were found previously that we have refuted in this study.

T. Y. Khong*

T. J. Toering*

J. J. H. M. Erwich†

**SA Pathology, Women's and Children's Hospital, Adelaide, South Australia; †Department of Obstetrics and Gynecology, University Medical Centre Groningen, The Netherlands*

Contact Professor T. Y. Khong.

E-mail: yee.khong@adelaide.edu.au

1. Stanek J. Placental haemosiderosis. *Pathology* 2010; 42: 499–501.

2. Khong TY, Toering TJ, Erwich JJHM. Haemosiderosis in the placenta does not appear to be related to chronic placental separation or adverse neonatal outcome. *Pathology* 2010; 42: 119–24.

DOI: 10.3109/00313025.2010.494298