University of Groningen

Risk and prognostic factors for non-specific musculoskeletal pain

Lakke, Sandra E.; Soer, Remko; Takken, Tim; Reneman, Michiel F.

Published in:
Pain

DOI:
[10.1016/j.pain.2009.08.032](https://doi.org/10.1016/j.pain.2009.08.032)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2009

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Lakke, S. E., Soer, R., Takken, T., & Reneman, M. F. (2009). Risk and prognostic factors for non-specific musculoskeletal pain: A synthesis of evidence from systematic reviews classified into ICF dimensions. *Pain*, 147(1-3), 153-164. <https://doi.org/10.1016/j.pain.2009.08.032>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Risk and prognostic factors for non-specific musculoskeletal pain: A synthesis of evidence from systematic reviews classified into ICF dimensions

Sandra E. Lakke^{a,b,c,*}, Remko Soer^{b,c}, Tim Takken^d, Michiel F. Reneman^{b,c}

^a Center for Applied Research and Innovation in Health Care and in Nursing, Hanze University Groningen, University of Applied Sciences, The Netherlands

^b Center for Rehabilitation, University Medical Center Groningen, Groningen, The Netherlands

^c Graduate School for Health Care Studies, University Medical Center Groningen, Groningen, The Netherlands

^d Health Sciences, Department of Physiotherapy Research, Utrecht University, The Netherlands

ARTICLE INFO

Article history:

Received 11 November 2008

Received in revised form 17 August 2009

Accepted 31 August 2009

Keywords:

Musculoskeletal pain

Probability

Low back

Shoulder

Neck

Systematic review

ABSTRACT

A wide variety of risk factors for the occurrence and prognostic factors for persistence of non-specific musculoskeletal pain (MSP) are mentioned in the literature. A systematic review of all these factors is not available. Thus a systematic review was conducted to evaluate MSP risk factors and prognostic factors, classified according to the dimensions of the International Classification of Functioning, Disability and Health. Candidate systematic reviews were identified in electronic medical journal databases, including the articles published between January 2000 and January 2008 that employed longitudinal cohort designs. The GRADE Working Group's criteria for assessing the overall level of evidence were used to evaluate the reviews. Nine systematic reviews were included, addressing a total of 67 factors. High evidence supported increased mobility of the lumbar spine and poor job satisfaction as risk factors for low back pain. There was also high evidence for intense pain during the onset of shoulder and neck pain and being middle aged as risk factors for shoulder pain. High evidence was also found for several factors that were not prognostic factors. For whiplash-associated disorders these factors were older age, being female, having angular deformity of the neck, and having an acute psychological response. Similarly, for persistence of low back pain, high evidence was found for having fear-avoidance beliefs and meagre social support at work. For low back pain, high evidence was found for meagre social support and poor job content at work as not being risk factors.

© 2009 International Association for the Study of Pain. Published by Elsevier B.V. All rights reserved.

1. Introduction

Work is viewed as being beneficial for health and for social economic status [168]. However, when musculoskeletal pain (MSP) is present, work can be burdensome, resulting in reduced productivity, increased sick leave, and high costs for society [28,56,121]. Obtaining better knowledge of risk factors for the onset of MSP and prognostic factors for the persistence of MSP could provide tailored interventions [59,94,138].

In a healthy population various risk factors of MSP exist. As soon as MSP emerges, it may run its normal course; but in some people, pain lasts longer and may become chronic. These influencing factors are called prognostic factors. Several theoretical models have been proposed that describe the development and prolongation of MSP [72,122,162]. Some reflect contradictory theoretical relationships between the cause and the consequence of MSP. For example,

Waddell's biopsychosocial model is based on neurophysiological or physiological dysfunction [162]. A work-related model is Kasarek's Job Control-Demand model [72]. This situation-centred psychosocial model assumes that a disbalance between high job demands and low worker control results in poor subjective health. A person-centred model is the catastrophizing hypothesis model, which posits that fear of pain results in self-limitation of activity and could therefore be a prognostic factor [122]. All these models have their own paradigm, which may possibly lead to confusion. The International Classification of Functioning, Disability and Health (ICF), however, lacks a paradigm [168]. Instead of explaining causal relationships, the ICF classifies them (Fig. 1) [168]. Therefore the ICF can be used to disentangle a diversity of relationships.

The variety and the number of factors stated in the different ICF dimensions make it difficult for healthcare professionals to judge the relative importance of different risk and prognostic factors [27]. Moreover, several medical disciplines have their own guideline recommendations for employers and patients. These guidelines focus on different risk and prognostic factors [12,84,161]. For example, occupational guidelines for preventing low back pain (LBP) list physically or psychologically demanding work as causal

* Corresponding author. Address: Hanze University Groningen, University of Applied Sciences, P.O. Box 3109, 9701 DC, Groningen, The Netherlands. Tel.: +31 50 4095944.

E-mail address: a.e.jorna-lakke@pl.hanze.nl (S.E. Lakke).

Fig. 1. International Classification of Functioning, Disability and Health [168].

factors of MSP [161]. By contrast, the Dutch physical therapist guideline for LBP lists pain behaviour, fear avoidance and patients' social environment as prognostic factors [12], whereas the clinical guideline of the Norwegian Back Pain Network lists heritage, life-style and low physical activity as risk factors for acute LBP [84]. The guidelines are based on several levels of evidence, from authority-based judgements to systematic reviews of longitudinal and transversal studies. Currently a thorough overview of these predictive factors, regardless of specialism, is lacking. This could result in clinicians being ill informed of how to correctly advice patients and employers to appropriately consider risk and prognostic factors during treatment.

The aim of this review was to qualify and classify the evidence presented in systematic reviews of risk and prognostic factors for non-specific MSP within the ICF. We summarised the evidence, providing a meta-perspective of existing evidence for factors. Missing components in the model may motivate further research into that specific classification domain.

2. Methods

2.1. Search strategy

A systematic review (SR) is considered to be the highest level of evidence [108]. Many overviews of risk and prognostic factors have been published. For this reason, only SRs were included in this review. To identify relevant SRs, we performed an electronic search of bibliographic literature databases (MEDLINE, CINAHL, EMBASE, and PsycINFO), using keywords, MeSH and free text words (Supplementary Appendix 1) from January 2000 up to January 2008. A sensitive search filter for SRs was used [59]. Additional references of guidelines of MSP and all identified SRs were screened for potential eligible studies.

2.2. Selection of studies

Only full reports written in English and meeting the following inclusion criteria (based on study design, population, and exposure) were selected.

2.2.1. Design

Longitudinal research is the preferred method for identifying causal relationships [94]. Therefore, SRs that summarised prospective or retrospective cohort studies were included in our present review. A SR was defined as a review of studies that systematically searched for evidence, that was based on methodological quality assessment of the included studies and that summarised the findings according to predetermined criteria. We considered a meta-analysis to be a type of SR that uses quantitative methods.

2.2.2. Population

Studies that examined adults, aged 18–70 years, with non-specific MSP (as an outcome variable or inclusion criterion) were in-

cluded. Non-specific MSP was defined as MSP not attributed to recognisable, known specific pathology (e.g., infection, tumour, osteoporosis, ankylosing spondylitis, fracture, inflammatory process, radicular syndrome, cauda equina syndrome, and pregnancy) [28,56]. For SRs analysing risk factors, we included those that examined working populations or community-based populations and that identified at least one risk factor and non-specific MSP as an outcome variable. For SRs analysing prognostic factors, we included studies that identified at least one prognostic factor for prolonged MSP. SRs that included workers on 100% sick leave at baseline assessment were excluded. Additional exclusion criteria, such as acute and chronic or severe and non-severe pain at baseline, were not formulated.

2.2.3. Exposure

We included SRs that investigated whether a person's exposure to various factors (body function and structures, activities, participation, personal and environmental factors) predicted MSP. SRs were excluded that examined the impact of treatments. If an SR summarised several factors, we only extracted the findings for factors based on longitudinal cohort studies.

2.3. Study outline

In the first stage, one reviewer (AEL) screened the title and abstract of candidate articles. In the second stage, two reviewers (AEL and RS) screened the full text of all potential relevant articles to determine whether the article met the inclusion criteria. Because the reviewers were familiar with some of the articles, no blinding of authors and institutes was performed.

2.4. Methodological quality assessment of the included systematic reviews

Two reviewers (AEL and TT) independently assessed the quality of the included SRs using the list of criteria for assessing quality, description of potential bias, internal validity, and statistical criteria (Supplementary Appendix 1) [6–8,68]. For each candidate SR, each criterion was rated as 'met' (+), 'unclear/partly met' (±), or 'not met' (–). The total score was calculated by summing up the numbers of 'met'. The total maximum score was 9 points. The methodological quality of an SR was labelled as 'minor limitation' if the quality score was at least 7 out of 9 points and as 'moderate limitations' if the quality score was at least 4 out of 9 points. SRs meeting less than four of the criteria were SRs with 'major limitations' [68]. The inter-rater agreement between the two reviewers was calculated with Cohen's kappa [33]. Agreement was resolved by consensus between AEL and TT. If disagreement persisted after the consensus meeting a third reviewer (MFR) made the final decision.

2.5. Extraction of data

The following data were used for analysis: population characteristics at baseline, date of ending search strategy, number of cohorts and included subjects, study design, methodological quality assessment of included cohort studies, consistency of the available evidence of factors, range of time over which follow-up measurements were made, and outcome measurements. The cohort studies of the included SRs were checked for double counting of extracted risk or prognostic factors based on repetition of cohort studies. When we encountered more than one SR that assessed the methodological quality of the same cohort study, we extracted the cohort study assessments from the SR that was of the highest methodological quality. Identified risk and prognostic factors were classified according to ICF [168]. One reviewer (AEL) extracted the

Table 2
GRADE level of evidence [7,68].

Level of Evidence Quality	Based on:
High-quality evidence	One or more updated, high-quality systematic reviews based on at least 2 high-quality cohort studies ^a with consistent ^b results
Moderate-quality evidence	One or more updated systematic reviews of high or moderate quality based on at least 1 high-quality cohort study based on at least 2 cohort studies of moderate quality with consistent results
Low-quality evidence	One or more systematic reviews of variable quality based on cohort studies of moderate quality based on inconsistent results in the reviews based on inconsistent results in cohort studies
No evidence	No systematic review identified

^a The assessment of the methodological quality of cohort studies was extracted from the included systematic review.

^b Consistent means more than 75% of the included cohorts pointed towards the same direction.

data. To verify accuracy, a second reviewer (RS) selected a random sample ($n = 3$) from the included SRs.

2.6. Level of evidence for each risk and prognostic factor across systematic reviews

The level of evidence and strength of recommendations were assessed according to the criteria assessed by the GRADE Working Group [6,68]. GRADE stands for Grading of Recommendations Assessment, Development and Evaluation. GRADE classifies the level of evidence (high, moderate, low, and none) based on (1) the methodological quality of the SR, (2) the quality of the cohort studies included in the SR, and (3) the consistency of the results of the cohort studies (Table 2). The GRADE level of evidence indicates the extent to which one can be confident that a specific factor predicts MSP or the consequences of MSP.

3. Results

3.1. Literature search

The results of the search strategy are presented in Fig. 2. The literature search of databases resulted in 7937 potentially relevant articles. Excluded on title, abstract and duplicate were 7881 articles. Another 48 articles were excluded after the full text was read. The main reason for exclusion was firstly allowing cross-sectional study design in the reviewed factor of the SR, and secondly non-attendance of methodological quality rating. Screening the refer-

ences of MSP guidelines, all selected articles, and all retrieved SRs resulted in one additional eligible SR. A total of nine SRs were included in the present review [35,53,55,65,66,82,122,140,158]. No meta-analyses were produced in the search.

3.2. Description of systematic reviews

Supplementary Table 3 presents the details of the included SRs. Nine SRs described MSP in predetermined body parts [35,53,55,65,66,82,122,140,158]. Two of the SRs included only prospective cohort studies [122,140], whereas the other seven SRs included both prospective and retrospective studies [35,53,55,65,66,82,158]. Only the risk factor body mass index (BMI) was extracted from the SR of Viikari-Juntura et al. [95,106,158,159], because the other factors assessed by these authors were based on a cross-sectional design. The SR of Scholten-Peeters et al. did not categorize the cohort studies' references for each factor [140]. This observation was confirmed (personal communication; G.M. Scholten-Peeters). Unfortunately these classifications were lost due to removal. Therefore the described cohort studies' references for each prognostic factor [15,23–25,30,38,46,47,58,60–62,73–75,101–105,114,115,117–119,125–131,137,139,146,148,163].

3.3. Double counting

Double counting was checked. Several cohort studies on whip-lash-associated disorders (WAD) were duplicates. Scholten-Peeters et al. [140] included 38 cohort studies on WAD in which the

Fig. 2. Selection of systematic reviews.

subjects' accident occurred less than six days before the start of the study. Coté et al. [35] included subjects that had experienced WAD for less than six weeks [35]. Cote et al.'s SR scored less than Scholten-Peeters et al.'s SR on the methodological quality assessment. Following the preset criteria, we added one cohort study assessed by Cote et al. [82]. For LBP, the risk factor 'social support at the work place' was reviewed in two articles [55,65]. Hartvigsen et al. assessed 10 cohort studies on social support at the work place [41,42,49,63,89,92,116,143,150,170]. Hoogendoorn et al. assessed five cohort studies on the same subject [19,57,92,116,133]. Hartvigsen et al. scored 1 point more than Hoogendoorn et al. on methodological quality. Therefore, Hartvigsen's methodological quality rating of the two duplicated cohort studies was extracted [92,116]. Hamberg-van Reenen et al. included three articles reporting large lumbar flexion [17,50,53,152]. Two of these articles, both rated as having high methodological quality, were related to the same cohort study [50,152]. Thus, both were mentioned but counted as one.

3.4. Participants

The number of subjects ranged from 465 to 27,923 per SR. The included population in SRs considering risk factors consisted of working and community-based subjects. The SRs considering prognostic factors included patients from private and primary care practices, hospital emergency departments, and population- and insurance-based cohorts (Supplementary Table 3).

3.5. Risk and prognostic factors

Five SRs assessed risk factors [53,55,65,66,158]. Two of these evaluated the ICF dimension environmental factors [55,65]; two SRs addressed the dimension of body functions and structure

[53,158]; and one SR assessed factors on the activity and participation dimension [66]. Five SRs assessed prognostic factors on several dimensions of the ICF [35,55,82,122,140]. One SR included cohort studies of both the risk and prognostic factors [55].

Several SRs set the cut-off points for a positive risk estimate at >2.0 and <0.5 [35,66,82,140]. One SR used the same cut-off points to indicate the strength of the association [55]. Another SR presented prognostic factors that used these cut-off points in at least one study [82]. One SR set the criteria for a positive risk or prognostic factor at a statistically significant p -value of 0.10 or less [53]. Three different SRs included statistical analyses in their methodological quality assessments [65,122,158].

3.6. Outcome measurements

A large variety of questionnaires were used to assess MSP in the cohort studies, ranging from self-reported pain, disability, recovery time, sick leave, and incidence of LBP to incidence of claims (Supplementary Appendix 2). The incidence of MSP was measured to determine the risk factors. The consequences of MSP were evaluated for prognostic factors. The outcome measures in Hamberg-van Reenen et al.'s SR varied from incidence of MSP to filing of insurance claims due to MSP [53]. Overall for prognostic factors, a large variety of baseline assessments and follow-up measurements were used. New episodes were not specifically operationalized. Pincus et al.'s criterion for inclusion was acute LBP in patients who had no pain during the preceding three months [122].

3.7. Methodological quality of systematic reviews

The methodological quality of SRs is described in Table 4. Cohen's kappa for overall agreement between the reviewers was

Table 4
Methodological quality of included systematic reviews [68].

	Hartvigsen et al., 2004 [55]	Ijmker et al., 2007 [66]	Kuijpers et al., 2004 [82]	Scholten-Peeters et al., 2003 [140]	Hamberg van Reenen et al., 2007 [53]	Hoogendoorn et al., 2000 [65]	Pincus et al., 2006 [122]	Cote et al., 2001 [35]	Viikari- Juntura et al., 2007 [158]
1 Is the search strategy described in enough detail for the search to be reproducible?	+	+	+	+	+	+	+	+	+
2 Was the search for evidence reasonably comprehensive?	+	+	+	+	+	+	+	±	-
3 Were the criteria used for deciding which studies to include in the review reported?	+	+	+	+	+	+	+	+	+
4 Was bias in the selection of articles avoided?	+	+	+	+	-	-	+	+	+
5 Were the criteria used for assessing the validity of the studies that were reviewed reported?	+	+	+	+	+	+	+	+	-
6 Was the validity of all of the studies referred to in the text assessed using appropriate criteria in analysing the studies that are cited?	+	+	+	+	+	+	+	+	+
7 Were the methods used to combine the findings of the relevant studies (to reach a conclusion) reported? (Best evidence synthesis)	+	+	+	+	+	+	-	-	-
8 Were the findings of the relevant studies combined (or not combined) and analysed appropriately relative to the primary question the review addresses and the available data?	+	+	+	+	+	+	-	-	-
9 Were the conclusions made by the author(s) supported by the data and/or the analysis reported in the review?	+	+	+	+	+	+	+	±	+
Total score	9	9	9	9	8	8	7	5	5

+ = criteria 'met'; ± = criteria 'unclear'/'partly met'; - = criteria 'not met'.

$K = 0.53$, which is considered to represent moderate agreement [4,86]. Full agreement for all criteria ($K = 1.00$) was reached during the consensus meeting. The third assessor did not come into operation. The methodological quality rating of SRs ranged from 5 to 9 points with a median of 8 points. Seven SRs had minor limitations [53,55,65,66,82,122,140]. Since they had a minimum score of 7 out of 9 points. Two had moderate limitations [35,158]. In two SRs, the selection bias could have occurred, because the selection of articles was done by one reviewer [53,65]. Three articles did not report the methods used to combine the findings, nor did these SRs combine the cohort studies appropriately [35,122,158].

3.8. Methodological quality of cohort studies

The methodological quality assessment of the cohort studies was reproduced from the included SRs. The methodological quality for each risk and prognostic factor across SRs varied widely. A criterion for clearly defining the objective of the cohort study was assessed in two SRs [55,65]. One SR described a criterion about the correct statement of the research question [35]. A clear description of the study population was a criterion in five SRs [35,55,65,82,140]. Inclusion and exclusion criteria were described in six SRs [35,53,55,82,140,122]. The response rate at baseline was an assessment criterion in six SRs and varied from a reported minimum of 80% [35,53,55,65,66,82]. A response rate less than 60% was an exclusion criterion in one SR [158]. The dropout-loss-to-follow-up rate was less than 20% in five SRs [65,66,82,140,122]. Two SRs qualitatively expressed the dropout-loss-to-follow-up as 'reasonable' but did not report a percentage [35,158]. Two other SRs rated the criteria positive for sufficient time between baseline and follow-up [53,55]. All included SRs described standardised methods for the data collection of acceptable quality of prognostic or risk factors. One SR judged the prognostic factors on clinical relevance [140]. Another SR assessed the intention of the prognostic factors, such as dose, level, and duration [35].

For the outcome measurements used in the SRs, adequate, standardised, valid, and reliable measure instruments scored one quality point in all SRs. Four SRs gained one quality assessment point, if comparison between the dropout group and the follow-up group at baseline was measured [35,53,82,122].

The data analyses described in the SRs were assessed for whether a multivariate analysis was done. A confounder control was assessed in all the SRs. Three SRs gained one quality point because the number of cases in the multivariate analyses was at least 10 times the number of independent variables [53,55,56]. Two other SRs reported sufficient numbers of subjects [82] and more than 200 subjects in the analysis sample [122].

3.9. Level of evidence based on GRADE dimensions

The level of evidence for risk and prognostic factors for MSP according to GRADE was classified within the ICF dimensions (Table 5). This level of evidence was based on the methodological quality of each SR, the methodological quality of the cohort studies included in the SRs, and the consistency of the results of the cohort studies (Table 5). Highly rated evidence is described in Section 3.9.

3.9.1. Body function and structure

3.9.1.1. Risk factors. In two SRs, 15 cohort studies reported mobility of the spine as risk or prognostic factor for MSP. The results for neck mobility were inconsistent (Table 5). One SR reported increased mobility of the lumbar spine as a risk factor for LBP [53]. The two cohort studies considered in this SR were deemed to have high methodological quality and showed the same positive direc-

tion [17,50,152]. Two articles researched the same cohort and were therefore counted as one. According to the GRADE-based assessment, high evidence was found for increased mobility of the lumbar spine is a risk factor for lumbar pain.

3.9.1.2. Prognostic factors. For two SRs [82,140], that included eight cohort studies, high evidence was found that intense pain intensity at the onset of shoulder and neck pain is a prognostic factor for the duration of symptoms [82,140]. Mental functions were investigated in a population with WAD [140]. Four included cohort studies found no association between 'high acute psychological response' after a car accident and prolonged WAD. One included cohort study found a positive association. Because more than 75% of the results pointed in the same direction, according to GRADE, it can be concluded that there is high evidence that 'high acute psychological response' is not a prognostic factor for WAD.

3.9.2. Activity and participation

3.9.2.1. Risk factors. None of the included SRs examined risk factors for MSP on the activities and participation dimension.

3.9.2.2. Prognostic factors. One SR [82] identified high-activity limitations and participation restrictions at baseline, and another SR [140] identified low workload in neck muscles and driving occupation as prognostic factors for neck and shoulder disorder [140]. The results of the included cohort studies in these SRs were all in the same positive direction for prognostic factors for neck and shoulder disorder. However, they were each based on only one high methodological quality cohort study; therefore, these SRs were rated as providing moderate evidence. High evidence could not be obtained in these SRs on the activities and participation dimension of the ICF.

3.9.3. Environmental factors

3.9.3.1. Risk factors. One high-quality SR examined low job satisfaction as a risk factor for LBP [65]. This SR included six cohort studies. Five cohort studies were rated as methodologically high quality. These five cohort studies showed positive results. One methodologically low-quality cohort study showed no results. High evidence was produced showing that low job satisfaction is a risk factor for LBP. Poor job content (defined as monotonous work, work with few possibilities for new learning and developing knowledge and skills) was rated as high evidence for not being a risk factor for low back disorder; this conclusion is based on one SR [65] that included four high-quality cohort studies showing no results. Poor social support at work (e.g., meagre social support from co-workers and supervisors, relationships at work, and problems with workmates and supervisors) was reviewed in two SRs [15,18] that assessed 13 cohort studies. According to GRADE, high evidence was produced showing that poor social support at work is not a risk factor for LBP.

3.9.3.2. Prognostic factors. There is high evidence that poor social support at work is not a prognostic factor for LBP; this conclusion is based on one SR [15] that included nine cohort studies.

3.9.4. Personal factors

3.9.4.1. Risk factors. No SR was included that measured personal factors as risk factors for MSP.

3.9.4.2. Prognostic factors. Contrary to environmental factors, personal factors are recognized but not classified in the ICF [168]. Personal factors are defined in the ICF as the background of an individual's life [168]. Fear-avoidance beliefs as a prognostic factor was measured in nine cohort studies as an individual's life

Table 5
GRADE-classified level of evidence for MSP risk and prognostic factors according to ICFdimensions.

Risk or prognostic factors					Quality of evidence							
ICF dimension		Factors identified	MSP of body part	QR	No. of cohort studies with positive or no results and methodological quality (high/low)				Summarised results	Risk factor	Prognostic factor	
					Pos		No results					
					high	low	high	low				
Body function and structure	muscle function	poor trunk muscle strength	low back	1 +[53]	5[9,10,16,50,99,100,149]		13[2,9,11,17,48,50,83,88,93,99,100,110,134,147,149,152]		no	low		
			neck	1 +[53]			1[52]		no	moderate		
		poor trunk muscle endurance	low back	1 +[53]	2[2,3,17]		7[2,3,48,77,83,88,93,100,149,147]		no	low		
			neck	1 +[53]			1[9,10]		no	moderate		
	spine mobility	increased mobility of the lumbar spine	low back	1 +[53]	2[17,50,152]				pos	high		
			low back	1 +[53]	1[17,100]		1[1]		pos	low		
		reduced mobility	neck	1 +[53]			1[52]		no	moderate		
			neck	1 +[140]	3*		2*		pos			
	pain	pain of neck or head before collision	neck	2(1+,1±)[140,35]	2*[141]		4*		1*	no	low	
			high initial pain at baseline	neck	1 +[140]	4*		1*		pos	high	
		high pain intensity at baseline	shoulder	1 +[82]	2[96,156]				pos	high		
		concomitant neck pain	shoulder	1 +[82]	1[156]				pos	moderate		
		radicular symptoms	neck	1 +[140]	1*		3*		2*	5*	no	low
		sleeping disturbance	neck	1 +[140]	2*		2*			equal	low	
		angular deformity	neck	1 +[140]			1*		3*	4*	no	high
		initial disc changes	neck	1 +[140]			2*		2*		no	low
		no cause of overuse	shoulder	1 +[82]	1[156]				2*		pos	moderate
		unusual activity	shoulder	1 +[82]	1[156]						pos	moderate
	body structure	no acute bursitis	shoulder	1 +[82]	1[156]						pos	
		BMI	shoulder	1 ±[158]	3[95,106,159]						pos	moderate
body position		unprepared for collision	neck	1 +[140]	2*		2*		2*	no	low	
		car accident (WAD)	turned head	neck	1 +[140]	2*		1*		pos	low	
	position	velocity change > 10 km/h	neck	1 +[140]	1*		1*		equal	low		
pre-existing changes	pre-existing spondylosis + degenerative changes	neck	1 +[140]	2*		1*		2*	4*	no	low	
mental functions	cognitive impairments	neck	1 +[140]	1*				1*	pos	low		
		neck	1 +[140]	1*				1*	pos	low		
	speed of information processing	neck	1 +[140]	1*				1*	pos	low		
		poor concentration	neck	1 +[140]	1*				1*	pos	low	
	neurotism	neck	1 +[140]	1				1	pos	low		

Activities and participation		increased acute psychological response	neck	1+[140]	1*		4*	1*	no		high
		reduced workload on neck muscle	neck	1+[140]	1*				pos		moderate
		driving occupation	neck	1+[140]	1*				pos		moderate
		high disability score at baseline	shoulder	1+[82]	1[96]	1[36]			pos		moderate
Personal factors		fear avoidance	low back	1+[122]			8[78,80,120,124,142,144,145,165]	1[29]	no		high
		depression	low back	1+[122]	1[145]				pos		moderate
		previous psychiatric problems	neck	1+[140]	1*				pos		moderate
		stress unrelated to accident	neck	1+[140]	1*			1*	pos		low
		nervousness	neck	1+[140]	1*				pos		moderate
		need to resume physiotherapy	neck	1+[140]		1*			pos		low
		need for cervical collar > 12 weeks	neck	1+[140]	1*			1*	pos		low
		older age	neck	1+[140]	2*		10*	10*	no		high
		female gender	neck	1+[140]	3*	2*	12*	9*	no		high
		middle age (45–54 yr)	shoulder	1+[82]	2[31,107]				pos		high
Environmental factors	work perception	perception of work	low back	1+[55]	1[157]	1[116]	3[43,44,63]	4[21,111,132,160,170]	no	low	
			low back	1+[55]	3[64,154,157]	3[42,153,155]	3[19,20,34,51]	10[20,45,54,57,67,85,112,116,151,167]	no		low
	work organisational aspects	low job satisfaction	low back	1+[65]	5[17,19,116,132,136]			1[57]	pos	high	
		stress at work	low back	1+[55]	1[43]		1[42]	3[49,132,170]	no	low	
			low back	1+[55]			1[42]	2[91,166]	no		moderate
		poor social support at work	low back	2+[55,65]	2[19,133]		4[42,49,57,63]	8[41,49,89,92,116,144,150,170]	no	high	
			low back	1+[55]	1[42]		4[64,153–155]	4[45,57,91,116]	no		high
		poor job content	low back	1+[65]	1[92]		4[17,92,133,136]		no	high	
		organisational aspects of work	low back	1+[55]	1[42]	1[92]	2[40,63]	5[21,49,89,143,150]	no	low	
			low back	1+[55]	2[51,57]		4[42,64,153,154]	3[142,151,166]	no		low
	work instruments	low job control	low back	1+[65]			1[92]		no	moderate	
		low decision latitude	low back	1+[65]	1[57]				pos	moderate	
		high pace of work	low back	1+[65]	2[57,133]		1[18]		no	low	
		High-quality job demand	low back	1+[65]	1[57]	1[14]			pos	moderate	
		total computer use time	neck	1+[66]	1[98]		1[79]	1[69,71]	no	low	
			upper extremity	1+[66]	1[98]			1[69,71]	pos	low	
		mouse use time	neck	1+[66]	1[5,26,81,87]				pos	moderate	
			upper extremity	1+[66]	1[5,26,81,87]	1[69,71]			pos	moderate	
social contact		keyboard use time	neck	1+[66]	1[5,26,81,87]				pos	moderate	
			upper extremity	1+[66]	1[5,26,81,87]				pos	moderate	
		overload at work	shoulder	1+[82]	1[106]				pos		moderate
		social support	low back	1+[65]		1[109]	1[19]		no	low	

(continued on next page)

Table 5 (continued)

Risk or prognostic factors			QR	No. of cohort studies with positive or no results and methodological quality (high/low)	Quality of evidence				
ICF dimension	Factors identified	MSP of body part			Summarised results	Risk factor	Prognostic factor		
					Pos		No results		
					high	low	high	low	
crash position	initial treatment	neck	1 + [140]		1*		1*	pos	low
	hospital crash rear-end collision	neck	1 + [140]	1*	1*	5*	7*	no	high
	accident on highway	neck	1 + [140]	1*				pos	moderate
	car stationary	neck	1 + [140]	1*		1*	1*	low	low
social security services	car stationary when hit rear-end	neck	1 + [140]		1*				
	compensation	neck	1 + [140]		1*	3*	3*	no	high

MBI = Body mass index; ICF = International Classification of Functioning, Disability and Health; MSP = musculoskeletal pain; QR = number and Quality of systematic review; + = minor limitations; ± = moderate limitations. Results of cohort studies: pos, evidence for risk or prognostic factor; no, no evidence for risk or prognostic factor; high, methodological high quality cohort study; low, methodological low quality cohort study. Summarised results: pos, evidence for a risk or prognostic factor; no, no evidence for a risk or prognostic factor; [15,23–25,30,38,46,47,58,60–62,73–75,101–105,114,115,117–119,125–131,137,139,146,148,150,163]. All the Cohort studies of Scholten-Peeters et al.

background and not as an impairment [122]. Therefore, in this SR, fear avoidance was classified on the personal factors dimension. One SR fulfilled our preset inclusion criteria [122]. Eight high-quality and one low-quality methodological cohort study concluded that fear-avoidance beliefs were not a prognostic factor for LBP. Following GRADE, we concluded that high evidence was present, showing that fear-avoidance beliefs are not a prognostic factor for LBP. High evidence was produced showing that being female and being old age are not the prognostic factors for WAD; this conclusion is based on one SR [140] that included several cohort studies. One SR [82] investigated the prognostic factor age (45–54 years) in two cohort studies rated as having high methodological quality. Following GRADE's criteria of evidence, we conclude that high evidence was produced showing that being middle aged is a prognostic factor for persistent shoulder pain.

4. Discussion

The first aim of this SR was to determine the quality of the evidence for MSP risk and prognostic factors by using findings from available SRs as a basis. There is high evidence that increased lumbar spine mobility and low job satisfaction are risk factors for the development of LBP. High evidence for prognostic factors for neck and shoulder pain are baseline neck and shoulder pain intensity, and a prognostic factor for shoulder pain is being middle aged. There is high evidence that older age, being female, angular deformity of the neck, and acute psychological response are not prognostic factors for persistent WAD. For LBP, there is high evidence that fear avoidance and poor social support at work are not prognostic factors for LBP. Poor social support at work and poor job content are not risk factors for LBP.

The second aim of this SR was to summarise the quality of evidence in terms of the ICF classification scheme to identify missing areas for further research. The ICF provides a systematic coding scheme for health information systems, establishing a common language to improve communication between different users; it also takes a neutral stand with regard to specialism and underlying theoretical models [168]. A limited number of cohort studies measured prognostic factors for MSP on the activities and participation dimension of the ICF, with all pointing towards the same positive direction for possible prognostic factors for MSP [96,36,140]. Due to the meagre number of cohort studies, none of these factors were graded as high level of evidence. In addition, no SR summarised risk factors on the ICF activities and participation dimension for the onset of MSP.

Another remarkable lack of factors could be recognized in the ICF framework. No included SRs measured the risk factors on the personal dimension. Furthermore, environmental risk and prognostic factors, such as 'work perception', were only found for LBP, not pain in other body parts. Firstly, because the present SR only included SRs, our main recommendations for future research agendas are to fill in the gaps in the ICF given in Table 5 with SRs. Secondly, if SRs are not feasible or not yet available, this table could be populated with single prospective cohort studies.

The strength of this SR lies in the number of participants included ($N = 119,849$) and in an exhaustive search of multiple electronic databases. This SR gives an overview of the systematically reviewed risk and prognostic factor literature, which consisted of longitudinal cohort studies that were all rated on methodological quality. The results of this SR with regard to prognostic factors are of clinical relevance and should have implications for practice. Psychosocial yellow flags in acute LBP are defined as risk factors for long-term disability and work loss [76]. Identification of at-risk individuals should lead to appropriate early management targeted towards the prevention of chronic pain and disability. The defini-

tion of prognostic factor is identical to these yellow flags. High neck and shoulder pain intensity could be added as yellow flags. On the other hand, with regard to LBP, fear-avoidance beliefs and poor social support at work perhaps should be removed as yellow flags [12,16,84].

As with all SRs, one limitation of the present SR is heterogeneity, which could cause effect bias. To limit the risk of bias, two reviewers independently assessed the methodological quality of the studies with a validated instrument [68], and two reviewers performed the search strategy for the second stage. Another problem inherent to all SRs is the publication bias. Because of the extent of the issue we assessed, publications could have been missed [40]. However, since we used a comprehensive search strategy, it is unlikely that any publications were missed.

The ICF defines personal factors in terms of the particular background of an individual's life and way of living and the domain mental functions as a manifestation of pathology [168]. One could argue about the ICF classification of the factors in this review. For example, the factor 'nervousness' was classified as a personal factor dimension and not as a mental impairment. Classifying these factors differently would affect the 'umbrella overview' of the existing evidence for factors, not the results of the overall quality of this SR.

Apart from the problems discussed thus far, limitations can also arise from the problems of the included SRs. For example, in assessing the risk factors for back pain, employees and community-based populations were summarised without considering the 'healthy workers effect' [97]. Indeed, workers with back pain may leave a job, resulting in a surviving workforce with healthier backs. This may introduce significant membership bias.

The outcome measurements of the primary studies were very diverse. Some measured sick leave, some measured self-reported symptoms. Self-reported physical or mental symptoms do not automatically translate to incapacity for work. One-third of the people reporting physical or mental symptoms function normally at work [162]. In the included SRs, the studies with outcome measures physical symptoms and sick leave were combined. This could have led to an effect bias. However, the variety in outcome measures and the amount of included cohort studies may have equalized possible effect bias.

In this review, cohort studies searching for prognostic factors included acute and chronic, and severe and non-severe MSP at baseline. However, we think that this heterogeneity in baseline characteristics does not significantly affect the findings of the current SR. MSP is an intermittent lifetime problem in which symptomatic periods alternate with symptom-free periods. To increase the clinical relevance, recommendations for future research should agree on outcome measures and baseline characteristics in prognostic cohort research [123].

Our recommendations for future research include performing SRs on initial pain as a prognostic factor for LBP, environmental causal factors for neck or shoulder pain, and causal personal factors for MSP. Furthermore, more methodologically high-quality cohort studies should be carried out to identify the prognostic factors categorized within the ICF activities and participation dimension. Future SRs should also assess and identify risk factors within this dimension. Effect modification of several dimensions of the ICF could occur. For example, personal factors could influence an environmental outcome variable such as job content [22,37,94]. Potential confounders and mediators such as age, gender, job satisfaction, or personal factors such as depressive feelings or motivation, should be taken into account. This SR does not provide a complete overview of the factors influencing MSP in different body parts. Thus, the next step would be to research additional SRs or to fill in the gaps given in Table 5 with cohort studies. A conceptual model of illustrating the relationship between ICF dimensions in

a working population should be built in order to gain insight into the coherence between the different dimensions in a specific population [13,164,169]. Without further research, we will not know whether modifying a person's risk factor would prevent MSP and reduce sick leave. Therefore, the risk factors 'increased mobility of the lumbar spine' and 'low job satisfaction' should not be used as selection criteria for engaging employees.

5. Conclusion

By applying the GRADE method of classifying the level of evidence, we determined that increased lumbar spine mobility and low job satisfaction are high evidence risk factors for LBP. There is high evidence that intense initial pain at baseline and being middle aged (45–54 years) are prognostic factors for neck and shoulder pain and for shoulder pain, respectively. Moreover, there is high evidence showing that older age, being female, angular deformity and acute psychological response are not prognostic factors for prolonged pain in WAD. High evidence also indicated that fear at early stages of pain and poor social support at work are not prognostic factors for LBP. In addition, high evidence indicated that poor job content and poor social support at work are not risk factors for LBP. Recommendations for future research are to systematically review prospective cohort studies on MSP risk factors on the ICF activities and participation dimension and personal dimension. Further recommendations include performing SRs on environmental risk factors for neck and shoulder pain and the prognostic factor initial pain for LBP. Finally, SRs on environmental risk and prognostic factors of MSP other than LBP are recommended.

Acknowledgements

The first author thanks Dr. Harriët Wittink, University of Applied Sciences Utrecht, The Netherlands, for her constructive participation in this study. Financial support for this systematic review has not been provided in any kind. No funds were received to support this work. No benefits in any form have been or will be received from a commercial party related directly or indirectly to the subject of this manuscript.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.pain.2009.08.032](https://doi.org/10.1016/j.pain.2009.08.032).

References

- [1] Adams MA, Mannion AF, Dolan P. Personal risk factors for first-time low back pain. *Spine* 1999;24:2497–505.
- [2] Alaranta H, Luoto S, Heliovaara M, Hurri H. Static back endurance and the risk of low-back pain. *Clin Biomech* 1995;10:323–4.
- [3] Alaranta H, Hurri H, Heliovaara M, Soukka A, Harju R. Flexibility of the spine: normative values of goniometric and tape measurements. *Scand J Rehabil Med* 1994;26:147–54.
- [4] Altman DG. *Practical statistics for medical research*. 1st ed. London, New York: Chapman and Hall; 1991.
- [5] Andersen JH, Thomsen JF, Overgaard E, Lassen CF, Brandt LP, Vilstrup I, Kryger AI, Mikkelsen S. Computer use and carpal tunnel syndrome: a 1-year follow-up study. *JAMA* 2003;289:2963–9.
- [6] Atkins D, Best D, Briss PA, Eccles M, Falck-Ytter Y, Flottorp S, Guyatt GH, Harbour RT, Haugh MC, Henry D, Hill S, Jaeschke R, Leng G, Liberati A, Magrini N, Mason J, Middleton P, Mrukowicz J, O'Connell D, Oxman AD, Phillips B, Schunemann HJ, Edejer TT, Varonen H, Vist GE, Williams Jr JW, Zaza S. Grading quality of evidence, strength of recommendations. *BMJ* 2004;328:1490.
- [7] Atkins D, Eccles M, Flottorp S, Guyatt GH, Henry D, Hill S, Liberati A, O'Connell D, Oxman AD, Phillips B, Schunemann H, Edejer TT, Vist GE, Williams Jr JW. Systems for grading the quality of evidence and the strength of recommendations I: critical appraisal of existing approaches The GRADE Working Group. *BMC Health Serv Res* 2004;4:38.

- [8] Atkins D, Briss PA, Eccles M, Flottorp S, Guyatt GH, Harbour RT, Hill S, Jaeschke R, Liberati A, Magrini N, Mason J, O'Connell D, Oxman AD, Phillips B, Schunemann H, Edejer TT, Vist GE, Williams Jr JW. Systems for grading the quality of evidence and the strength of recommendations II: pilot study of a new system. *BMC Health Serv Res* 2005;5:25.
- [9] Barnekow-Bergkvist M, Hedberg G, Janlert U, Jansson E. Development of muscular endurance and strength from adolescence to adulthood and level of physical capacity in men and women at the age of 34 years. *Scand J Med Sci Sports* 1996;6:145–55.
- [10] Barnekow-Bergkvist M, Hedberg GE, Janlert U, Jansson E. Determinants of self-reported neck–shoulder and low back symptoms in a general population. *Spine* 1998;23:235–43.
- [11] Batti'e MC, Bigos SJ, Fisher LD, Hansson TH, Jones ME, Wortley MD. Isometric lifting strength as a predictor of industrial back pain reports. *Spine* 1989;14:851–6.
- [12] Bekkering GE, Hendriks HJM, Koes BW, Oostendorp RAB, Ostelo RWJG, Thomassen J, van Tulder MW. Dutch physiotherapy guidelines for low back pain. *Physiotherapy* 2003;89:82–96.
- [13] Bennett GG, Wolin KY, Avrunin JS, Stoddard AM, Sorensen G, Barbeau E, Emmons KM. Does race/ethnicity moderate the association between job strain and leisure time physical activity? *Ann Behav Med* 2006;32:60–7.
- [14] Bergenudd H, Nilsson B. Back pain in middle age; occupational workload and psychologic factors: an epidemiologic survey. *Spine* 1988;13:58–60.
- [15] Berglund A, Alfredsson L, Cassidy JD, Jensen I, Nygren A. The association between exposure to a rear-end collision and future neck or shoulder pain: a cohort study. *J Clin Epidemiol* 2000;53:1089–94.
- [16] Biering-Sorensen F. Physical measurements as risk indicators for low-back trouble over a one-year period. *Spine* 1984;9:106–19.
- [17] Biering-Sorensen F, Thomsen CE, Hilden J. Risk indicators for low back trouble. *Scand J Rehabil Med* 1989;21:151–7.
- [18] Biering-Sorensen F, Thomsen C. Medical, social and occupational history as risk indicators for low-back trouble in a general population. *Spine* 1986;11:720–5.
- [19] Bigos SJ, Battie MC, Spengler DM, Fisher LD, Fordyce WE, Hansson TH, Nachemson AL, Wortley MD. A prospective study of work perceptions and psychosocial factors affecting the report of back injury. *Spine* 1991;16:1–6.
- [20] Bigos SJ, Battie MC, Spengler DM, Fisher LD, Fordyce WE, Hansson T, Nachemson AL, Zeh J. A longitudinal, prospective study of industrial back injury reporting. *Clin Orthop Relat Res* 1992;279:21–34.
- [21] Bildt C, Alfredsson L, Michelsen H. Occupational and nonoccupational risk indicators for incident and chronic pain in a sample of the Swedish general population during a 4-year period: on influence of depression? *Int Behav Med* 2000;7:372–92.
- [22] Bongers PM, Kremer AM, ter LJ. Are psychosocial factors, risk factors for symptoms and signs of the shoulder, elbow, or hand/wrist? A review of the epidemiological literature. *Am J Ind Med* 2002;41:315–42.
- [23] Borchgrevink G, Smevik O, Haave I, Haraldseth O, Nordby A, Lereim I. MRI of cerebrum and cervical column within two days after whiplash neck sprain injury. *Injury* 1997;28:331–5.
- [24] Borchgrevink GE, Smevik O, Nordby A, Rinck PA, Stiles TC, Lereim I. MR imaging and radiography of patients with cervical hyperextension–flexion injuries after car accidents. *Acta Radiol* 1995;36:425–8.
- [25] Borchgrevink GE, Stiles TC, Borchgrevink PC, Lereim I. Personality profile among symptomatic and recovered patients with neck sprain injury, measured by MCMI-I acutely and 6 months after car accidents. *J Psychosom Res* 1997;42:357–67.
- [26] Brandt LP, Andersen JH, Lassen CF, Kryger A, Overgaard E, Vilstrup I, Mikkelsen S. Neck and shoulder symptoms and disorders among Danish computer workers. *Scand J Work Environ Health* 2004;30:399–409.
- [27] Breen AC, van Tulder MW, Koes BW, Jensen I, Reardon R, Bronfort G. Mono-disciplinary or multidisciplinary back pain guidelines? How can we achieve a common message in primary care? *Eur Spine J* 2006;15:641–7.
- [28] Burton AK, Balague F, Cardon G, Eriksen HR, Henrotin Y, Lahad A, Leclerc A, Muller G, van der Beek AJ, Chapter 2. European guidelines for prevention in low back pain: November 2004. *Eur Spine J* 2006;15:S136–68.
- [29] Burton AK, McClune TD, Clarke RD, Main CJ. Long-term follow-up of patients with low back pain attending for manipulative care: outcomes and predictors. *Man Ther* 2004;9:30–5.
- [30] Cassidy JD, Carroll LJ, Cote P, Lemstra M, Berglund A, Nygren A. Effect of eliminating compensation for pain and suffering on the outcome of insurance claims for whiplash injury. *N Engl J Med* 2000;342:1179–86.
- [31] Cassou B, Derriennic F, Monfort C, Norton J, Touranchet A. Chronic neck and shoulder pain, age, and working conditions: longitudinal results from a large random sample in France. *Occup Environ Med* 2002;59:537–44.
- [32] Chaffin DB, Herrin GD, Keyserling WM. Preemployment strength testing: an updated position. *J Occup Med* 1978;20:403–8.
- [33] Cohen J. Coefficient of agreement for nominal scales. *Educ Psychol Meas* 1960;20:37–46.
- [34] Coste J, Delecoeur G, Cohen de LA, Le Parc JM, Paolaggi JB. Clinical course and prognostic factors in acute low back pain: an inception cohort study in primary care practice. *BMJ* 1994;308:577–80.
- [35] Cote P, Cassidy JD, Carroll LJ, Frank JW, Bombardier C. A systematic review of the prognosis of acute whiplash and a new conceptual framework to synthesize the literature. *Spine* 2001;26:E445–58.
- [36] Croft P, Pope D, Silman A. The clinical course of shoulder pain: prospective cohort study in primary care. Primary care rheumatology society shoulder study group. *BMJ* 1996;313:601–2.
- [37] Davis KG, Heaney CA. The relationship between psychosocial work characteristics and low back pain: underlying methodological issues. *Clin Biomech* 2000;15:389–406.
- [38] Drottning M, Staff PH, Levin L, Malt UFR. Acute emotional response to common whiplash predicts subsequent pain complaints. *Nord J Psychiatry* 1995;49:293–9.
- [39] Dueker JA, Ritchie SM, Knox TJ, Rose SJ. Isokinetic trunk testing and employment. *J Occup Med* 1994;36:42–8.
- [40] Easterbrook PJ, Berlin JA, Gopalan R, Matthews DR. Publication bias in clinical research. *Lancet* 1991;337:867–72.
- [41] Elfering A, Semmer NK, Schade V, Grund S, Boos N. Supportive colleague, unsupportive supervisor: the role of provider-specific constellations of social support at work in the development of low back pain. *J Occup Health Psychol* 2002;7:130–40.
- [42] Elfering A, Grebner S, Semmer NK, Gerber H. Time control, catecholamines and back pain among young nurses. *Scand J Work Environ Health* 2002;28:386–93.
- [43] Feuerstein M, Berkowitz SM, Huang GD. Predictors of occupational low back disability: implications for secondary prevention. *J Occup Environ Med* 1999;41:1024–31.
- [44] Feyer AM, Heribson P, Williamson AM, de Silva I, Mandryk J, Hendrie L, Hely MC. The role of physical and psychological factors in occupational low back pain: a prospective cohort study. *Occup Environ Med* 2000;57:116–20.
- [45] Fransen M, Woodward M, Norton R, Coggan C, Dawe M, Sheridan N. Risk factors associated with the transition from acute to chronic occupational back pain. *Spine* 2002;27:92–8.
- [46] Gargan M, Bannister G, Main C, Hollis S. The behavioural response to whiplash injury. *J Bone Joint Surg Br* 1997;79:523–6.
- [47] Gargan MF, Bannister GC. The rate of recovery following whiplash injury. *Eur Spine J* 1994;3:162–4.
- [48] Gibbons LE, Videman T, Battie MC. Isokinetic and psychophysical lifting strength, static back muscle endurance, and magnetic resonance imaging of the paraspinal muscles as predictors of low back pain in men. *Scand J Rehabil Med* 1997;29:187–91.
- [49] Gonge H, Jensen LD, Bonde JP. Do psychosocial strain and physical exertion predict onset of low-back pain among nursing aides? *Scand J Work Environ Health* 2001;27:388–94.
- [50] Griffin AB, Troup JD, Lloyd DC. Tests of lifting and handling capacity. Their repeatability and relationship to back symptoms. *Ergonomics* 1984;27:305–20.
- [51] Hagen KB, Tambs K, Bjerkedal T. A prospective cohort study of risk factors for disability retirement because of back pain in the general working population. *Spine* 2002;27:1790–6.
- [52] Härmäläinen O, Vanharanta H, Bloigu R. +Gz-related neck pain: a follow-up study. *Aviat Space Environ Med* 1994;65:16–8.
- [53] Hamberg-van Reenen HH, Ariens GA, Blatter BM, van Tulder MW, Bongers PM. A systematic review of the relation between physical capacity and future low back and neck/shoulder pain. *Pain* 2007;130:93–107.
- [54] Harkapaa K. Psychosocial factors as predictors for early retirement in patients with chronic low back pain. *J Psychosom Res* 1992;36:553–9.
- [55] Hartvigsen J, Lings S, Leboeuf-Yde C, Bakketeig L. Psychosocial factors at work in relation to low back pain and consequences of low back pain; a systematic, critical review of prospective cohort studies. *Occup Environ Med* 2004;61:e2.
- [56] Helliwell PS, Bennett RM, Littlejohn G, Muirhead KD, Wigley RD. Towards epidemiological criteria for soft-tissue disorders of the arm. *Occup Med* 2003;53:313–9.
- [57] Hemingway H, Shipley MJ, Stansfeld S, Marmot M. Sickness absence from back pain, psychosocial work characteristics and employment grade among office workers. *Scand J Work Environ Health* 1997;23:121–9.
- [58] Herrstrom P, Lannerbro-Geijer G, Hogstedt B. Whiplash injuries from car accidents in a Swedish middle-sized town during 1993–95. *Scand J Prim Health Care* 2000;18:154–8.
- [59] Higgins JPT, Green S. Cochrane handbook for systematic reviews of interventions 4.2.6 [updated September 2006]. In: The Cochrane Library, Issue 4. Chichester, UK: John Wiley & Sons, Ltd.; 2006.
- [60] Hildingsson G, Toolanen G. Outcome after soft-tissue injury of the cervical spine. A prospective study of 93 car-accident victims. *Acta Orthop Scand* 1990;61:357–9.
- [61] Hodgson SP, Grundy M. Whiplash injuries: their long prognosis and its relationship to compensation. *Neuro-orthopedics* 1989;7:88–91.
- [62] Hohl M. Soft-tissue injuries of the neck in automobile accidents. Factors influencing prognosis. *J Bone Joint Surg Am* 1974;56:1675–82.
- [63] Hoogendoorn WE, Bongers PM, de Vet HC, Houtman IL, Ariens GA, van Tulder MW, Bouter LM. Psychosocial work characteristics and psychological strain in relation to low-back pain. *Scand J Work Environ Health* 2001;27:258–67.
- [64] Hoogendoorn WE, Bongers PM, de Vet HC, Ariens GA, van Tulder MW, Bouter LM. High physical work load and low job satisfaction increase the risk of sickness absence due to low back pain: results of a prospective cohort study. *Occup Environ Med* 2002;59:323–8.
- [65] Hoogendoorn WE, van Poppel MN, Bongers PM, Koes BW, Bouter LM. Systematic review of psychosocial factors at work and private life as risk factors for back pain. *Spine* 2000;25:2114–25.

- [66] Ijmker S, Huysmans MA, Blatter BM, van der Beek AJ, van Tulder MW, Bongers PM. Should office workers spend fewer hours at their computer? A systematic review of the literature. *Occup Environ Med* 2007;64:211–22.
- [67] Infante RC, Lortie M. Prognostic factors for return to work after a first compensated episode of back pain. *Occup Environ Med* 1996;53:488–94.
- [68] Jamtvedt G, Dahm KT, Christie A, Moe RH, Haavardsholm E, Holm I, Hagen KB. Physical therapy interventions for patients with osteoarthritis of the knee: an overview of systematic reviews. *Phys Ther* 2008;88:123–36.
- [69] Jensen C. Development of neck and hand-wrist symptoms in relation to duration of computer use at work. *Scand J Work Environ Health* 2003;29:197–205.
- [70] Josephson M, Hagberg M, Hjelm EW. Self-reported physical exertion in geriatric care. A risk indicator for low back symptoms? *Spine* 1996;21:2781–5.
- [71] Juul-Kristensen B, Sogaard K, Stroyer J, Jensen C. Computer users' risk factors for developing shoulder, elbow and back symptoms. *Scand J Work Environ Health* 2004;30:390–8.
- [72] Karasek RA. Job demands, job decision latitude, and mental strain: implications for job redesign. *Adv Sci Quart* 1979;24:285–308.
- [73] Karlsborg M, Smed A, Jespersen H, Stephensen S, Cortsen M, Jennum P, Herning M, Korftsen E, Werdelin L. A prospective study of 39 patients with whiplash injury. *Acta Neurol Scand* 1997;95:65–72.
- [74] Kasch H, Bach FW, Jensen TS. Handicap after acute whiplash injury: a 1-year prospective study of risk factors. *Neurology* 2001;56:1637–43.
- [75] Kasch H, Stengaard-Petersen K, Arendt-Nielsen L, Jensen TS. Headache, neck pain and neck mobility after acute whiplash injury: a prospective study. *Spine* 2001;26:1246–51.
- [76] Kendall NAS, Linton SJ, Main, CJ. Guide to assessing psycho-social yellow flags in acute low back pain: risk factors for long-term disability and work loss. Wellington, New Zealand: Accident Compensation Corporation and the New Zealand Guidelines Group; 1997 [October 2004 ed.].
- [77] Klaber Moffett JA, Hughes GI, Griffiths P. A longitudinal study of low back pain in student nurses. *Int J Nurs Stud* 1993;30:197–212.
- [78] Koleck M, Mazaux JM, Rascle N, Bruchon-Schweitzer M. Psycho-social factors and coping strategies as predictors of chronic evolution and quality of life in patients with low back pain: a prospective study. *Eur J Pain* 2006;10:1–11.
- [79] Korhonen T, Ketola R, Toivonen R, Luukkonen R, Hakkanen M, Viikari-Juntura E. Work related and individual predictors for incident neck pain among office employees working with video display units. *Occup Environ Med* 2003;60:475–82.
- [80] Kovacs FM, Abaira V, Zamora J, Fernandez C. The transition from acute to subacute and chronic low back pain: a study based on determinants of quality of life and prediction of chronic disability. *Spine* 2005;30:1786–92.
- [81] Kryger AI, Andersen JH, Lassen CF, Brandt LP, Vilstrup I, Overgaard E, Thomsen JF, Mikkelsen S. Does computer use pose an occupational hazard for forearm pain; from the NUDATA study. *Occup Environ Med* 2003;60:e14.
- [82] Kuijpers T, van der Windt DA, van der Heijden GJ, Bouter LM. Systematic review of prognostic cohort studies on shoulder disorders. *Pain* 2004;109:420–31.
- [83] Kujala UM, Taimela S, Viljanen T, Jutila H, Viitasalo JT, Videman T, Battie MC. Physical loading and performance as predictors of back pain in healthy adults. A 5-year prospective study. *Eur J Appl Physiol Occup Physiol* 1996;73:452–8.
- [84] Laerum E, Dullerud R, Kirkesola G, Mengshoel AM, Nygaard OP, Skouen JS, Stig LC, Werner E. Acute low back pain. Interdisciplinary clinical guidelines. Oslo: The Norwegian Back Pain Network; 2002.
- [85] Lancourt J, Kettelhut M. Predicting return to work for lower back pain patients receiving worker's compensation. *Spine* 1992;17:629–40.
- [86] Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics* 1977;33:159–74.
- [87] Lassen CF, Mikkelsen S, Kryger AI, Brandt LP, Overgaard E, Thomsen JF, Vilstrup I, Andersen JH. Elbow and wrist/hand symptoms among 6, 943 computer operators: a 1-year follow-up study (the NUDATA study). *Am J Ind Med* 2004;46:521–33.
- [88] Latikka P, Battie MC, Videman T, Gibbons LE. Correlations of isokinetic and psychophysical back lift and static back extensor endurance tests in men. *Clin Biomech* 1995;10:325–30.
- [89] Latza U, Pfahlerberg A, Gefeller O. Impact of repetitive manual materials handling and psychosocial work factors on the future prevalence of chronic low-back pain among construction workers. *Scand J Work Environ Health* 2002;28:314–23.
- [90] Lee JH, Hoshino Y, Nakamura K, Kariya Y, Saita K, Ito K. Trunk muscle weakness as a risk factor for low back pain. A 5-year prospective study. *Spine* 1999;24:54–7.
- [91] Lehmann TR, Spratt KF, Lehmann KK. Predicting long-term disability in low back injured workers presenting to a spine consultant. *Spine* 1993;18:1103–12.
- [92] Leino PI, Hanninen V. Psychosocial factors at work in relation to back and limb disorders. *Scand J Work Environ Health* 1995;21:134–42.
- [93] Leino P, Aro S, Hasan J. Trunk muscle function and low back disorders: a ten-year follow-up study. *J Chronic Dis* 1987;40:289–96.
- [94] Linton SJ, Gross D, Schultz IZ, Main C, Cote P, Pransky G, Johnson W. Prognosis and the identification of workers risking disability: research issues and directions for future research. *J Occup Rehabil* 2005;15:459–74.
- [95] Luime JJ, Kuiper JJ, Koes BW, Verhaar JA, Miedema HS, Burdorf A. Work-related risk factors for the incidence and recurrence of shoulder and neck complaints among nursing-home and elderly-care workers. *Scand J Work Environ Health* 2004;30:279–86.
- [96] Macfarlane GJ, Hunt IM, Silman AJ. Predictors of chronic shoulder pain: a population based prospective study. *J Rheumatol* 1998;25:1612–5.
- [97] Manek NJ, Macgregor AJ. Epidemiology of back disorders: prevalence, risk factors, and prognosis. *Curr Opin Rheumatol* 2005;17:134–40.
- [98] Marcus M, Gerr F, Monteilh C, Ortiz DJ, Gentry E, Cohen S, Edwards A, Ensor C, Kleinbaum D. A prospective study of computer users: II. Postural risk factors for musculoskeletal symptoms and disorders. *Am J Ind Med* 2002;41:236–49.
- [99] Masset DF, Piette AG, Malchaire JB. Relation between functional characteristics of the trunk and the occurrence of low back pain. Associated risk factors. *Spine* 1998;23:359–65.
- [100] Mayer TG, Tencer AF, Kristoferson S, Mooney V. Use of noninvasive techniques for quantification of spinal range-of-motion in normal subjects and chronic low-back dysfunction patients. *Spine* 1984;9:588–95.
- [101] Mayou R, Bryant B, Duthie R. Psychiatric consequences of road traffic accidents. *BMJ* 1993;307:647–51.
- [102] Mayou R, Bryant B. Outcome of 'whiplash' neck injury. *Injury* 1996;27:617–23.
- [103] Meenen NM, Katzer A, Dihlmann SW, Held S, Fyfe I, Jungbluth KH. Whiplash injury of the cervical spine – on the role of pre-existing degenerative diseases. *Unfallchirurgie* 1994;20:138–48.
- [104] Miles KA, Maimaris C, Finlay D, Barnes MR. The incidence and prognostic significance of radiological abnormalities in soft tissue injuries to the cervical spine. *Skelet Radiol* 1988;17:493–6.
- [105] Minton R, Murray P, Stephenson W, Galasko CS. Whiplash injury – are current head restraints doing their job? *Accid Anal Prev* 2000;32:177–85.
- [106] Miranda H, Viikari-Juntura E, Heistaro S, Heliovaara M, Riihimäki H. A population study on differences in the determinants of a specific shoulder disorder versus nonspecific shoulder pain without clinical findings. *Am J Epidemiol* 2005;161:847–55.
- [107] Miranda H, Viikari-Juntura E, Martikainen R, Takala EP, Riihimäki H. A prospective study of work related factors and physical exercise as predictors of shoulder pain. *Occup Environ Med* 2001;58:528–34.
- [108] Moore A, Jull G. The systematic review of systematic reviews has arrived. *Man Ther* 2006;11:91–2.
- [109] Muramatsu N, Liang J, Sugisawa H. Transitions in chronic low back pain in Japanese older adults: a sociomedical perspective. *J Gerontol B Psychol Sci Soc Sci* 1997;52:S222–34.
- [110] Newton M, Thow M, Somerville D, Henderson I, Waddell G. Trunk strength testing with iso-machines. Part 2: experimental evaluation of the Cybex II Back Testing System in normal subjects and patients with chronic low back pain. *Spine* 1993;18:812–24.
- [111] Niedhammer I, Lert F, Marne MJ. Back pain and associated factors in French nurses. *Int Arch Occup Environ Health* 1994;66:349–57.
- [112] Nordin M, Skovron ML, Hiebert R. Early predictors of delayed return to work in patients with low back pain. *J Musculoskelet Pain* 1997;5:2–27.
- [113] Norlander S, Gustavsson BA, Lindell J, Nordgren B. Reduced mobility in the cervico-thoracic motion segment—a risk factor for musculoskeletal neck-shoulder pain: a two-year prospective follow-up study. *Scand J Rehabil Med* 1997;29:167–74.
- [114] Norris SH, Watt I. The prognosis of neck injuries resulting from rear-end vehicle collisions. *J Bone Joint Surg Br* 1983;65:608–11.
- [115] Obelieniene D, Schrader H, Bovim G, Miseviciene I, Sand T. Pain after whiplash: a prospective controlled inception cohort study. *J Neurol Neurosurg Psychiatry* 1999;66:279–83.
- [116] Papageorgiou AC, Macfarlane GJ, Thomas E, Croft PR, Jayson MI, Silman AJ. Psychosocial factors in the workplace – do they predict new episodes of low back pain? Evidence from the South Manchester Back Pain Study. *Spine* 1997;22:1137–42.
- [117] Partheni M, Constantoyannis C, Ferrari R, Nikiforidis G, Voulgaris S, Papadakis N. A prospective cohort study of the outcome of acute whiplash injury in Greece. *Clin Exp Rheumatol* 2000;18:67–70.
- [118] Pearce JM. Headaches in the whiplash syndrome. *Spinal Cord* 2001;39:228–33.
- [119] Pettersson K, Hildingsson C, Toolanen G, Fagerlund M, Bjornebrink J. Disc pathology after whiplash injury. A prospective magnetic resonance imaging and clinical investigation. *Spine* 1997;22:283–7.
- [120] Picavet HS, Vlaeyen JW, Schouten JS. Pain catastrophizing and kinesiophobia: predictors of chronic low back pain. *Am J Epidemiol* 2002;156:1028–34.
- [121] Picavet HS, Schouten JS. Musculoskeletal pain in the Netherlands: prevalences, consequences and risk groups, the DMC(3)-study. *Pain* 2003;102:167–78.
- [122] Pincus T, Vogel S, Burton AK, Santos R, Field AP. Fear avoidance and prognosis in back pain: a systematic review and synthesis of current evidence. *Arthritis Rheum* 2006;54:3999–4010.
- [123] Pincus T, Santos R, Breen A, Burton AK, Underwood M. A review and proposal for a core set of factors for prospective cohorts in low back pain: a consensus statement. *Arthritis Rheum* 2008;59:14–24.
- [124] Poiraudau S, Rannou F, Le HA, Coudeyre E, Rozenberg S, Huas D, Martineau C, Jolivet-Landreau I, Revel M, Ravaud P. Outcome of subacute low back pain: influence of patients' and rheumatologists' characteristics. *Rheumatology* 2006;45:718–23.
- [125] Radanov BP, Di SG, Schnidrig A, Ballinari P. Role of psychosocial stress in recovery from common whiplash. *Lancet* 1991;338:712–5.

- [126] Radanov BP, Sturzenegger M, Di SG, Schnidrig A, Aljinovic M. Factors influencing recovery from headache after common whiplash. *BMJ* 1993;307:652–5.
- [127] Radanov BP, Sturzenegger M, Di SG, Schnidrig A, Mumenthaler M. Results of a 1-year follow-up study of whiplash injury. *Schweiz Med Wochenschr* 1993;123:1545–52.
- [128] Radanov BP, Di SG, Schnidrig A, Sturzenegger M, Augustiny KF. Cognitive functioning after common whiplash. A controlled follow-up study. *Arch Neurol* 1993;50:87–91.
- [129] Radanov BP, Di SG, Schnidrig A, Sturzenegger M. Common whiplash: psychosomatic or somatopsychic? *J Neurol Neurosurg Psychiatry* 1994;57:486–90.
- [130] Radanov BP, Sturzenegger M, De SG, Schnidrig A. Relationship between early somatic, radiological, cognitive and psychosocial findings and outcome during a one-year follow-up in 117 patients suffering from common whiplash. *Br J Rheumatol* 1994;33:442–8.
- [131] Radanov BP, Sturzenegger M, Di SG. Long-term outcome after whiplash injury. A 2-year follow-up considering features of injury mechanism and somatic, radiologic, and psychosocial findings. *Medicine* 1995;74:281–97.
- [132] Ready AE, Boreskie SL, Law SA, Russell R. Fitness and lifestyle parameters fail to predict back injuries in nurses. *Can J Appl Physiol* 1993;18:80–90.
- [133] Riihimäki H, Viikari-Juntura E, Moneta G, Kuha J, Videman T, Tola S. Incidence of sciatic pain among men in machine operating, dynamic physical work, and sedentary work. A three-year follow-up. *Spine* 1994;19:138–42.
- [134] Riihimäki H, Wickstrom G, Hanninen K, Luopajarvi T. Predictors of sciatic pain among concrete reinforcement workers and house painters – a five-year follow-up. *Scand J Work Environ Health* 1989;15:415–23.
- [135] Rissanen A, Heliovaara M, Alaranta H, Taimela S, Malkia E, Knekt P, Reunanen A, Aromaa A. Does good trunk extensor performance protect against back-related work disability? *J Rehabil Med* 2002;34:62–6.
- [136] Rossignol M, Lortie M, Ledoux E. Comparison of spinal health indicators in predicting spinal status in a 1-year longitudinal study. *Spine* 1993;18:54–60.
- [137] Ryan GA, Taylor GW, Moore VM, Dolinis J. Neck strain in car occupants: injury status after 6 months and crash-related factors. *Injury* 1994;25:533–7.
- [138] Sackett DL, Haynes RB, Tugwell P. *Clinical epidemiology a basic science for clinical medicine*. Boston: Little, Brown and Co.; 1985.
- [139] Satoh S, Naito S, Konishi T, Yoshikawa M, Morita N, Okada T, Kageyama T, Matsuzaki I. An examination of reasons for prolonged treatment in Japanese patients with whiplash injuries. *J Musculoskelet Pain* 1997;5:71–84.
- [140] Scholten-Peeters GG, Verhagen AP, Bekkering GE, van der Windt DA, Barnsley L, Oostendorp RA, Hendriks EJ. Prognostic factors of whiplash-associated disorders: a systematic review of prospective cohort studies. *Pain* 2003;104:303–22.
- [141] Schrader H, Obelieniene D, Bovim G, Surkiene D, Mickeviciene D, Miseviciene I, Sand T. Natural evolution of late whiplash syndrome outside the medicolegal context. *Lancet* 1996;347:1207–11.
- [142] Schultz IZ, Crook JM, Berkowitz J, Meloche GR, Milner R, Zuberbier OA, Meloche W. Biopsychosocial multivariate predictive model of occupational low back disability. *Spine* 2002;27:2720–5.
- [143] Shannon HS, Woodward CA, Cunningham CE, McIntosh J, Lendrum B, Brown J, Rosenbloom D. Changes in general health and musculoskeletal outcomes in the workforce of a hospital undergoing rapid change: a longitudinal study. *J Occup Health Psychol* 2001;6:3–14.
- [144] Sieben JM, Vlaeyen JW, Tuerlinckx S, Portegijs PJ. Pain-related fear in acute low back pain: the first two weeks of a new episode. *Eur J Pain* 2002;6:229–37.
- [145] Sieben JM, Vlaeyen JW, Portegijs PJ, Verbunt JA, van Riet-Rutgers S, Kester AD, Von KM, Arntz A, Knottnerus JA. A longitudinal study on the predictive validity of the fear-avoidance model in low back pain. *Pain* 2005;117:162–70.
- [146] Sterner Y, Toolanen G, Knibestol M, Gerdle B, Hildingsson C. Prospective study of trigeminal sensibility after whiplash trauma. *J Spinal Disord* 2001;14:479–86.
- [147] Stevenson JM, Weber CL, Smith JT, Dumas GA, Albert WJ. A longitudinal study of the development of low back pain in an industrial population. *Spine* 2001;26:1370–7.
- [148] Sturzenegger M, Radanov BP, Di SG. The effect of accident mechanisms and initial findings on the long-term course of whiplash injury. *J Neurol* 1995;242:443–9.
- [149] Takala EP, Viikari-Juntura E. Do functional tests predict low back pain? *Spine* 2000;25:2126–32.
- [150] Torp S, Riise T, Moen BE. The impact of psychosocial work factors on musculoskeletal pain: a prospective study. *J Occup Environ Med* 2001;43:120–6.
- [151] Tousignant M, Rossignol M, Goulet L, Dassa C. Occupational disability related to back pain: application of a theoretical model of work disability using prospective cohorts of manual workers. *Am J Ind Med* 2000;37:410–22.
- [152] Troup JD, Foreman TK, Baxter CE, Brown D. 1987 Volvo award in clinical sciences. The perception of back pain and the role of psychophysical tests of lifting capacity. *Spine* 1987;12:645–57.
- [153] Tubach F, Leclerc A, Landre MF, Pietri-Taleb F. Risk factors for sick leave due to low back pain: a prospective study. *J Occup Environ Med* 2002;44:451–8.
- [154] van der Giesen AM, Bouter LM, Nijhuis FJ. Prediction of return-to-work of low back pain patients sicklisted for 3–4 months. *Pain* 2000;87:285–94.
- [155] van der Weide WE, Verbeek JH, Salle HJ, van Dijk FJ. Prognostic factors for chronic disability from acute low-back pain in occupational health care. *Scand J Work Environ Health* 1999;25:50–6.
- [156] van der Windt DA, Koes BW, Boeke AJ, Deville W, de Jong BA, Bouter LM. Shoulder disorders in general practice. prognostic indicators of outcome. *Br J Gen Pract* 1996;46:519–23.
- [157] van Poppel MN, Koes BW, Deville W, Smid T, Bouter LM. Risk factors for back pain incidence in industry: a prospective study. *Pain* 1998;77:81–6.
- [158] Viikari-Juntura E, Shiri R, Solovieva S, Karppinen J, Leino-Arjas P, Varonen H, Kalso E, Ukkola O. Risk factors of atherosclerosis and shoulder pain—is there an association? A systematic review. *Eur J Pain* 2008;12:412–26.
- [159] Viikari-Juntura E, Riihimäki H, Takala EP, Rauas S, Leppänen A, Malmivaara A. Factors predicting pain in the neck, shoulders and upper limbs in forestry work. *People Work Res Rep* 1993;7:223–53.
- [160] Viikari-Juntura E, Vuori J, Silverstein BA, Kalimo R, Kuosma E, Videman T. A life-long prospective study on the role of psychosocial factors in neck-shoulder and low-back pain. *Spine* 1991;16:1056–61.
- [161] Waddell G, Burton AK. Occupational health guidelines for the management of low back pain at work: evidence review. *Occup Med* 2001;51:124–35.
- [162] Waddell G. Preventing incapacity in people with musculoskeletal disorders. *Br Med Bull* 2006;77–78:55–69.
- [163] Warren RA, Warren MA. Shiplash injury sustained in motor vehicle accidents: factors influencing time of work. *J Orthop Med* 2001;23:50–4.
- [164] Wempe KM, Rosvall M. Work related and non-work related stress in relation to low leisure time physical activity in a Swedish population. *J Epidemiol Community Health* 2005;59:377–9.
- [165] Werneke M, Hart DL. Centralization phenomenon as a prognostic factor for chronic low back pain and disability. *Spine* 2001;26:758–64.
- [166] Wickstrom GJ, Pentti J. Occupational factors affecting sick leave attributed to low-back pain. *Scand J Work Environ Health* 1998;24:145–52.
- [167] Williams RA, Pruitt SD, Doctor JN, Epping-Jordan JE, Wahlgren DR, Grant I, Patterson TL, Webster JS, Slater MA, Atkinson JH. The contribution of job satisfaction to the transition from acute to chronic low back pain. *Arch Phys Med Rehabil* 1998;79:366–74.
- [168] World Health Organization. International classification of functioning, disability and health: ICF. Geneva: World Health Organization; 2001.
- [169] Wu B, Porell F. Job characteristics and leisure physical activity. *J Aging Health* 2000;12:538–59.
- [170] Yip YB. The association between psychosocial work factor and future low back pain among nurses in Hong Kong: a prospective study. *Psychol Health Med* 2002;7:223–33.