University of Groningen

The Resolution of 2-Hydroxy-5,5-dimethyl-4-phenyl-1,3,2-dioxaphosphorinan 2-Oxide (Phencyphos) by Preferential Crystallization

Leeman, Michel; Querniard, Florian; Kaptein, Bernard; Kellogg, Richard M.

Published in:
Organic Process Research & Development

DOI:
[10.1021/op900171k](https://doi.org/10.1021/op900171k)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Final author's version (accepted by publisher, after peer review)

Publication date:
2009

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Leeman, M., Querniard, F., Kaptein, B., & Kellogg, R. M. (2009). The Resolution of 2-Hydroxy-5,5-dimethyl-4-phenyl-1,3,2-dioxaphosphorinan 2-Oxide (Phencyphos) by Preferential Crystallization. *Organic Process Research & Development*, 13(6), 1379-1381. <https://doi.org/10.1021/op900171k>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

The Resolution of 2-Hydroxy-5,5-dimethyl-4-phenyl-1,3,2-dioxaphosphorinan 2-oxide (Phencyphos) by Preferential Crystallization.

Michel Leeman, Florian Querniard, Bernard Kaptein and Richard M. Kellogg

Methods

Materials

All chemicals were obtained from commercial sources and used without further purification.

General Information

Chiral HPLC analysis of phencyphos was carried out on a Chiralpak QN-AX column with MeOH:AcOH 97:3 + 0.25g NH₄OAc/100 mL as mobile phase at room temperature and 1.5 mL·min⁻¹. UV-vis detection was performed at 254 nm. The phencyphos was dissolved in MeOH and injected as such. (+)-(S)-phencyphos R_f: 7.34 min, (-)-(R)-phencyphos R_f: 8.44 min.

IR spectra were recorded on a Thermo Nicolet 380 FT-IR with Smart Orbit.

Procedure for the construction of the ternary phase diagrams

Just enough (±)-phencyphos·0.84H₂O to achieve a suspension, was slurried in known mixtures of water and DMSO, DMF or MeOH for a couple of days at 20°C. After removal of the solid phase the composition of the mother liquor was determined by weighing the weight loss after evaporation of the solvents *in vacuo*.

Determination of the IR spectra of (±)-phencyphos

Solvates of (±)-phencyphos with either MeOH, EtOH or IPA should give different IR signatures. (±)-phencyphos was recrystallized from these solvents and the solids were analyzed by FT-IR. However, the signatures were similar which indicates that MeOH, EtOH and IPA do not give solvates with (±)-phencyphos as depicted in Figure 1. The recrystallization of (±)-phencyphos from water delivered the monohydrate which, indeed, has a different spectrum (best visible between 1050 and 1250 cm⁻¹) than the (±)-phencyphos recrystallized from MeOH, EtOH or IPA.

Figure 1 IR spectra from (±)-phencyphos crystallized from several solvents. The peaks at 2350 cm⁻¹ are from CO₂.

Determination of the solubility of (±)-phencyphos

The solubility of (±)-phencyphos in several pure solvents as described in the main article was performed in a similar fashion to the procedure described above for the ternary phase diagrams and the results are given in Table 1.

Table 1 Solubilities of (±)-phencyphos in several water miscible solvents.

Entry	Solvent	Solubility (mg (±)- phencyphos/mL solvent)
1	MeCN	< 1
2	1,4-dioxane	1.1
3	water	3.2
4	ethylene glycol	4.0
5	acetic acid	4.8
6	1-butanol	5.7
7	sec-butanol	5.8
8	IPA	8.4
9	MeOH	17
10	DMF	81
11	DMSO	142

Procedure for the preferential crystallization of phencyphos hydrate on 35L scale

A temperature controlled (Huber Unistat 510) 40L double jacketed glass reactor was charged with (±)-phencyphos·0.84H₂O (861 g, 3.35 mol, 1.0 eq), water (9.25 L) and MeOH (27.75 L) and stirred with a propeller shaped stirrer at 300 rpm throughout the preferential crystallization.

The mixture was biased with (–)-phencyphos hydrate (137 g, 0.52 mol, 0.16 eq) and a temperature program was run where the internal temperature was heated from 28.9°C to 54.0°C as fast as possible (41 minutes) and kept at this temperature for another 5 minutes. Complete dissolution was observed. Then the mixture was cooled to 28.9°C as fast as possible (44 minutes) and kept at this temperature for another 37 minutes. Then the temperature program was repeated. The temperature profile is depicted in Figure 2.

The solids were collected (P2) 10 minutes prior to the reheating to 54.0°C. The filter cakes were sucked dry but not washed. Chiral HPLC analyses (as described above) on the solids were carried out. If the solids had an *ee* greater than 85%, (±)-phencyphos·H₂O (270 g, 1.04 mol, 0.31 eq) was added to the filtrate. If the collected solids had 0% *ee* anhydrous phencyphos was isolated and thus, more water was added and the preferential crystallization repeated by reintroducing the filter cake in the reactor. If the collected solids had an *ee* between 30% and 85%, the concentration of the undesired enantiomer was too high and the preferential crystallization was repeated with the filter cake and replacing some filtrate by MeOH. In a typical run, solids were collected with an average *ee* of 93%. Combined enriched batches were heated and subsequently cooled in a mixture of 30% wt. H₂O in MeOH to give crystals with >99% *ee* after filtration. In practice the whole resolution process yielded 41% of each enantiomer of optically pure phencyphos hydrate.

Figure 2 Temperature program for the resolution by preferential crystallization of phencyphos hydrate on 35L scale. The program repeats itself after 127 min.