University of Groningen

Effect of nodule characteristics on variability of semiautomated volume measurements in pulmonary nodules detected in a lung cancer screening program

Wang, Y.; van Klaveren, R.J.; van der Zaag-Loonen, H.J.; de Bock, G.H.; Gietema, H.A.; Xu, D.M.; Leusveld, A.L.; de Koning, H.J.; Scholten, E.T.; Verschakelen, J.

Published in:
Radiology

DOI:
[10.1148/radiol.2482070957](https://doi.org/10.1148/radiol.2482070957)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2008

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Wang, Y., van Klaveren, R. J., van der Zaag-Loonen, H. J., de Bock, G. H., Gietema, H. A., Xu, D. M., Leusveld, A. L., de Koning, H. J., Scholten, E. T., Verschakelen, J., Prokop, M., & Oudkerk, M. (2008). Effect of nodule characteristics on variability of semiautomated volume measurements in pulmonary nodules detected in a lung cancer screening program. *Radiology*, 248(2), 625-631.
<https://doi.org/10.1148/radiol.2482070957>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Effect of Nodule Characteristics on Variability of Semiautomated Volume Measurements in Pulmonary Nodules Detected in a Lung Cancer Screening Program¹

Ying Wang, MD
 Rob J. van Klaveren, MD, PhD
 Hester J. van der Zaag-Loonen, MD, PhD
 Geertruida H. de Bock, PhD
 Hester A. Gietema, MD, PhD
 Dong Ming Xu, MD, PhD
 Anne L. M. Leusveld, MD
 Harry J. de Koning, MD, PhD
 Ernst T. Scholten, MD
 Johnny Verschakelen, MD, PhD
 Mathias Prokop, MD, PhD
 Matthijs Oudkerk, MD, PhD

¹ From the Departments of Radiology (Y.W., H.J.v.d.Z., D.M.X., A.L.M.L., M.O.) and Epidemiology (G.H.d.B.), University Medical Center Groningen, University of Groningen, Hanzeplein 1, 9700 RB Groningen, the Netherlands; Departments of Pulmonology (R.J.v.K.) and Public Health (H.J.d.K.), Erasmus Medical Center, Rotterdam, the Netherlands; Department of Radiology, University of Utrecht Medical Center, Utrecht, the Netherlands (H.A.G., M.P.); Department of Radiology, Kennemer Gasthuis, Haarlem, the Netherlands (E.T.S.); and Department of Radiology, University Hospital Gasthuisberg, Leuven, Belgium (J.V.). From the 2006 RSNA Annual Meeting. Received June 4, 2007; revision requested August 6; revision received January 10, 2008; final version accepted February 21, 2008.

Address correspondence to M.O. (e-mail: m.oudkerk@rad.umcg.nl).

© RSNA, 2008

Purpose:

To retrospectively assess volume measurement variability in solid pulmonary nodules (volume, 15–500 mm³) detected at lung cancer screening and to quantify the independent effects of nodule morphology, size, and location.

Materials and Methods:

This retrospective study was a substudy of the screening program that was approved by the Dutch Ministry of Health, and all participants provided written informed consent. Two independent readers used semiautomated software to measure the volume of pulmonary nodules detected in 6774 participants aged 50–75 years (5917 men). Nodules were classified according to their location (purely intraparenchymal, pleural based, juxtavascular, or fissure attached), morphology (smooth, polylobulated, spiculated, or irregular), and size (≤ 50 mm³ or > 50 mm³). The level of agreement was expressed by using the absolute values of the relative volume differences (RVDs). Multivariate logistic regression analysis was performed, and odds ratios (ORs) were computed to quantify the independent effects of morphology, location, and size on RVD categories.

Results:

Altogether, 4225 nodules in 2239 participants were included. Complete agreement in volume was obtained for 3646 (86%) of the nodules. Disagreement was small (absolute value of RVD $< 5\%$) for 173 (4%) nodules, moderate (absolute value of RVD $\geq 5\%$ but $< 15\%$) for 232 (6%), and large (absolute value of RVD $\geq 15\%$) for 174 (4%). Multivariate analysis showed that the ORs of volume disagreement were 15.7, 3.1, and 1.9 for irregular, spiculated, and polylobulated nodules, respectively; 3.5, 2.6, and 2.1 for juxtavascular, pleural-based, and fissure-attached nodules, respectively; and 1.3 for large nodules compared with smooth, purely intraparenchymal, and small reference nodules.

Conclusion:

Nodule morphology, location, and size influence volume measurement variability, particularly for juxtavascular and irregular nodules.

© RSNA, 2008

With the introduction of multidetector computed tomography (CT) and the ongoing advances in multidetector CT technology, the detection of small pulmonary nodules has become common, with detection rates ranging from 23% to 74% in lung cancer CT screening studies (1–5). Today, the challenge is to identify the malignant nodules among the majority of benign nodules. Volume doubling time has received attention as a diagnostic tool with which to differentiate malignant nodules from benign nodules, especially subcentimeter lesions in which additional diagnostic techniques—such as positron emission tomography, contrast material-enhanced CT, and percutaneous needle biopsy—are less suitable (6–8). Malignant solid pulmonary nodules typically have volume doubling times of 20–400 days, whereas benign nodules usually have volume doubling times of more than 400 days (9–14).

The value of volume doubling time in clinical practice is, however, largely dependent on measurement variability. Two-dimensional volumetric assessments have been found to be unreliable in the detection of volume changes (15,16), while the results obtained with computer-aided three-dimensional (3D) volumetric software have been found to be superior to the results obtained with two-dimensional software (12,17–22).

Advances in Knowledge

- Complete agreement in software-generated volume measurement is dependent on nodule morphology and location and varies from 91% for purely intraparenchymal nodules to 70% for juxtavascular nodules and from 90% for smooth nodules to 34% for irregular nodules.
- Compared with smooth, small, and purely intraparenchymal nodules, irregular and juxtavascular nodules have the largest volume variability; in spiculated, pleural-based, and fissure-attached nodules, variability is moderate; and in lobulated and larger (>50 mm³) nodules, variability is small.

In a retrospective study, Revel et al (18) reported that software-generated volumetric analyses yielded similar results in nine repeated measurements in 35 (67%) of 52 nodules. Gietema et al (23) found complete agreement for 89% of the 430 solid 15–500-mm³ nodules that were situated entirely within the lung parenchyma. However, not all nodules are purely intraparenchymal. They may also be pleural based, juxtavascular, or fissure attached, and these different locations may influence the accuracy and variability of volumetric measurements. Furthermore, nodule morphology may influence volume measurement variability (23,24).

To our knowledge, quantitative assessment of the independent effects of nodule morphology, size, and location on volume measurement variability has not been performed. Thus, the purpose of our study was to retrospectively assess volume measurement variability in solid pulmonary nodules (volume, 15–500 mm³) detected at lung cancer screening and to quantify the independent effects of nodule morphology, size, and location.

Materials and Methods

Study Group

Between April 2004 and December 2005, 6774 participants underwent baseline low-dose multidetector CT screening for lung cancer. Participants (5917 men, 857 women) were current or former smokers aged 50–75 years with a history of smoking more than 15 cigarettes per day for more than 25 years or more than 10 cigarettes per day for more than 30 years. The prospective screening study was approved by the Dutch Ministry of Health and the medi-

cal ethical board at each of the four participating hospitals, and written informed consent was obtained from all participants. Our current retrospective study was performed by using CT data acquired in the prospective screening study. The original approval and informed consent for the prospective study included permission to use data for future retrospective research.

Imaging

All examinations were performed by using 16-detector helical CT scanners (Sensation 16, Siemens Medical Systems, Forchheim, Germany; Mx8000 IDT or Brilliance 16P, Philips Medical Systems, Cleveland, Ohio), a low radiation dose, and the following parameters: 0.5-second tube rotation, 0.75-mm single-section collimation, 15- or 18-mm table feed per rotation, and 1.3–1.5 pitch. Scanning was performed in a caudocranial direction, without the use of contrast material. Images were obtained from the level of the lung bases (posterior recesses) to the level of the lung apex with the help of a scout view. Exposure settings (20–30 mAs, 100–140 kVp) depended on the weight of the participant. These settings corresponded to an effective radiation dose of less than 1.6 mSv. We reconstructed axial images with a 1.0-mm thickness at 0.7-mm in-

Published online

10.1148/radiol.2482070957

Radiology 2008; 248:625–631

Abbreviations:

CI = confidence interval
OR = odds ratio
RVD = relative volume difference
3D = three-dimensional

Author contributions:

Guarantors of integrity of entire study, Y.W., R.J.v.K., H.J.v.d.Z., J.V., M.O.; study concepts/study design or data acquisition or data analysis/interpretation, all authors; manuscript drafting or manuscript revision for important intellectual content, all authors; manuscript final version approval, all authors; literature research, Y.W., R.J.v.K., H.J.v.d.Z., H.A.G.; clinical studies, Y.W., R.J.v.K., H.J.v.d.Z., H.A.G., D.M.X., A.L.M.L., E.T.S., J.V., M.P.; statistical analysis, Y.W., H.J.v.d.Z., G.H.d.B., D.M.X.; and manuscript editing, Y.W., R.J.v.K., H.J.v.d.Z., H.A.G., D.M.X., H.J.d.K., J.V., M.P., M.O.

Authors stated no financial relationship to disclose.

Implication for Patient Care

- Visual verification of growth, in addition to volumetric measurement, is recommended for pleural-based, fissure-attached, or juxtavascular nodules with ill-defined margins to avoid false-positive results.

crements. A moderately sharp kernel was used for reconstruction (B30 filter, Siemens Medical Solutions, Erlangen, Germany; B filter, Philips Medical Systems). All images were reconstructed with a field of view large enough to cover the complete lung cross section.

Nodule Detection and Volume Measurement

For nodule detection and volume measurement, digital workstations (Leonardo; Siemens Medical Systems) were used with Food and Drug Administration–approved commercially available software to obtain semiautomated volume measurements (LungCare; Siemens Medical Solutions) at all four screening sites. Pulmonary nodules were identified on axial thin-slab maximum intensity projections with the cine mode at standardized window width and window center settings of 1500 and –500 HU, respectively, and a slab thickness of 6 or 8 mm. After the nodule was marked with a mouse click, the program automatically defined the volume of interest around the nodule. A 3D template that represented the nodule in its most optimal way was generated, without inclusion of surrounding structures, if possible.

A second mouse click initiated the automated volume measurement, which included the following steps (22): First, a fixed attenuation threshold of –400 HU was applied to extract a 3D-connected structure of interest. This structure may have included surrounding structures, such as vessels or parts of the chest wall. Apart from this process, a small spherical 3D template that originated from the click point was gradually expanded, and its cross correlation with the segmented nodule was computed for each step. The peak value of the cross-correlation curve was determined, and an empirical cutoff value close to the peak value was used to separate the nodule from its potential adjacent structures. Finally, segmentation was completed by fusing the optimal 3D template and the structure of interest. The segmented nodule was then shown in yellow on the volume-rendered display as the volume of interest. If an ob-

server disagreed with the fit of the 3D template with the volume of interest, he or she could manually modify the segmentation by increasing or decreasing the volume. This could have introduced bias in the estimation of volume measurement variability; therefore, if readers manually modified the volume in the prospective lung cancer screening study, the readings and nodules they evaluated were excluded from our retrospective study.

Nodule Evaluation

In the prospective lung cancer screening program, which was based on the study protocol, all solid nodules larger than 15 mm³ were prospectively evaluated and entered into our central database by a local reader and a second central reader working independently (25). The 13 local readers had 1–20 years (median, 6 years) experience reading thoracic CT images. There were two central readers, each of whom had 6 years of experience. All readers measured the volume with the same software and recorded the nodule characteristics independently.

The nodules were classified into the following four subgroups on the basis of location: The first subgroup consisted of purely intraparenchymal nodules that did not have contact with the pleura, vessels, or fissures or that had contact with less than 50% of the nodule diameter (average of long and short axis). The second subgroup consisted of pleural-based nodules that had pleural contact with more than 50% of the nodule diameter. The third subgroup consisted of fissure-attached nodules that had fissural contact with more than 50% of the nodule diameter. The fourth subgroup consisted of juxtavascular nodules that had vessel contact with more than 50% of the nodule diameter.

Nodules were classified as smooth, polylobulated, spiculated, or irregular on the basis of morphologic findings (26). Lobulation was defined as an abrupt bulging of the lesion contour (27). Spiculation was defined as the presence of thicker strands extending from the nodule margin into the lung parenchyma without reaching the pleu-

ral surface (28). A nodule was deemed irregular if it was not smooth, polylobulated, or spiculated. Nodules were further classified as small (15–49 mm³; effective diameter, 3.1–4.6 mm) or large (50–500 mm³; effective diameter, 4.7–9.8 mm), corresponding to two nodule categories (insignificantly small and indeterminate) in our screening program (25). In our retrospective study, we retrieved all information on solid nodules, with the exception of complete and central calcified nodules between 15 and 500 mm³ (effective diameter, 3.1–9.8 mm) detected by both readers in our central database. Nodules larger than 500 mm³ were excluded because the software we used was not developed for use with these larger nodules. We evaluated all CT images. Because two of the 13 local readers manually modified the volume, all of their readings were excluded from our retrospective study; thus, there were 11 local readers. In cases of disagreement between the first and second readers on nodule characteristics, two readers (D.M.Y., Y.W.) retrospectively reevaluated the images to reach consensus.

Statistical Analysis

Because the true nodule volume was unknown, we used the Bland-Altman method (29) to estimate volume measurement variability. The difference between the first and second volume measurements was computed for all nodules and was defined as the absolute volume difference. The relative volume difference (RVD) was defined as the absolute volume difference divided by the mean volume of the two measurements. Volume measurement variability was assessed for both absolute volume difference and RVD, and means and 95% confidence intervals (CIs) were calculated. Thereafter, the absolute value of RVD was calculated because the percentage volume change is directly related to growth (30). Disagreement was defined as small when the absolute value of RVD was less than 5%, moderate when it was between 5% and 15%, and large when it was more than 15%. The percentage of nodules with volume disagreement of more than 25% was com-

puted because 25% was used as the growth criterion in our lung cancer screening program.

To evaluate the influence of nodule characteristics on measurement variability, we performed univariate logistic regression analysis by using volume disagreement as a dependent variable and by using location, morphology, and size as independent variables. Volume disagreement was defined as an absolute value of RVD that did not equal zero, whereas volume agreement was defined as an absolute value of RVD that equaled zero, and odds ratios (ORs) and 95% CIs were computed. In the next step, the independent variables that contributed significantly in the univariate analyses were included in the multivariate logistic regression analysis. Again, ORs with 95% CIs were calculated. Because the definition of volume agreement was rather strict, the following factors were also considered: (a) whether a nodule had a disagreement of more than 5% (absolute value of RVD > 5% or absolute value of RVD ≤ 5%) and (b) whether a nodule had a disagreement of more than 15% (absolute value of RVD > 15% or absolute value of RVD ≤ 15%). All analyses were performed with statistical software (SPSS, version 14.0.1; SPSS, Chi-

cago, Ill). $P < .05$ indicated a significant difference.

Results

Of the 6774 participants who underwent baseline screening, 2367 had 4477 solid noncalcified nodules with a volume of 15–500 mm³ detected by both readers. Of these nodules, 252 were excluded because they were evaluated by readers who manually changed the software-generated volume. Thus, we included 4225 nodules in 2239 participants in our analyses. The median volume was 40 mm³ (range, 15–485 mm³). The majority of nodules were purely intraparenchymal (68%, $n = 2853$), had a smooth outer contour (80%, $n = 3375$), and were small (62%, $n = 2603$) (Table 1).

The mean absolute volume difference for all nodules was 0.6 mm³ ± 9.1 (standard deviation) (95% CI: −17.2 mm³, 18.4 mm³) (Fig 1a), and the mean RVD was 0.5% (95% CI: −13.4%, 14.5%) (Fig 1b). Complete volume agreement between the two readers was recorded in 3646 (86%) nodules. Volume disagreement was small (absolute value of RVD < 5%) for 173 (4%) nodules, moderate (absolute value of RVD ≥ 5% but < 15%) for 232 (6%),

and large (absolute value of RVD ≥ 15%) for 174 (4%). For 75 nodules (2% [95% CI: 1.4%, 2.3%]), the positive value of RVD was larger than 25%.

All three characteristics contributed significantly to the volume disagreement ($P < .001$) seen at univariate analysis (Table 2). In multivariate analysis, with volume disagreement as a dependent variable, nodule morphology and location remained highly significant ($P < .001$) contributors to disagreement, while nodule size just reached significance ($P = .026$). With disagreement of more than 5% and disagreement of more than 15% as dependent variables, size was not a significant predictor ($P = .611$ and $P = .194$, respectively), whereas morphology and location were significant predictors ($P < .001$ for both) (Table 3).

In multivariate regression analysis, irregular nodules (Fig 2) had ORs of 15.7 (95% CI: 11.0, 22.2), 12.5 (95% CI: 8.8, 17.7), and 9.6 (95% CI: 6.1, 15.1) for volume discrepancies of more than 0%, more than 5%, and more than 15%, respectively, compared with the smooth reference category. Juxtavascular nodules (Fig 3) had ORs for volume disagreement of 3.5 (95% CI: 2.4, 5.0), 5.5 (95% CI: 3.8, 8.0), and 4.6 (95% CI: 2.8, 7.6), respectively, compared with the purely intraparenchymal reference category. Pleural-based, fissure-attached, spiculated, lobulated, and larger nodules had lower ORs compared with the corresponding reference category of nodules (Table 3).

Table 1

RVD and Absolute Volume Difference according to Location, Morphology, and Size in 4225 Solid Pulmonary Nodules Detected at Baseline Screening for Lung Cancer

Nodule Characteristic	No. of Nodules*	Median Volume (mm ³)	RVD (%)†	Absolute Volume Difference (mm ³)†
Location				
Purely intraparenchymal	2853 (68)	36	0.4 ± 5.4	0.5 ± 6.2
Pleural based	717 (17)	50	1.3 ± 9.4	1.4 ± 15.2
Juxtavascular	199 (5)	68	0 ± 12.5	0 ± 14.2
Fissure attached	456 (11)	46	0.2 ± 8.9	0.2 ± 7.9
Morphology				
Smooth	3375 (80)	36	0.4 ± 5.1	0.2 ± 4.6
Lobulated	611 (14)	64	1.4 ± 1.1	2.1 ± 17.4
Spiculated	68 (2)	181	0 ± 6.0	−0.7 ± 7.8
Irregular	171 (4)	40	2.0 ± 16.8	3.6 ± 21.9
Size				
Small	2603 (62)	28	0.2 ± 5.1	0.7 ± 1.6
Large	1622 (38)	78	0.4 ± 9.5	1.4 ± 14.4

* Data in parentheses are percentages.

† Data are means ± standard deviations.

Discussion

Our results show that in 86% of the solid nodules detected in a lung cancer screening trial, complete volume agreement between two readers can be achieved by using semiautomated volumetric software. Large volume discrepancies (absolute value of RVD ≥ 25%) occurred in only 2% of nodules (95% CI: 1.4%, 2.3%). An increase in volume of at least 25% within a 3-month interval was defined as growth in one screening trial (25), and it might have led to a false-positive test result at 3-month follow-up in 2% of the pulmonary nodules

Figure 1

Figure 1: Bland-Altman plot for 4225 nodules shows the (a) absolute volume difference and (b) RVD between the two observers against mean nodule volume. The solid line indicates mean difference, and the dashed lines indicate 95% CIs. Although most nodules had volume disagreement close to zero, a small percentage of nodules had large volume disagreement.

Table 2

ORs of Nodule Characteristics for Volume Disagreement at Univariate Analysis

Nodule Characteristics	OR	95% CI
Location*		
Purely intraparenchymal	1	...
Pleural based	2.0	1.6, 2.6
Juxtavascular	4.1	2.9, 5.7
Fissure attached	3.7	2.9, 4.7
Morphology*		
Smooth	1	...
Lobulated	2.3	1.8, 3.0
Spiculated	2.9	1.6, 5.1
Irregular	18.1	13.0, 25.4
Size*		
Small	1	...
Large	1.7	1.4, 2.0

* $P < .001$. Comparisons were made within each group.

Table 3

Multivariate Logistic Regression Analysis of the Relationships of Morphology, Location, and Size with Volume Variability

Nodule Characteristic	Positive Value of RVD Greater than 0%	Positive Value of RVD Greater than 5%	Positive Value of RVD Greater than 15%
Morphology*			
Smooth	1	1	1
Polylobulated	1.9 (1.5, 2.4)	1.7 (1.3, 2.3)	2.6 (1.7, 3.8)
Spiculated	3.1 (1.7, 5.6)	2.2 (1.0, 4.7)	2.5 (0.9, 7.2)
Irregular	15.7 (11.0, 22.2)	12.5 (8.8, 17.7)	9.6 (6.1, 15.1)
Location*			
Purely intraparenchymal	1	1	1
Pleural based	2.1 (1.7, 2.7)	2.4 (1.8, 3.2)	2.5 (1.7, 3.7)
Fissure attached	2.6 (2.0, 3.4)	2.7 (2.0, 3.7)	1.6 (1.0, 2.6)
Juxtavascular	3.5 (2.4, 5.0)	5.5 (3.8, 8.0)	4.6 (2.8, 7.6)
Size†			
Small	1	1	1
Large	1.3 (1.0, 1.5)	1.1 (0.8, 1.3)	1.2 (0.9, 1.7)

Note.—Data are ORs, and data in parentheses are 95% CIs.

* Overall P value was less than .001 for positive value of RVD greater than 0%, positive value of RVD greater than 5%, and positive value of RVD greater than 15%.

† Overall P value was less than .05 for positive value of RVD greater than 0% and not significant for positive value of RVD greater than 5% and positive value of RVD greater than 15%.

detected. We also found that irregular and juxtavascular nodules had the largest risk of volume variability; spiculated, fissure-attached, and pleural-based nodules had a moderate risk of volume variability; and lobulated and larger nodules had a small risk of volume

variability. Semiautomated software-generated segmentation should be visually verified and manually adjusted if the observer believes segmentation has not

been successful, especially with irregular and juxtavascular nodules.

Compared with previous studies on volume measurement variability,

our study not only was larger, but it also covered the complete spectrum of solid nodules detected at baseline screening. To our knowledge, ours was also the only study in which the independent effects of nodule characteristics on volume measurement variability were quantified. Gietema et al (23) used a smaller data set with only

purely intraparenchymal nodules. They found complete agreement between two readers regarding a purely intraparenchymal location for 89% of the nodules, which is almost identical to our finding of complete agreement for 91% of the nodules. In the study conducted by Wormanns et al (22), two consecutive CT acquisitions were

performed within a 10-minute interval in 151 nodules (mean diameter, 7.4 mm) in patients with known pulmonary metastases. The interobserver 95% CI was -5.5% to 6.6% , which was less than that in our study. However, no details on the morphology and location of these nodules were provided. As pulmonary metastases tend to be purely intraparenchymal, smooth, and round, the conclusion reached by Wormanns et al (22) might be applicable to only that type of nodule. In the other studies (13,18,21) on volume measurement error published thus far, the number of nodules ranged from 13 to 62. The nodules were usually smaller than 10 mm and incidentally found on routine CT images. Because different volumetric software programs were used, comparison with our results is impossible.

An explanation for the observed volume disagreement between the two readers is related to the fact that semi-automatic volumetric measurements may vary according to the positioning of the seed point by the observer, which is the only nonautomated part of the procedure. When a spherical 3D template gradually expands from the seed point, different starting positions within the nodules may lead to different volumetric results. In nonsmooth nodules, it is difficult to identify the correct borderline, and in nonspherical nodules, the segmentation tends to vary with the seed point owing to the spherical 3D template used in the software program. In pleural-based, fissure-attached, and juxtavascular nodules, both measurement errors occur, whereas in large nodules, the larger disposable seed point area leads to more measurement variability. The software program used in our study provides a postprocessing option for unsatisfactory segmentation. Two (Y.W., D.M.X.) of the original 13 readers in our study were accustomed to changing the segmentation in cases of an unsatisfactory software-generated volume. The nodules these two readers evaluated were excluded because they performed the first reading and their results were not comparable to the results of the second readers, who used only the semiauto-

Figure 2

Figure 2: Volume-rendered CT displays show two volume estimates of an irregular purely intraparenchymal nodule. Nodule volume varied from 197 mm³ (right) to 248 mm³ (left), with a relative volume difference of 22.8%. The segmented nodule (yellow) is shown as the volume of interest.

Figure 3

Figure 3: Volume-rendered CT displays show two volume estimates of a polylobulated juxtavascular nodule. Nodule volume varied from 46 mm³ (right) to 64 mm³ (left), with a relative volume difference of 31.7%. The segmented nodule (yellow) is shown as the volume of interest.

tomatic measurement and could have introduced additional variability. The number of nodules evaluated by these two readers ($n = 252$) was relatively small compared with the total number of nodules ($n = 4477$).

A limitation of our study was that there was no reference standard for the true nodule volume. However, in practice, this is impossible to achieve with noninvasive means. Furthermore, we wanted to assess volume measurement variability between two independent readers. Another limitation of our study was that our results are applicable only for the specific software used, although they may be applicable for other software if identical procedures are used to measure the volume. Furthermore, we evaluated only the variability between two observers and not the variability between two successive CT examinations, which may be considerably larger and relevant to the assessment of volumetric growth over time.

In conclusion, semiautomated software-generated volumetric measurement was completely reproducible in 86% of the solid 15–500-mm³ nodules. Nodule morphology, location, and size influenced volume measurement variability, particularly among the juxtavascular and irregular nodules, in which visual verification of growth is recommended to avoid false-positive results.

Acknowledgments: We thank our colleagues from the Department of Radiology, University Medical Center Groningen, University of Groningen, especially Yingru Zhao, MD, and Fenneke van de Wiel, MD, for their support in the evaluation of the nodules; Dr Estelle J. K. Noach, MD, PhD, for her critical review of the manuscript; and Dr Peter Paul Taverne, MD, Dr Peter van Ooijen, PhD, and Wim Tukker, PhD, for their advice and technical support.

References

- Swensen SJ, Jett JR, Hartman TE, et al. Lung cancer screening with CT: Mayo Clinic experience. *Radiology* 2003;226:756–761.
- Swensen SJ, Jett JR, Hartman TE, et al. CT screening for lung cancer: 5-year prospective experience. *Radiology* 2005;235:259–265.
- Henschke CI, McCauley DI, Yankelevitz DF, et al. Early Lung Cancer Action Project: overall design and findings from baseline screening. *Lancet* 1999;354:99–105.
- Fischbach F, Knollmann F, Griesshaber V, Freund T, Akkol E, Felix R. Detection of pulmonary nodules by multislice computed tomography: improved detection rate with reduced slice thickness. *Eur Radiol* 2003;13:2378–2383.
- Diederich S, Wormanns D, Semik M, et al. Screening for early lung cancer with low-dose spiral CT: prevalence in 817 asymptomatic smokers. *Radiology* 2002;222:773–781.
- Coleman RE, Laymon CM, Turkington TG. FDG imaging of lung nodules: a phantom study comparing SPECT, camera-based PET, and dedicated PET. *Radiology* 1999;210:823–828.
- Li H, Boieselle PM, Shepard JO, Trotman-Dickenson B, McLoud TC. Diagnostic accuracy and safety of CT-guided percutaneous needle aspiration biopsy of the lung: comparison of small and large pulmonary nodules. *AJR Am J Roentgenol* 1996;167:105–109.
- Swensen SJ, Viggiano RW, Midthun DE, et al. Lung nodule enhancement at CT: multicenter study. *Radiology* 2000;214:73–80.
- Lillingston GA. Management of solitary pulmonary nodules. *Dis Mon* 1991;37:271–318.
- Revel MP, Merlin A, Peyrard S, et al. Software volumetric evaluation of doubling times for differentiating benign versus malignant pulmonary nodules. *AJR Am J Roentgenol* 2006;187:135–142.
- Aoki T, Nakata H, Watanabe H, et al. Evolution of peripheral lung adenocarcinomas: CT findings correlated with histology and tumor doubling time. *AJR Am J Roentgenol* 2000;174:763–768.
- Yankelevitz DF, Reeves AP, Kostis WJ, Zhao BS, Henschke CI. Small pulmonary nodules: volumetrically determined growth rates based on CT evaluation. *Radiology* 2000;217:251–256.
- Yankelevitz DF, Gupta R, Zhao B, Henschke CI. Small pulmonary nodules: evaluation with repeat CT—preliminary experience. *Radiology* 1999;212:561–566.
- Hasegawa M, Sone S, Takashima S, et al. Growth rate of small lung cancers detected on mass CT screening. *Br J Radiol* 2000;73:1252–1259.
- Revel MP, Bissery A, Bienvenu M, Aycard L, Lefort C, Fria G. Are two-dimensional CT measurements of small noncalcified pulmonary nodules reliable? *Radiology* 2004;231:453–458.
- Erasmus JJ, Gladish GW, Broemeling L, et al. Interobserver and intraobserver variability in measurement of non-small-cell carcinoma lung lesions: implications for assessment of tumor response. *J Clin Oncol* 2003;21:2574–2582.
- Jennings SG, Winer-Muram HT, Tarver RD, Farber MO. Lung tumor growth: assessment with CT—comparison of diameter and cross-sectional area with volume measurements. *Radiology* 2004;231:866–871.
- Revel MP, Lefort C, Bissery A, et al. Pulmonary nodules: preliminary experience with three-dimensional evaluation. *Radiology* 2004;231:459–466.
- Ko JP, Rusinek H, Jacobs EL, et al. Small pulmonary nodules: volume measurement at chest CT—phantom study. *Radiology* 2003;228:864–870.
- Goo JM, Tongdee T, Tongdee R, Yeo K, Hildebolt CF, Bae KT. Volumetric measurement of synthetic lung nodules with multi-detector row CT: effect of various image reconstruction parameters and segmentation thresholds on measurement accuracy. *Radiology* 2005;235:850–856.
- Goodman LR, Gulsun M, Washington L, Nagy PG, Piacsek KL. Inherent variability of CT lung nodule measurements in vivo using semiautomated volumetric measurements. *AJR Am J Roentgenol* 2006;186:989–994.
- Wormanns D, Kohl G, Klotz E, et al. Volumetric measurements of pulmonary nodules at multi-row detector CT: in vivo reproducibility. *Eur Radiol* 2004;14:86–92.
- Gietema HA, Wang Y, Xu D, et al. Pulmonary nodules detected at lung cancer screening: interobserver variability of semiautomated volume measurements. *Radiology* 2006;241:251–257.
- Petrou M, Quint LE, Nan B, Baker LH. Pulmonary nodule volumetric measurement variability as a function of CT slice thickness and nodule morphology. *AJR Am J Roentgenol* 2007;188:306–312.
- Xu DM, Gietema H, de Koning H, et al. Nodule management protocol of the NELSON randomised lung cancer screening trial. *Lung Cancer* 2006;54:177–184.
- Erasmus JJ, Connolly JE, McAdams HP, Roggli VL. Solitary pulmonary nodules. I. Morphologic evaluation for differentiation of benign and malignant lesions. *RadioGraphics* 2000;20:43–58.
- Takashima S, Maruyama Y, Hasegawa M, et al. CT findings and progression of small peripheral lung neoplasms having a replacement growth pattern. *AJR Am J Roentgenol* 2003;180:817–826.
- Takashima S, Sone S, Li F, et al. Small solitary pulmonary nodules ($\leq$ or $=$ 1 cm) detected at population-based CT screening for lung cancer: reliable high-resolution CT features of 513 benign lesions. *AJR Am J Roentgenol* 2003;180:955–964.
- Bland JM, Altman DG. Measuring agreement in method comparison studies. *Stat Methods Med Res* 1999;8:135–160.
- Kostis WJ, Yankelevitz DF, Reeves AP, Fluttre SC, Henschke CI. Small pulmonary nodules: reproducibility of three-dimensional volumetric measurement and estimation of time to follow-up CT. *Radiology* 2004;231:446–452.

Radiology 2008

This is your reprint order form or pro forma invoice

(Please keep a copy of this document for your records.)

Reprint order forms and purchase orders or prepayments must be received 72 hours after receipt of form either by mail or by fax at 410-820-9765. It is the policy of Cadmus Reprints to issue one invoice per order.

Please print clearly.

Author Name _____
Title of Article _____
Issue of Journal _____ Reprint # _____ Publication Date _____
Number of Pages _____ KB # _____ Symbol Radiology
Color in Article? Yes / No (Please Circle)

Please include the journal name and reprint number or manuscript number on your purchase order or other correspondence.

Order and Shipping Information

Reprint Costs (Please see page 2 of 2 for reprint costs/fees.)

_____ Number of reprints ordered \$ _____

_____ Number of color reprints ordered \$ _____

_____ Number of covers ordered \$ _____

Subtotal \$ _____

Taxes \$ _____

(Add appropriate sales tax for Virginia, Maryland, Pennsylvania, and the District of Columbia or Canadian GST to the reprints if your order is to be shipped to these locations.)

First address included, add \$32 for
each additional shipping address \$ _____

TOTAL \$ _____

Shipping Address (cannot ship to a P.O. Box) Please Print Clearly

Name _____

Institution _____

Street _____

City _____ State _____ Zip _____

Country _____

Quantity _____ Fax _____

Phone: Day _____ Evening _____

E-mail Address _____

Additional Shipping Address* (cannot ship to a P.O. Box)

Name _____

Institution _____

Street _____

City _____ State _____ Zip _____

Country _____

Quantity _____ Fax _____

Phone: Day _____ Evening _____

E-mail Address _____

* Add \$32 for each additional shipping address

Payment and Credit Card Details

Enclosed: Personal Check _____

Credit Card Payment Details _____

Checks must be paid in U.S. dollars and drawn on a U.S. Bank.

Credit Card: ☐ VISA ☐ Am. Exp. ☐ MasterCard

Card Number _____

Expiration Date _____

Signature: _____

Please send your order form and prepayment made payable to:

Cadmus Reprints

P.O. Box 751903

Charlotte, NC 28275-1903

Note: Do not send express packages to this location, PO Box.

FEIN #:541274108

Signature _____

Date _____

Signature is required. By signing this form, the author agrees to accept the responsibility for the payment of reprints and/or all charges described in this document.

Invoice or Credit Card Information

Invoice Address Please Print Clearly

Please complete Invoice address as it appears on credit card statement

Name _____

Institution _____

Department _____

Street _____

City _____ State _____ Zip _____

Country _____

Phone _____ Fax _____

E-mail Address _____

Cadmus will process credit cards and *Cadmus Journal* Services will appear on the credit card statement.

If you don't mail your order form, you may fax it to 410-820-9765 with your credit card information.

Radiology 2008

Black and White Reprint Prices

Domestic (USA only)						
# of Pages	50	100	200	300	400	500
1-4	\$221	\$233	\$268	\$285	\$303	\$323
5-8	\$355	\$382	\$432	\$466	\$510	\$544
9-12	\$466	\$513	\$595	\$652	\$714	\$775
13-16	\$576	\$640	\$749	\$830	\$912	\$995
17-20	\$694	\$775	\$906	\$1,017	\$1,117	\$1,220
21-24	\$809	\$906	\$1,071	\$1,200	\$1,321	\$1,471
25-28	\$928	\$1,041	\$1,242	\$1,390	\$1,544	\$1,688
29-32	\$1,042	\$1,178	\$1,403	\$1,568	\$1,751	\$1,924
Covers	\$97	\$118	\$215	\$323	\$442	\$555

International (includes Canada and Mexico)						
# of Pages	50	100	200	300	400	500
1-4	\$272	\$283	\$340	\$397	\$446	\$506
5-8	\$428	\$455	\$576	\$675	\$784	\$884
9-12	\$580	\$626	\$805	\$964	\$1,115	\$1,278
13-16	\$724	\$786	\$1,023	\$1,232	\$1,445	\$1,652
17-20	\$878	\$958	\$1,246	\$1,520	\$1,774	\$2,030
21-24	\$1,022	\$1,119	\$1,474	\$1,795	\$2,108	\$2,426
25-28	\$1,176	\$1,291	\$1,700	\$2,070	\$2,450	\$2,813
29-32	\$1,316	\$1,452	\$1,936	\$2,355	\$2,784	\$3,209
Covers	\$156	\$176	\$335	\$525	\$716	\$905

Minimum order is 50 copies. For orders larger than 500 copies, please consult Cadmus Reprints at 800-407-9190.

Reprint Cover

Cover prices are listed above. The cover will include the publication title, article title, and author name in black.

Shipping

Shipping costs are included in the reprint prices. Domestic orders are shipped via UPS Ground service. Foreign orders are shipped via a proof of delivery air service.

Multiple Shipments

Orders can be shipped to more than one location. Please be aware that it will cost \$32 for each additional location.

Delivery

Your order will be shipped within 2 weeks of the journal print date. Allow extra time for delivery.

Color Reprint Prices

Domestic (USA only)						
# of Pages	50	100	200	300	400	500
1-4	\$223	\$239	\$352	\$473	\$597	\$719
5-8	\$349	\$401	\$601	\$849	\$1,099	\$1,349
9-12	\$486	\$517	\$852	\$1,232	\$1,609	\$1,992
13-16	\$615	\$651	\$1,105	\$1,609	\$2,117	\$2,624
17-20	\$759	\$787	\$1,357	\$1,997	\$2,626	\$3,260
21-24	\$897	\$924	\$1,611	\$2,376	\$3,135	\$3,905
25-28	\$1,033	\$1,071	\$1,873	\$2,757	\$3,650	\$4,536
29-32	\$1,175	\$1,208	\$2,122	\$3,138	\$4,162	\$5,180
Covers	\$97	\$118	\$215	\$323	\$442	\$555

International (includes Canada and Mexico))						
# of Pages	50	100	200	300	400	500
1-4	\$278	\$290	\$424	\$586	\$741	\$904
5-8	\$429	\$472	\$746	\$1,058	\$1,374	\$1,690
9-12	\$604	\$629	\$1,061	\$1,545	\$2,011	\$2,494
13-16	\$766	\$797	\$1,378	\$2,013	\$2,647	\$3,280
17-20	\$945	\$972	\$1,698	\$2,499	\$3,282	\$4,069
21-24	\$1,110	\$1,139	\$2,015	\$2,970	\$3,921	\$4,873
25-28	\$1,290	\$1,321	\$2,333	\$3,437	\$4,556	\$5,661
29-32	\$1,455	\$1,482	\$2,652	\$3,924	\$5,193	\$6,462
Covers	\$156	\$176	\$335	\$525	\$716	\$905

Tax Due

Residents of Virginia, Maryland, Pennsylvania, and the District of Columbia are required to add the appropriate sales tax to each reprint order. For orders shipped to Canada, please add 7% Canadian GST unless exemption is claimed.

Ordering

Reprint order forms and purchase order or prepayment is required to process your order. Please reference journal name and reprint number or manuscript number on any correspondence. You may use the reverse side of this form as a proforma invoice. Please return your order form and prepayment to:

Cadmus Reprints

P.O. Box 751903
Charlotte, NC 28275-1903

Note: Do not send express packages to this location, PO Box. FEIN #:541274108

Please direct all inquiries to:

Rose A. Baynard

800-407-9190 (toll free number)
410-819-3966 (direct number)
410-820-9765 (FAX number)
baynardr@cadmus.com (e-mail)

Reprint Order Forms and purchase order or prepayments must be received 72 hours after receipt of form.