University of Groningen

Re

Reijneveld, Sijmen A.

Published in:
Child Abuse & Neglect

DOI:
[10.1016/j.chiabu.2007.04.001](https://doi.org/10.1016/j.chiabu.2007.04.001)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2007

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Reijneveld, S. A. (2007). Re: Age-related incidence curve of hospitalized Shaken Baby Syndrome cases: Convergent evidence for crying as a trigger to shaking (Barr, Trent, & Cross, 2006). *Child Abuse & Neglect*, 31(6), 601-602. <https://doi.org/10.1016/j.chiabu.2007.04.001>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

ELSEVIER

Child Abuse & Neglect 31 (2007) 601–602

Child Abuse
& Neglect

Letter to the Editor

Re: Age-related incidence curve of hospitalized Shaken Baby Syndrome cases: Convergent evidence for crying as a trigger to shaking (Barr, Trent, & Cross, 2006)

To the Editor:

In their recent paper on infant crying as a trigger to shaking, Barr, Trent, and Cross (2006) show the age-related incidence curve of Shaken Baby Syndrome hospitalizations to be of a shape very similar to that of normal crying. The ages at which both curves peak differ somewhat, but a likely explanation for these differences is that multiple shakings may have occurred before the shaking leading to hospitalization.

I agree with Barr et al. that it is indeed very likely that the crying of the infant is the precipitating factor in at least a part of the cases of Shaken Baby Syndrome. However, the evidence is even more compelling than these authors suggest. The results of a previous study on 3,345 infants show that parents report to undertake a number of actions detrimental for infant health to stop infant from crying (Reijneveld, Brugman, Hira Sing, & Verloove-Vanhorick, 2004). For instance, 3.35% of the parents of 6-month-old infants report to have shaken their baby at least once to stop their crying. For parents of 1-month-old infants, this percentage is 1.10%. Of the parents of 6-month-old infants, 5.60% report to have shaken their infant to stop their crying or to have undertaken similar detrimental actions like smothering or slapping. Probably, these percentages are actually higher because of parental underreporting, which is likely to occur even though it concerns an anonymous and confidential questionnaire (Reijneveld et al., 2004). And most important: the worries of parents about the crying of their infant and their judging this crying excessive are much more predictive than the actual amount of crying, measured by Wessel's criteria for its duration (Reijneveld, Van der Wal, Brugman, Hira Sing, & Verloove-Vanhorick, 2004; Wessel, Cobb, Jackson, Harris, & Detwiller, 1954).

This evidence strongly supports the association between infant crying and the Shaken Baby Syndrome as suggested by Barr et al. (2006). Moreover, it provides important items for its prevention. This prevention should especially focus on effect of the crying on the parents, and on their skills to better cope with both the crying and the stress that it provokes. Just diminishing the crying is probably less adequate because the judgment of the crying appears to be more important than its actual duration (Reijneveld et al., 2004). These interventions should be especially targeted at parents that report their infant to cry excessively. Improving the coping skills of these parents may be beneficial both for the prevention of the Shaken Baby Syndrome and for the prevention of abuse in later child life.

References

- Barr, R. G., Trent, R. B., & Cross, J. (2006). Age-related incidence curve of hospitalized Shaken Baby Syndrome cases: Convergent evidence for crying as a trigger to shaking. *Child Abuse & Neglect*, 30, 7–16.
- Reijneveld, S. A., Brugman, E., Hira Sing, R. A., & Verloove-Vanhorick, S. P. (2004). Parental reports of excessive infant crying: The impact of varying definitions. *Pediatrics*, 108, 893–897.
- Reijneveld, S. A., Van der Wal, M. F., Brugman, E., Hira Sing, R. A., & Verloove-Vanhorick, S. P. (2004). Infant crying and abuse. *Lancet*, 364, 1440–1442.
- Wessel, M. A., Cobb, J. C., Jackson, E. B., Harris, G. S., & Detwiller, A. C. (1954). Paroxysmal fussing in infancy, sometimes called “colic”. *Pediatrics*, 14, 421–434.

Sijmen A. Reijneveld^{a,b,*}

^a Department of Health Sciences
University Medical Center Groningen
University of Groningen
Groningen, The Netherlands

^b TNO Quality of Life
Department of Child Health
Leiden, The Netherlands

* Corresponding author address: University Medical Center
Groningen, University of Groningen, Department of Health Sciences
PO Box 196, 9700 AD Groningen, The Netherlands.

Available online 29 May 2007