University of Groningen

Are the outcomes of clinical pathways evidence-based? A critical appraisal of clinical pathway evaluation research

El Baz, N.; Middel, B.; Van Dijk, J.P.; Oosterhof, A.; Boonstra, P.W.; Reijneveld, S.A.

Published in:
Journal of Evaluation in Clinical Practice

DOI:
[10.1111/j.1365-2753.2006.00774.x](https://doi.org/10.1111/j.1365-2753.2006.00774.x)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2007

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

El Baz, N., Middel, B., Van Dijk, J. P., Oosterhof, A., Boonstra, P. W., & Reijneveld, S. A. (2007). Are the outcomes of clinical pathways evidence-based? A critical appraisal of clinical pathway evaluation research. *Journal of Evaluation in Clinical Practice*, 13(6), 920-929. <https://doi.org/10.1111/j.1365-2753.2006.00774.x>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Are the outcomes of clinical pathways evidence-based? A critical appraisal of clinical pathway evaluation research

Noha El Baz MSc,¹ Berrie Middel PhD,² Jitse P. van Dijk MD PhD,³ Andre Oosterhof RN MBA,⁴ Piet W. Boonstra MD PhD⁵ and Sijmen A. Reijneveld MD PhD⁶

¹PhD Student, ²Senior Lecturer Research Methodology, Department of Health Sciences, Subdivision Care Sciences, University Medical Center Groningen, University of Groningen, The Netherlands

³Associate Professor in Public Health, ⁶Professor of Public Health, Head of the Department of Health Sciences, Department of Health Sciences, Subdivision Public Health, University Medical Center Groningen, University of Groningen, The Netherlands

⁴Head Nurse Thorax Center, ⁵Professor of Thoracic Surgery, Head of the Thorax Center, Thorax Center, University Medical Center Groningen, University of Groningen, The Netherlands

Keywords

clinical pathway, complication, length of stay, methodological quality, readmission, systematic review

Correspondence

Berrie Middel
Department of Health Sciences
University Medical Center Groningen
University of Groningen
PO Box 196
9700 AD Groningen
The Netherlands
E-mail: b.middel@med.umcg.nl

Accepted for publication: 19 June 2006

doi:10.1111/j.1365-2753.2006.00774.x

Abstract

Aim and objective To evaluate the validity of study outcomes of published papers that report the effects of clinical pathways (CP).

Method Systematic review based on two search strategies, including searching Medline, CINAHL, Embase, Psychinfo and Picarta from 1995 till 2005 and ISI Web of KnowledgeSM. We included randomized controlled or quasi-experimental studies evaluating the efficacy of clinical pathway application. Assessment of the methodological quality of the studies included randomization, power analysis, selection bias, validity of outcome indicators, appropriateness of statistical tests, direct (matching) and indirect (statistical) control for confounders. Outcomes included length of stay, costs, readmission rate and complications. Two reviewers independently assessed the methodological quality of the selected papers and recorded the findings with an evaluation tool developed from a set of items for quality assessment derived from the Cochrane Library and other publications.

Results The study sample comprised of 115 publications. A total of 91.3% of the studies comprised of retrospective studies and 8.7% were randomized controlled studies. Using a quality-scoring assessment tool, 33% of the papers were classified as of good quality, whereas 67% were classified as of low quality. Of the studies, 10.4% controlled for confounding by matching and 59.1% adopted parametric statistical tests without testing variables on normal distribution. Differences in outcomes were not always statistically tested.

Conclusion Readers should be cautious when interpreting the results of clinical pathway evaluation studies because of the confounding factors and sources of contamination affecting the evidence-based validity of the outcomes.

Introduction

Clinical pathways have been developed in health care as multidisciplinary care plans that outline the sequence and timing of actions necessary for achieving expected patient outcomes and organizational goals regarding quality, costs, patient satisfaction and efficiency. The concept of *clinical pathways* (CP) refers to specific guidelines for care that describe patient treatment goals and define a sequence and timing of intervention for meeting those goals efficiently [1]. They can also be defined as care plans that detail essential steps in patient care with a view to describing the expected progress of the patient [2]. They are also known as critical pathways, integrated care pathway, critical path, care mapsTM and care paths and they are being embraced in many systems.

In an attempt to evaluate the efficacy of integrated care pathways, Campbell *et al.* posed the question 'Are clinical pathways effective in improving patient care?' [3]. They used the results of a comprehensive review performed by the National Health Service in Wales in 1996, which was comprised of approximately 4000 references to integrated care pathways and related topics worldwide. The studies that were found mainly described benefits that were experienced and addressed concerns associated with the use of pathways or practical barriers to implementation. Most of the studies they found were uncontrolled 'before-after' studies and no randomized controlled studies were found. The authors came to the conclusion that 'these reports do not provide reliable evidence and publication bias is highly likely, favoring publications reporting favorable experience'.

Every *et al.* reported that in *cardiovascular medicine*, although the studies they evaluated were somewhat under-powered, the overall experience had been promising [1]. Clinical pathways applied to patients with a cardiovascular disease showed a tendency towards a decreased treatment variation, improved guideline compliance and reduced costs. However, the evidence of the effectiveness of clinical pathways in cardiovascular medicine cannot be generalized because of the insufficient number of controlled studies. Renholm *et al.* concluded in a review article that clinical pathways had positive effects on patient-care outcome, although some studies did suggest that the use of clinical pathways had no influence on patient-care outcomes, while by the same token they also stated that there was no evidence at all that they had any negative effect [2].

Similarly, Van Herck *et al.* concluded that clinical pathways did have a positive effect on patient outcome, but they did not take methodological weaknesses into consideration because they analysed most of the manuscripts (55.5%) by means of abstracts [4]. Additionally, they expressed their concerns about 'publication bias since clinical pathways with no, few, or even negative results hardly ever get published'.

Kim *et al.* conducted a systematic review which focused on the effectiveness of clinical pathways for total knee and total hip arthroplasty [5]. They included 11 papers and identified only one randomized controlled study. They reported a decrease in length of stay (LOS) and in costs with either reduced or unchanged rates of complications and either improvement or no change in patient-reported outcomes. Furthermore, they concluded that, although the data in their review supported the effectiveness of clinical pathways, 'definitive conclusions cannot be made because of methodological limitations'.

Another systematic review was conducted by Kwan *et al.* regarding clinical pathways for stroke patients [6,7]. They included both randomized and non-randomized studies and found no evidence that clinical pathways provided any significant additional benefit over standard medical care in terms of major clinical outcomes (death or discharge destination). Moreover, they concluded that stroke patients in clinical pathway groups were more dependent on discharge, while the effect on LOS and hospitalization costs remained unclear.

Despite these uncertainties, clinical pathways have been widely used in many institutions throughout the United States, the United Kingdom and some parts of Europe for patients undergoing various treatments, surgeries or diagnostic procedures. According to the results of the majority of publications in which clinical pathways were compared with standard care, there is an overall tendency to come to the conclusion that clinical pathways are effective in reducing LOS, costs, complication rates and readmissions. On the other hand, clinical pathways may positively affect patients' health-related functioning, quality of life or patient satisfaction.

However, some critical studies [1,4,5,8] have introduced serious doubts about the amount of the evidence, and this criticism stimulated us to undertake this study. The purpose of this study was to appraise the methodological qualities of clinical pathway evaluation studies in order to evaluate the validity of the evidence for the efficacy of clinical pathways and to suggest improvements in future study design. Methodological quality was defined through a set of parameters related to the design and conduct of

the study that determines the internal and external validity of the study [9–12].

Methods

Two *search strategies* were employed to find published studies on the efficacy of clinical pathways. The first strategy involved computerized database searches using Medline, CINAHL, Embase, Psycinfo and Picarta from 1995 till 2005. Studies were identified by a broad range of keywords: clinical pathways, critical pathway, clinical paths and integrated care pathway, effect, adults, paediatric diseases, patient satisfaction, length of stay, complications, readmission, quality of care, quality of life, costs, longitudinal studies, experimental studies, randomized controlled studies, cohort studies, double-blind methods, systematic review, evaluation and comparative studies.

The second strategy concerned a snowball sampling method using the databases of ISI Web of KnowledgeSM. We identified studies that were cited retrospectively in a publication on the effectiveness of clinical pathways, as well as studies that cited the selected papers in the years following their publication.

Inclusion and exclusion of publications

Information from abstracts and titles of the papers that were detected using both search strategies was used to include or exclude manuscripts. Papers were excluded if they were:

- manuscripts that addressed the noun 'pathway' but were not related to studies investigating the efficacy of a clinical pathway in terms of specified guidelines or outlines for care that describe patient-treatment goals and define a sequence and timing of interventions to meet those goals efficiently;
- manuscripts concerning our definition of a clinical pathway which did not provide empirical quantitative results, such as letters to the editor, brief reports, case studies, qualitative designs, opinions of experts, etc.

After exclusion of inappropriate manuscripts, two of the investigators (NEB and BM) independently assessed the remaining publications in full text against two criteria:

- 1 The study evaluated the efficacy of a clinical pathway by means of quantitative methods (e.g. a meta-analysis or systematic review).
- 2 The design of the effect study included a clinical pathway group and a control group.

Papers were included if one or both investigators unequivocally considered the publication as appropriate for analysis. Differences were resolved through discussion with reference to a third reviewer (JPvD) if necessary. Methodological appraisal included papers that satisfied both criteria.

Assessment of methodological quality

Two reviewers (NEB and BM) independently assessed the methodological quality of the selected papers and recorded their findings with an evaluation tool or scoring system comprising of a set of items for quality assessment from the Cochrane Library and from other publications on quality assessment of studies [9–11,13]. The quality score was based on eight items that evaluated

Table 1 The evaluation items used to assess the methodological quality of the studies investigated

Was a power analysis performed?	Yes/No/Insufficient information
Was a method of randomization performed?	Yes/No/Insufficient information
Did a medical ethics committee assess the study protocol?	Yes/No/Insufficient information
Were the groups similar at baseline in regards to the most important prognostic indicators?	Yes/No/Insufficient information
Were the eligibility criteria specified adequately?	Yes/No/Insufficient information
Were the operational definitions of outcomes valid?	Yes/No/Insufficient information
Were the statistical tests appropriate to the data?	Yes/No/Insufficient information
Were potential confounders controlled in the study design (e.g. matching) or statistical analysis (multivariate analysis)?	Yes/No/Insufficient information
Were dropouts or losses in follow-up analysed appropriately?*	Yes/No/Insufficient information

*Because of the fact that only 10 studies reported the occurrence of patient dropout or losses in follow-up, this parameter was not incorporated into the final score.

the following methodological characteristics of individual studies as presented in Table 1.

The appraisal of the overall methodological quality was based on a weighted score in which the similarity of groups at baseline, randomization and adequate statistical methods were assigned three points; the use of power analysis and control of confounders were assigned two points; and assessment by a medical ethics committee, unbiased outcome measure and eligibility criteria were assigned one point. We calculated overall quality scores for each study by summing up the score-weighted 'yes' scores.

We established the cut-off point between 'high' and 'low' quality studies by following the method developed by Verhagen *et al.* [12] They set the cut-off point at 50% of the maximum achievable score of 16 points, meaning high-quality studies scored ≥ 8.0 and low-quality studies scored ≤ 7.0 points. In addition to this dichotomous scale, we arranged the quality scale scores into the following ordinal categories: invalid studies (scores 0–3), weak to medium quality (scores 4–7), good quality (scores 8–11) and high quality (scores 12–16).

Analysis

Analyses were performed using SPSS version 12.0.1 (SPSS Inc., Chicago, IL, USA) and for all tests $P < 0.05$ was considered significant. We calculated 95% confidence intervals for the differences in proportions [14]. Chi-square and Fisher's exact tests were used for associations between categorical variables. Apart from the methodological parameters in Table 1, outcomes such as decrease in LOS, costs, readmission and complications and their statistical significance were observed.

Results

Of the 556 publications that were analysed, 310 papers (55.7%) were not related to our definition of a clinical pathway at all as they addressed either metabolic, molecular or genetic pathways, letters to the editor and editorials or investigated the effect of a pharmacological therapy, or aspects of surgical techniques. Of the remaining sample of 246 publications, 131 manuscripts were excluded because they either did not meet the inclusion criterion regarding required study design (74 papers contained no evaluation study, 52 papers addressed the definition or phases of development of clinical pathways and five were not in English). A total of 115 studies out of 556 investigated the efficacy of clinical path-

Figure 1 Flowchart describing study selection and excluded studies.

ways according to our definition of clinical pathways and these were used for the assessment of methodological quality (see Fig. 1).

Disease treatments

In the period between 1995 and 2005, 246 publications addressed relevant aspects of clinical pathways in terms of our definition. In Table 2, these publications are shown sorted by disease or intervention category. The most often studied category was in the field of cardiovascular surgery and diseases in both the gross sample and study sample (21.5% and 17.4%, respectively). Twenty publications evaluating cardiovascular clinical pathways were included in the study sample [15–34]. The relative number (%) of publications included in the study sample varied between 5% and 16% in the following domains: (1) respiratory diseases, therapy and thoracic surgery [28,35–51]; (2) gastrointestinal surgery, endoscopic

Table 2 Studies on clinical pathways across diseases and interventions published between 1995 and 2005

Disease and interventions	Gross sample of publications <i>n</i> (%)	Study sample <i>n</i> (%)	95% CI
1. Cardiovascular surgery and diseases	53 (21.5)	20 (17.4)	-12.7-4.5
2. Respiratory diseases, therapy and thoracic surgery	20 (8.1)	18 (15.6)	-14.0-0.3
3. Gastrointestinal surgery, endoscopic surgery and diseases (peritonitis)	24 (9.8)	19 (16.5)	-14.5-0.1
4. Orthopaedic surgery and multiple trauma	22 (8.9)	13 (11.3)	-9.2-4.4
5. Oncological diseases and surgery	14 (5.7)	9 (7.8)	-7.8-3.5
6. Vascular surgery	8 (3.2)	7 (6.1)	-7.7-2.0
7. Neurological trauma, disorders, diseases and pain management	18 (7.3)	8 (6.9)	-5.3-6.1
8. Urological diseases, surgery and catheterization	15 (6.1)	5 (4.3)	-3.1-6.5
9. Gynaecological diseases, surgery and maternity care	6 (2.4)	6 (5.2)	-7.3-1.7
10. Paediatric conditions (emergencies, neonatal care, oral rehydration and tonsillectomy)	4 (1.6)	3 (2.6)	-2.3-4.3
11. Head and neck surgery	3 (1.2)	2 (1.8)	-2.2-3.3
12. Burns, decubitus ulcer reconstructive surgery	3 (1.2)	2 (1.8)	-2.2-3.3
13. Psychological disturbances and mental health	10 (4.1)	2 (1.8)	-5.7-1.1
14. Metabolic diseases	5 (2.0)	1 (0.9)	-3.6-1.3
15. Critical pathway evaluation and job satisfaction	6 (2.4)	0 (0)	-4.3-0.5
16. Bone marrow transplant	1 (0.4)	0 (0)	-1.2-0.4
17. Not in English language	7 (2.8)	0 (0)	-4.9-0.8
18. Not related to a specific disease or medical treatment	27 (11.0)	0 (0)	
Total	246 (100)	115 (100)	

surgery and diseases [52-70]; (3) orthopedic surgery and multiple trauma [71-83]; (4) oncological diseases and surgery [84-92]; (5) neurological trauma, disorders, diseases and pain management [93-100]; (6) vascular surgery [15,19,101-105]; and (7) gynaecological diseases, surgery and maternity care [106-111].

Categories of diseases or treatment, which represent less than 5% of the study sample, comprised studies on urological diseases, surgery and procedure [112-116]; psychological disturbances and mental health [117,118]; metabolic diseases [119]; paediatric conditions [120-122]; burn and skin reconstructive surgery [123,124]; and head and neck surgery [125,126].

We tested whether selection bias affected the sample of manuscripts used for methodological analysis with 95% confidence intervals for differences in proportions. The differences between the proportion of papers addressing the effects of clinical pathways on job satisfaction and papers not in English showed underrepresentation in the final sample (used in the current study) as compared with the total sample of clinical pathway-related publications.

The study sample of papers shows a similar distribution across diseases as compared with the gross sample, indicating a good representation of the population of studies published between 1995 and 2005.

Designs

Of the 246 papers, 131 were excluded because of the fact that they were descriptive studies or review articles and only the remaining 115 publications were included based on the criteria of study design. Ten randomized controlled studies were found and the majority ($n = 105$) were comprised of studies with a retrospective comparative research design or were cross-sectional retrospective studies that compared the differences in patient outcome during a

period before and after implementation of clinical pathways. The following retrospective designs were used:

1 A majority of these studies ($n = 96$) used a historical control group and were conducted at the same hospital (before-after design).

2 Three studies were conducted with a historical control group from a different hospital and at different time periods.

3 Six studies were conducted using concurrent control and experimental groups either in the same hospital, though using separate wards, or in different hospitals.

Ten studies, labelled as randomized controlled studies, comprised studies that randomly selected hospitals [45,127] where a pathway was implemented or that assigned patients randomly to either pathway care or standard care.

These randomized controlled studies followed up patients after discharge for outcomes like quality of life, pain, readmission, mortality and complications. One paper assessed patients at baseline, 3, 6 and 12 months [117], and another measured patients 1 week after assignment to CP or conventional care and then at 4, 12 and up to 26 weeks [96-98]. The remaining papers followed up patients after discharge from the health care facility at a time ranging from 10 days up to 12 weeks [42,45,56,72,127,128].

Randomization and matching

We detected 12 retrospective studies (10.4%) that controlled for confounding through matching, of which three studies used a random sample from a clinical pathway group which was matched with controls from the pre-pathway period [25,30,48,50,58,79,81,107,110,111,122,129]. Furthermore, 10 randomized controlled studies were found, of which two studies randomly assigned hospitals either to implement a clinical pathway or to remain on standard care [45,127]. Eight studies randomly

assigned patients to either a clinical pathway or standard care [42,56,72,96–98,117,128].

Power analysis

The question of what sample size to use constitutes a crucial part of any research proposal. However, only 16.5% ($n = 19$) of the studies conducted a power analysis to determine in advance the required number of observations which would be sufficient to provide the required precision of results. Among the 115 studies, 25% of the samples were very small ($n < 50$), 25% ranged from 51 to 100 patients, 25% ranged from 100 to 200 patients and 25% had samples greater than 200 patients in either the clinical pathway or the control group. No statistically significant association was found between sample size ($n < 100$ vs. $n > 100$) and performance for a statistical power analysis (Chi-square, $P = 0.56$).

Control for confounding

In less than half of the studies (42.1%), the authors did pay attention to the problem of controlling for potential confounders and either applied direct control with randomization or matching (17.5%), and/or used a control for confounders (e.g. co-morbidity, age and gender) in regression models (24.6%). The other (57.9%) did not take the potential risk of confounders into account at all.

Selection bias may occur when inclusion and exclusion criteria are obscure. In 57 of the studies that were appraised (49.6%), misleading conclusions were prone to arise based on the fact that patients in 'experimental' groups differed from the control group patients regarding characteristics such as differences in age, gender, disease severity or co-morbidity. The other half of the studies meticulously described criteria for inclusion and exclusion.

Accuracy and validity of outcome measures

Length of stay was evaluated in 108 publications (93.3%). However, more than a quarter (28.1%) of these studies gave no accurate or meticulous description of its operationalization or a clear description of the way it was assessed. Costs and hospital charges were assessed in 73 papers (63.5%), among which 62 (53.9%) stated a clear description of the charges and costs calculated.

- *Readmission rates* were calculated in 53 papers (46.1%), of which 50 (43.5%) precisely defined readmission within a time frame.
- *Complications* were evaluated in 70 papers (60.9%), of which 64 (55.6%) defined and clearly stated the complications.
- *Quality of life* was assessed in 10 (8.7%) of the studies with a validated measure. Functional health-related functioning was measured in six (5.2%) studies, and both quality of life and health status were measured in two papers. Three studies (2.6%) assessed psychological distress (anxiety and depression). However, only two studies used a validated measure: the Hospital and Anxiety and Depression Scale [48,98].
- *Patient satisfaction* was assessed in 15 studies (13.0%), but was measured with a multi-item tool in 13 studies (11.3%). *Work satisfaction* was evaluated in four (3.5%) of the studies, and three studies presented an accurate description of this construct. Clinical quality-of-care indicators were evaluated in 57 (49.6%) papers and were accurately defined in all cases.

Appropriateness of statistical methods

More than half (59.1%) of the studies adopted parametric statistical tests without question, but the rest (40.9%) tested variables over normal distribution and, depending on the outcome, used non-parametric tests. Reduction of LOS, costs, readmission rates and number of complications belong to the most relevant targets for implementing clinical pathways.

However, decreases in LOS, costs, readmission rates and number of complications were not statistically tested in 12.3%, 28.8%, 20.8% and 27.1% of the studies, respectively [16–19, 22,23,25–28,32,33,36–38,42,44–47,49,53,61,75,85,88,91,92,99–103,105,110,113,120]. There were studies that used a statistical test to decide whether a difference between a clinical pathway group and a control group was due to sample fluctuation, but they also reported other differences without this test. For the main outcome parameters, LOS, readmission rates, costs and number of complications (65%) of the studies tested for each outcome (23%) did not test for all differences between clinical pathways and controls, while (12%) did not apply any test at all.

Quality of studies related to statistically significant outcomes

We found 92 publications that reported a decrease in LOS and 60 that reported a decrease in costs. All the good-quality studies reported a statistically significant result in both LOS and costs. However, among the low-quality studies (84%) of the papers reported a reduction in LOS that was statistically significant, and only 68% of the publications reported a decrease in costs which was statistically significant (Fisher's exact test, $P = 0.02$ and $P = 0.03$, respectively). There was no association between quality of the studies and the statistical significance of the reduction in complications and in readmission rates (see Table 3).

Overall quality related to other study characteristics

According to the dichotomous threshold of Verhagen *et al.* [12], one-third of the papers (33%) analysed were classified as high-quality papers, while 67% were classified as studies of low quality. However, the ordinal quality scale showed that 35.7% of the studies reflected low quality, that is, invalid studies (31.1%) were appraised as weak- to medium-quality studies, and 24.3% as good-quality studies. Finally, 8.7% of the sample reflected very good quality.

No statistically significant associations were found between the quality of the studies and the sample size (Mann–Whitney U -test/ Wilcoxon W -test, $Z = -0.48$, $P = 0.63$). Across the diseases shown in Table 1, the differences between proportions of low- or high-quality studies were not statistically significant. Furthermore, the quality of the studies was not associated with treatment in terms of surgery versus non-surgery (Fisher's exact test, $P = 0.31$).

Data extracted from hospital records were used in 81% of the studies and 19% used self-reported questionnaires or interviews in combination with data from the hospital records.

Both dichotomous and ordinal categorization confirmed that studies which qualified as 'good quality' were more likely to use patient record information in combination with self-report

Table 3 Comparison of statistically significant results on the two main outcome measures between studies with low and good methodological quality

Statistically significant reduction in:	Low quality	Good quality	Total	<i>P</i> *
Length of stay				
Yes	52 (83.9%)	30 (100%)	82	0.02
No	10 (16.1%)	0	10	
Total	62	30	92	
Costs				
Yes	32 (68.1%)	13 (100%)	45	0.03
no	15 (31.9%)	0	15	
Total	47	13	60	

*Fisher's exact test.

questionnaires or interviews (Fisher's exact test, $P = 0.01$). Because of the fact that the majority of studies (91.3%) used electronic databases, the question of whether dropouts were analysed appropriately did not play a significant role in our analysis. A significant association was found between the quality of the study and its design. Only 27% of the studies with a retrospective design were classified as of good quality (Fisher's exact test, $P = 0.0001$), whereas all randomized controlled studies were of high quality.

Discussion

The majority of the publications on CP we analysed were classified as studies of low quality (67%), which raise questions about the validity of the evidence for the implementation of clinical pathway in the last decade.

The methodology of studies assessing the efficacy of clinical pathways has been criticized in regard to their research designs, poor reporting of the methodology and under-powered sample sizes [1,3,5–8]. Therefore, we performed an appraisal and analysis of the methodological qualities of a large number of clinical pathway evaluation studies in order to investigate the validity of their outcomes.

In relation to the internal and external validity of the studies in the sample, our analysis revealed several factors that influenced the quality of clinical pathway evaluation. Most studies in our sample concerned non-randomly selected small patient populations without a power analysis [19,23,25,36,52,54,56,58,59,61,76,81,86–88,91,99,107,111,118,122,123,126,128].

Furthermore, in many studies selection bias has occurred. Patients selected for a clinical pathway were likely to differ from patients managed with standard care using characteristics known to be associated with LOS and hospital delay, and consequently included costs such as age and co-morbidity. In half of the study sample (49.6%), the inclusion and exclusion criteria of patients were obscure and difficult to relate to the population's characteristics. According to our findings, only 12 retrospective studies (10.4%) controlled for selection bias by matching, out of which three studies matched a random sample from a clinical pathway group with controls from the pre-pathway period group [25,30,48,50,58,79,81,107,110,111,122,129].

In regard to the outcomes measured, our analysis also revealed that most studies focused on cost issues and reductions in LOS. However, clinically relevant outcomes such as mortality, discharge disposition, quality of care as seen through the eyes of the patient, psychological distress (anxiety), care dependency and use of health services after discharge were largely ignored. A few studies focused on outcome in terms of: (1) destination of discharge [21,24,69,72,78,98,101,104,122]; (2) delay [44,109]; (3) patient education [32,124]; (4) quality of care [22,74]; and (5) psychological distress [48,98].

In relation to the use of appropriate statistical tests, more than half (59.1%) of the studies adopted parametric statistical tests without question, but 40.9% tested variables on normal distribution and, depending on the outcome, used non-parametric tests. Before–after differences in frequently measured outcomes such as LOS, costs, readmission and complications were statistically tested in few of the studies. Other studies used statistical tests to decide whether differences between groups were related to sample fluctuations, while also reporting other differences without statistical tests. After reviewing all papers included, it was clear that there was an inevitable risk of misrepresentation of the true costs. Reduced LOS in the hospital may lead to admission to another health care facility, for example, a rehabilitation unit or extended care facility. Although costs related to these services are not considered as a part of hospital charges, they still add to direct health care costs and may also lead to out-of-pocket expenses for the patient. Furthermore, studies that report reduction in both LOS and costs do not take into consideration that both are interrelated, which leads to contamination in statistical analysis.

Moreover, from a methodological point of view both the investigators (who assessed the outcomes) and the health care providers should have been blinded to the use of a clinical pathway since this might have biased their observations and assignment procedure.

In the light of these findings, it can be concluded that readers must be extremely cautious when interpreting the results of clinical pathway evaluation studies because of the confounding factors and the sources of contamination affecting the internal and external validity of most of the published studies.

Recommendations for future research

After reviewing a large number of clinical pathway evaluation studies and having stated our conclusions, we recommend that:

1 More (randomized) controlled studies should be conducted, in which patients are randomly assigned to the condition of either a pathway or standard should be conducted. However, such randomized controlled studies in the same hospital invite contamination because many of the same doctors, as well as care staff, are involved in treating the same population of patients. To avoid such Hawthorne effects, we suggest establishing multi-centre trials with randomization after pre-stratification of confounding factors (e.g. gender, co-morbidity) with a clearly defined method of randomization, concealment of allocation or blinding with the appropriate balancing method [130].

2 Standardization of the total direct costs is specified by clearly defined cost components and a standardized operational definition of LOS. LOS should not be confined to the hospital setting, but should be extended to include whether patients are discharged home or to an extended health care facility, and should include

whether this is a permanent or temporary arrangement. An accurate calculation in this case will reflect the true effect of clinical pathways on LOS and subsequent related costs.

3 More attention should be paid to measuring relevant patient outcomes such as quality of life, hospital anxiety, patient expectations and satisfaction with standardized validated tools, which reflect the true effect with use of appropriate statistical methods.

Until the results of good-quality research are more widely available, CPs should be considered as a potentially effective (but not evidence-based) practice to improve patient care.

References

- Every, N. R., Hochman, J., Becker, R., Kopecky, S. & Cannon, C. P. (2000) AHA scientific statement. Critical pathways: a review. *Circulation*, 101, 461–465.
- Renholm, M., Leino-Kilpi, H. & Suominen, T. (2002) Critical pathways: a systematic review. *Journal of Nursing Administration*, 32, 196–202.
- Campbell, H., Hotchkiss, R., Bradshaw, N. & Porteous, M. (1998) Integrated care pathways. *British Medical Journal*, 316, 133–137.
- Van Herck, P., Vanhaecht, K. & Sermeus, W. (2004) Effects of clinical pathways: do they work? *Journal of Integrated Care Pathways*, 8, 95–105.
- Kim, S., Losina, E., Solomon, D. H., Wright, J. & Katz, J. N. (2003) Effectiveness of clinical pathways for total knee and total hip arthroplasty – literature review. *Journal of Arthroplasty*, 18, 69–74.
- Kwan, J. & Sandercock, P. (2004) In-hospital care pathways for stroke. *Cochrane Database of Systematic Reviews*, 18, CD002924.
- Kwan, J. & Sandercock, P. (2005) In-hospital care pathways for stroke. *Stroke*, 36, 1348–1349.
- Kwan, J. & Sandercock, P. (2003) In-hospital care pathways for stroke: a Cochrane systematic review. *Stroke*, 34, 587–588.
- Verhagen, A. P., de Vet, H. C., de Bie, R. A., Bouter, L. M. & Knipschild, P. G. (1998) The Delphi list: a criteria list for quality assessment of randomized clinical trials for conducting systematic reviews developed by Delphi consensus. *Journal of Clinical Epidemiology*, 51, 1235–1241.
- Higgins, J. P. T. & Green, S. (2005) Cochrane Handbook for Systematic Reviews of Interventions 4.2.5 [updated May 2005]. Chichester: John Wiley & Sons.
- Hadorn, D. C., Baker, D., Hodges, J. S. & Hicks, N. (1996) Rating the quality of evidence for clinical practice guidelines. *Journal of Clinical Epidemiology*, 49, 749–754.
- Verhagen, A. P., de Bie R. A., Lenssen A. F., de Vet H. C., Kessels A. G. H., Boers M. & Van den Brant P. A. (2000) Impact of quality items on study outcome. Treatments in acute lateral ankle sprains. *International Journal of Technology Assessment in Health Care*, 16, 1136–1146.
- Dawes, M., Davies, P., Gray, A., Mant, J., Seers, K. & Snowball, R. (1999) Evidence-Based Practice. A Primer for Health Care Professionals. Edinburgh: Churchill Livingstone.
- Newcombe, R. G. & Altman, D. G. (2005) Proportions and their differences. In *Statistics with Confidence* (eds D. G. Altman, D. Machin, T. N. Bryant & M. J. Gardner), pp. 45–56. Bristol: British Medical Journal.
- Back, M. R., Harward, T. R., Huber, T. S., Carlton, L. M., Flynn, T. C. & Seeger, J. M. (1997) Improving the cost-effectiveness of carotid endarterectomy. *Journal of Vascular Surgery*, 26, 456–462.
- Barnes, R. V., Lawton, L. & Briggs, D. (1994) Clinical benchmarking improves clinical paths – experience with coronary-artery bypass-grafting. *Joint Commission Journal on Quality Improvement*, 20, 267–276.
- Cohn, L. H., Rosborough, D. & Fernandez, J. (1997) Reducing costs and length of stay and improving efficiency and quality of care in cardiac surgery. *Annals of Thoracic Surgery*, 64, S58–S60.
- Halm, M. A. (1997) Collaborative care: improving patient outcomes in cardiovascular surgery. *Progress in Cardiovascular Nursing*, 12, 15–23.
- Jano, S. & Harlin, S. A. (2000) Designing a carotid endarterectomy critical pathway for your organization. *Military Medicine*, 165, 385–389.
- Kim, M. H., Deeb, G. M., Morady, F., *et al.* (2001) Effect of postoperative atrial fibrillation on length of stay after cardiac surgery. The Postoperative Atrial Fibrillation in Cardiac Surgery study [PACS(2)]. *American Journal of Cardiology*, 87, 881–885.
- Lazar, H. L., Fitzgerald, C. A., Ahmad, T., Bao, Y. S., Colton, T., Shapira, O. M. & Shemin, R. T. (2001) Early discharge after coronary artery bypass graft surgery: are patients really going home earlier? *Journal of Thoracic and Cardiovascular Surgery*, 121, 950.
- Nikas, D. J., Freeman, J. E., Luterma, A. R., Warnken, L. C., Nensiel, R. D., Farrell, M. & Alpert, M. A. (1996) Use of a national data base to assess perioperative risk, morbidity, mortality, and cost savings in coronary artery bypass grafting. *Southern Medical Journal*, 89, 1074–1077.
- Porter, H. B. (1998) Health resources utilization and quality of life outcomes of low risk coronary artery bypass graft patients: a comparison study. *Canadian Journal of Cardiovascular Nursing*, 9, 10–15.
- Rumble, S. J., Jernigan, M. H. & Rudisill, P. T. (1996) Determining the effectiveness of critical pathways for coronary artery bypass graft patients: retrospective comparison of readmission rates. *Journal of Nursing Care Quality*, 11, 34–40.
- Serna, D. L., Chen, J. C. & Milliken, J. C. (1998) Accelerated recovery after coronary artery bypass surgery in patients with poor left ventricular function: preliminary report. *American Surgeon*, 64, 942–946.
- Velasco, F. T., Ko, W., Rosengart, T., Altorki, N., Lang, S., Gold, J. P., Krieger, K. H. & Isom, O. W. (1996) Cost containment in cardiac surgery: results with a critical pathway for coronary bypass surgery at the New York hospital – Cornell Medical Center. *Best Practices and Benchmarking in Healthcare*, 1, 21–28.
- Walters, J., Schwartz, C., Monaghan, H., Watts, J., Schlafer, G., Deeb, G. & Bolling, S. F. (1998) Long-term outcome following case management after coronary artery bypass surgery. *Journal of Cardiac Surgery*, 13, 123–128.
- Wright, C. D., Wain, J. C., Grillo, H. C., Moncure, A. C., Macaluso, S. M. & Mathisen, D. J. (1997) Pulmonary lobectomy patient care pathway: a model to control cost and maintain quality. *Annals of Thoracic Surgery*, 64, 299–302.
- Cardozo, L. & Aherns, S. (1999) Assessing the efficacy of a clinical pathway in the management of older patients hospitalized with congestive heart failure. *Journal for Healthcare Quality*, 21, 12–16.
- Cheah, J. (2000) Clinical pathways – an evaluation of its impact on the quality of care in an acute care general hospital in Singapore. *Singapore Medical Journal*, 41, 335–346.
- Holmboe, E. S., Meehan, T. P., Radford, M. J., Wang, Y., Marciniak, T. & Krumholz, H. (1999) Use of critical pathways to improve the care of patients with acute myocardial infarction. *American Journal of Medicine*, 107, 324–331.
- Hoskins, L. M., Clark, H. M., Schroeder, M. A., Walton-Moss, B. & Thiel, L. (2001) A clinical pathway for congestive heart failure . . . ‘clinical pathways versus a usual plan of care: what’s the difference’ in last month’s issue of HHN, part 2. *Home Healthcare Nurse*, 19, 207–217.
- Nichol, G., Walls, R., Goldman, L., *et al.* (1997) A critical pathway for management of patients with acute chest pain who are at low risk

- for myocardial ischemia: recommendations and potential impact. *Annals of Internal Medicine*, 127, 996–1005.
34. Ranjan, A., Tarigopula, L., Srivastava, R. K., Obasanjo, O. O. & Obah, E. (2003) Effectiveness of the clinical pathway in the management of congestive heart failure. *Southern Medical Journal*, 96, 661–663.
 35. Abisheganaden, J., Chee, C. B., Goh, S. K., Yeo, L. S., Prabhakaran, L., Earnest, A. & Wang, Y. T. (2001) Impact of an asthma carepath on the management of acute asthma exacerbations. *Annals Academy of Medicine Singapore*, 30, 22–26.
 36. Bailey, R., Weingarten, S., Lewis, M. & Mohsenifar, Z. (1998) Impact of clinical pathways and practice guidelines on the management of acute exacerbations of bronchial asthma. *Chest*, 113, 28–33.
 37. Burns, S. M., Marshall, M., Burns, J. E., Ryan, B., Wilmoth, D., Carpenter, R., Aloia, A., Wood, M. & Truwit, J. D. (1998) Design, testing, and results of an outcomes-managed approach to patients requiring prolonged mechanical ventilation. *American Journal of Critical Care*, 7, 45–57.
 38. Burns, S. M., Earven, S., Fisher, C., Lewis, R., Merrell, P., Schubart, J. R., Truwit, J. D. & Bleck, T. P. (2003) Implementation of an institutional program to improve clinical and financial outcomes of mechanically ventilated patients: one-year outcomes and lessons learned. *Critical Care Medicine*, 31, 2752–2763.
 39. Burns, S. M. & Dempsey, E. (2000) Long-term ventilator management strategies: experiences of two hospitals. *AACN Clinical Issues*, 11, 424–441.
 40. Chin, R., Browne, G. J., Lam, L. T., McCaskill, M. E., Fasher, B. & Hort, J. (2002) Effectiveness of a croup clinical pathway in the management of children with croup presenting to an emergency department. *Journal of Paediatrics and Child Health*, 38, 382–387.
 41. Estrada, C. A., Unterborn, J. N., Price, J., Thompson, D. & Gibson, L. (2000) Judging the effectiveness of clinical pathways for pneumonia: the role of risk adjustment. *Effective Clinical Practice*, 3, 221–228.
 42. Johnson, K. B., Blaisdell, C. J., Walker, A. & Eggleston, P. (2000) Effectiveness of a clinical pathway for inpatient asthma management. *Pediatrics*, 106, 1006–1012.
 43. Kwan-Gett, T. S., Lozano, P., Mullin, K. & Marcuse, E. K. (1997) One-year experience with an inpatient asthma clinical pathway. *Archives of Pediatrics and Adolescent Medicine*, 151, 684–689.
 44. Lee, S. C., Tseng, H. Y., Wang, K. Y. & Lee, L. C. (2002) Effect of a clinical pathway on selected clinical outcomes of pulmonary lobectomy. *Zhonghua Yi Xue Za Zhi (Taipei)*, 65, 7–12.
 45. Marrie, T. J., Lau, C. Y., Wheeler, S. L., Wong, C. J., Vandervoort, M. K. & Feagan, B. G. (2000) A controlled trial of a critical pathway for treatment of community-acquired pneumonia. *JAMA*, 283, 749–755.
 46. Phillips, K. & Crain, H. (1998) Effectiveness of a pneumonia clinical pathway: quality and financial outcomes. *Outcomes Management for Nursing Practice*, 2, 16–23.
 47. Price, B. J., Bernard, G. R., Drew, K., Foss, J. & Wheeler, A. P. (1998) Impact of a critical care pathway for unstable mechanically ventilated patients. *Critical Care Nursing Clinics of North America*, 10, 75–85.
 48. Santamaria, N., Connors, A., Osteraas, J., Ham, J. & Boodram, B. (2004) A prospective cohort study of the effectiveness of clinical pathways for the in-patient management of acute exacerbation of chronic obstructive pulmonary disease (COPD). *Collegian*, 11, 12–16.
 49. Soria, V., Coll, A., Flores, B., Miguel, J., Torralba, J. A., Andres, B., Carrasco, M. & Aguayo, J. L. (2005) Clinical pathway for the treatment of primary spontaneous pneumothorax in a general surgery department. *American Journal of Medical Quality*, 20, 268–276.
 50. Wilson, S. D., Dahl, B. B. & Wells, R. D. (2002) An evidence-based clinical pathway for bronchiolitis safely reduces antibiotic overuse. *American Journal of Medical Quality*, 17, 195–199.
 51. Zehr, K. J., Dawson, P. B., Yang, S. C. & Heitmiller, R. F. (1998) Standardized clinical care pathways for major thoracic cases reduce hospital costs. *Annals of Thoracic Surgery*, 66, 914–919.
 52. Archer, S. B., Burnett, R. J., Flesch, L. V., Hobler, S. C., Bower, R. H., Nussbaum, M. S. & Fischer, J. E. (1997) Implementation of a clinical pathway decreases length of stay and hospital charges for patients undergoing total colectomy and ileal pouch/anal anastomosis. *Surgery*, 122, 699–703.
 53. Bradshaw, B. G., Liu, S. S. & Thirlby, R. C. (1998) Standardized perioperative care protocols and reduced length of stay after colon surgery. *Journal of the American College of Surgeons*, 186, 501–506.
 54. Cooney, R. N., Bryant, P., Haluck, R., Rodgers, M. & Lowery, M. (2001) The impact of a clinical pathway for gastric bypass surgery on resource utilization. *Journal of Surgical Research*, 98, 97–101.
 55. Correa, A. J., Reinisch, L., Paty, V. A., Sanders, D. L. & Duncavage, J. A. (1999) Analysis of a critical pathway in osteoplastic flap for frontal sinus obliteration. *Laryngoscope*, 109, 1212–1216.
 56. Delaney, C. P., Zutshi, M., Senagore, A. J., Remzi, F. H., Hammel, J. & Fazio, V. W. (2003) Prospective, randomized, controlled trial between a pathway of controlled rehabilitation with early ambulation and diet and traditional postoperative care after laparotomy and intestinal resection. *Diseases of the Colon and Rectum*, 46, 851–859.
 57. Joh, H., Moon, I., Park, H., Kim, N. & Yang, S. (2003) The effect of the critical pathway for inguinal hernia repair. *Yonsei Medical Journal*, 44, 81–88.
 58. Kelly, R.-E. J., Wenger, A., Horton, C. J., Nuss, D., Croitoru, D. P. & Pestian, J. P. (2000) The effects of a pediatric unilateral inguinal hernia clinical pathway on quality and cost. *Journal of Pediatric Surgery*, 35, 1045–1048.
 59. Matsumoto, A., Kanda, K. & Shimegatsu, H. (2002) Development and implementation of a critical pathway for abdominal aortic aneurysms in Japan. *Journal of Vascular Nursing*, 20, 14–21.
 60. Melbert, R. B., Kimmins, M. H., Isler, J. T., Billingham, R. P., Lawton, D., Salvadalena, G., Cortezzo, M. & Rowbotham, R. (2002) Use of a critical pathway for colon resections. *Journal of Gastrointestinal Surgery*, 6, 745–752.
 61. Michalsky, M. P., Pratt, D., Caniano, D. A. & Teich, S. (2002) Streamlining the care of patients with hypertrophic pyloric stenosis: application of a clinical pathway. *Journal of Pediatric Surgery*, 37, 1072–1075.
 62. Pfau, P. R., Cooper, G. S., Carlson, M. D., Chak, A., Sivak, M. V. Jr, Gonet, J. A., Boyd, K. K. & Wong, R. C. (2004) Success and shortcomings of a clinical care pathway in the management of acute nonvariceal upper gastrointestinal bleeding. *American Journal of Gastroenterology*, 99, 425–431.
 63. Podila, P. V., Ben Menachem, T., Batra, S. K., Oruganti, N., Posa, P. & Fogel, R. (2001) Managing patients with acute, nonvariceal gastrointestinal hemorrhage: development and effectiveness of a clinical care pathway. *American Journal of Gastroenterology*, 96, 208–219.
 64. Porter, G. A., Pisters, P. W., Mansur, C., Bisanz, A., Reyna, K., Stanford, P., Lee, J. E. & Evans, D. B. (2000) Cost and utilization impact of a clinical pathway for patients undergoing pancreaticoduodenectomy. *Annals of Surgical Oncology*, 7, 484–489.
 65. Pritts, T. A., Nussbaum, M. S., Flesch, L. V., Fogelman, E. J., Parikh, A. A. & Fischer, J. E. (1999) Implementation of a clinical pathway decreases length of stay and cost for bowel resection. *Annals of Surgery*, 230, 728–733.
 66. Stephen, A. E. & Berger, D. L. (2003) Shortened length of stay and hospital cost reduction with implementation of an accelerated clinical care pathway after elective colon resection. *Surgery*, 133, 277–282.
 67. Takegami, K., Kawaguchi, Y., Nakayama, H., Kubota, Y. & Nagawa, H. (2003) Impact of a clinical pathway and standardization of treatment for acute appendicitis. *Surgery Today*, 33, 336–341.

68. Uchiyama, K., Takifuji, K., Tani, M., Onishi, H. & Yamaue, H. (2002) Effectiveness of the clinical pathway to decrease length of stay and cost for laparoscopic surgery. *Surgical Endoscopy*, 16, 1594–1597.
69. Walsh, M. D., Barry, M., Scott, T. E., Lamorte, W. W. & Menzoian, J. O. (2001) The role of a nurse case manager in implementing a critical pathway for infrainguinal bypass surgery. *Joint Commission Journal on Quality Improvement*, 27, 230–238.
70. Warner, B. W., Kulick, R. M., Stoops, M. M., Mehta, S., Stephan, M. & Kotagal, U. R. (1998) An evidenced-based clinical pathway for acute appendicitis decreases hospital duration and cost. *Journal of Pediatric Surgery*, 33, 1371–1375.
71. Choong, P. F., Langford, A. K., Dowsey, M. M. & Santamaria, N. M. (2000) Clinical pathway for fractured neck of femur: a prospective, controlled study. *Medical Journal of Australia*, 172, 423–426.
72. Dowsey, M. M., Kilgour, M. L., Santamaria, N. M. & Choong, P. F. (1999) Clinical pathways in hip and knee arthroplasty: a prospective randomised controlled study. *Medical Journal of Australia*, 170, 59–62.
73. Gholive, P. A., Kosygan, K. P., Sturdee, S. W. & Faraj, A. A. (2005) Multidisciplinary integrated care pathway for fractured neck of femur. A prospective trial with improved outcome. *Injury*, 36, 93–98.
74. Ireson, C. L. (1997) Critical pathways: effectiveness in achieving patient outcomes. *Journal of Nursing Administration*, 27, 16–23.
75. Koval, K. J., Chen, A. L., Aharonoff, G. B., Egol, K. A. & Zuckerman, J. D. (2004) Clinical pathway for hip fractures in the elderly: the Hospital for Joint Diseases experience. *Clinical Orthopaedics*, 425, 72–81.
76. Mabrey, J. D., Toohey, J. S., Armstrong, D. A., Lavery, L. & Wammack, L. A. (1997) Clinical pathway management of total knee arthroplasty. *Clinical Orthopaedics*, 345, 125–133.
77. Macario, A., Horne, M., Goodman, S., Vitez, T., Dexter, F., Heinen, R. & Brown, B. (1998) The effect of a perioperative clinical pathway for knee replacement surgery on hospital costs. *Anesthesia and Analgesia*, 86, 978–984.
78. Pearson, S., Moraw, I. & Maddern, G. J. (2000) Clinical pathway management of total knee arthroplasty: a retrospective comparative study. *Australian and New Zealand Journal of Surgery*, 70, 351–354.
79. Pennington, J. M., Jones, D. P. & McIntyre, S. (2003) Clinical pathways in total knee arthroplasty: a New Zealand experience. *Journal of Orthopedic Surgery*, 11, 166–173.
80. Roberts, H. C., Pickering, R. M., Onslow, E., Clancy, M., Powell, J., Roberts, A., Hughes, A., Coulson, D. & Bray, J. (2004) The effectiveness of implementing a care pathway for femoral neck fracture in older people: a prospective controlled before and after study. *Age Ageing*, 33, 178–184.
81. Santamaria, N., Houghton, L., Kimmel, L. & Graham, A. (2003) Clinical pathways for fractured neck of femur: a cohort study of health related quality of life, patient satisfaction, and clinical outcome. *Australian Journal of Advanced Nursing*, 20, 24–29.
82. Wammack, L. & Mabrey, J. D. (1998) Outcomes assessment of total hip and total knee arthroplasty: critical pathways, variance analysis, and continuous quality improvement. *Clinical Nurse Specialist*, 12, 122–129.
83. Sesperez, J., Wilson, S., Jalaludin, B., Seger, M. & Sugrue, M. (2001) Trauma case management and clinical pathways: prospective evaluation of their effect on selected patient outcomes in five key trauma conditions. *Journal of Trauma*, 50, 643–649.
84. Chang, S. S., Smith, J. A. Jr., Girasole, C., Baumgartner, R. G., Roth, B. J. & Cookson, M. S. (2002) Beneficial impact of a clinical care pathway in patients with testicular cancer undergoing retroperitoneal lymph node dissection. *Journal of Urology*, 168, 87–92.
85. Gendron, K. M., Lai, S. Y., Weinstein, G. S., Chalian, A. A., Husbands, J. M., Wolf, P. F., Di Donato, L. & Weber, R. S. (2002) Clinical care pathway for head and neck cancer: a valuable tool for decreasing resource utilization. *Archives of Otolaryngology-Head and Neck Surgery*, 128, 258–262.
86. Hanna, E., Schultz, S., Doctor, D., Vural, E., Stern, S. & Suen, J. (1999) Development and implementation of a clinical pathway for patients undergoing total laryngectomy: impact on cost and quality of care. *Archives of Otolaryngology-Head and Neck Surgery*, 125, 1247–1251.
87. Hwang, T. G., Wilkins, E. G., Lowery, J. C. & Gentile, J. (2000) Implementation and evaluation of a clinical pathway for TRAM breast reconstruction. *Plastic and Reconstructive Surgery*, 105, 541–548.
88. Levin, R. J., Ferraro, R. E., Kodosky, S. R. & Fedok, F. G. (2000) The effectiveness of a 'critical pathway' in the management of laryngectomy patients. *Head and Neck-Journal for the Sciences and Specialties of the Head and Neck*, 22, 694–699.
89. Ramanujam, L. N. & Cheah, W. K. (2005) Improvements in health care for patients undergoing thyroidectomy. *Asian Journal of Surgery*, 28, 266–270.
90. Santoso, U., Iau, P. T., Lim, J., Koh, C. S. & Pang, Y. T. (2002) The mastectomy clinical pathway: what has it achieved? *Annals of the Academy of Medicine*, 31, 440–445.
91. Sherman, D., Matthews, T. W., Lampe, H. & LeBlanc, S. (2001) Laryngectomy clinical pathway: development and review. *Journal of Otolaryngology*, 30, 115–120.
92. Yueh, B., Weaver, E. M., Bradley, E. H., Krumholz, H. M., Heagerty, P., Conley, A. & Sasakii, C. T. (2003) A critical evaluation of critical pathways in head and neck cancer. *Archives of Otolaryngology-Head and Neck Surgery*, 129, 89–95.
93. Dufault, M. A. & Willey-Lessne, C. (1999) Using a collaborative research utilization model to develop and test the effects of clinical pathways for pain management. *Journal of Nursing Care Quality*, 13, 19–33.
94. Kwan, J., Hand, P., Dennis, M. & Sandercock, P. (2004) Effects of introducing an integrated care pathway in an acute stroke unit. *Age Ageing*, 33, 362–367.
95. Spain, D. A., McIlvoy, L. H., Fix, S. E., Carrillo, E. H., Boaz, P. W., Harpring, J. E., Rague, G. H. & Miller, F. B. (1998) Effect of a clinical pathway for severe traumatic brain injury on resource utilization. *Journal of Trauma*, 45, 101–104.
96. Sulch, D., Evans, A., Melbourn, A. & Kalra, L. (2002) Does an integrated care pathway improve processes of care in stroke rehabilitation? A randomized controlled trial. *Age Ageing*, 31, 175–179.
97. Sulch, D., Melbourn, A., Perez, I. & Kalra, L. (2002) Integrated care pathways and quality of life on a stroke rehabilitation unit. *Stroke*, 33, 1600–1604.
98. Sulch, D., Perez, I., Melbourn, A. & Kalra, L. (2000) Randomized controlled trial of integrated (managed) care pathway for stroke rehabilitation. *Stroke*, 31, 1929–1934.
99. Vitaz, T. W., McIlvoy, L., Raque, G. H., Spain, D. A. & Shields, C. B. (2001) Development and implementation of a clinical pathway for spinal cord injuries. *Journal of Spinal Disorders*, 14, 271–276.
100. Vitaz, T. W., McIlvoy, L., Raque, G. H., Spain, D. & Shields, C. B. (2001) Development and implementation of a clinical pathway for severe traumatic brain injury. *Journal of Trauma*, 51, 369–375.
101. Aragon, D., Burton, V., Byers, J. F. & Cohen, M. (2002) The effect of a critical pathway on patients' outcomes after carotid endarterectomy. *American Journal of Critical Care*, 11, 250–258.
102. Calligaro, K. D., Doerr, K. J., McAfee-Bennett, S., Mueller, K. & Dougherty, M. J. (2004) Critical pathways can improve results with carotid endarterectomy. *Seminars in Vascular Surgery*, 17, 253–256.
103. Calligaro, K. D., Dougherty, M. J., Raviola, C. A., Musser, D. J. & DeLaurentis, D. A. (1995) Impact of clinical pathways on hospital costs and early outcome after major vascular surgery. *Journal of Vascular Surgery*, 22, 649–657.

104. Stanley, A. C., Barry, M., Scott, T. E., Lamorte, W. W., Woodson, J. & Menzoian, J. O. (1998) Impact of a critical pathway on postoperative length of stay and outcomes after infrainguinal bypass. *Journal of Vascular Surgery*, 27, 1056–1064.
105. Sweeney, A. B., Flora, H. S., Chaloner, E. J., Buckland, J., Morrice, C. & Barker, S. G. (2002) Integrated care pathways for vascular surgery: an analysis of the first 18 months. *Postgraduate Medical Journal*, 78, 175–177.
106. Brandsma, C., Calhoun, B. C. & Vannatta, J. E. (2000) Uncomplicated pregnancy: clinical pathway genesis based on the nursing process. *Military Medicine*, 165, 839–843.
107. Broder, M. S. & Bovone, S. (2002) Improving treatment outcomes with a clinical pathway for hysterectomy and myomectomy. *Journal of Reproductive Medicine*, 47, 999–1003.
108. Chang, W. C. & Lin, C. C. (2003) A clinical pathway for laparoscopically assisted vaginal hysterectomy. Impact on costs and clinical outcome. *Journal of Reproductive Medicine*, 48, 247–251.
109. Chang, W. C., Lee, C. C., Wu, H. C. & Yeh, L. S. (2003) Laparoscopy-assisted vaginal hysterectomy clinical pathway. A multivariate analysis of impact on costs and quality of care. *Gynecologic and Obstetric Investigation*, 55, 231–234.
110. Ghosh, K., Downs, L. S., Padilla, L. A., Murray, K. P., Twiggs, L. B., Letourneau, C. M. & Carson, L. F. (2001) The implementation of critical pathways in gynecologic oncology in a managed care setting: a cost analysis. *Gynecologic Oncology*, 83, 378–382.
111. Morris, M., Levenback, C., Burke, T. W., Dejesus, Y., Lucas, K. R., Gershenson, D. M. (1997) An outcomes management program in gynecologic oncology. *Obstetrics and Gynecology*, 89, 485–492.
112. Chang, P. L., Wang, T. M., Huang, S. T., Hsieh, M. L., Tsui, K. H. & Lai, R. H. (1999) Effects of implementation of 18 clinical pathways on costs and quality of care among patients undergoing urological surgery. *Journal of Urology*, 161, 1858–1862.
113. Chang, P. L., Wang, T. M., Huang, S. T., Hsieh, M. L., Chuang, Y. C. & Chang, C. H. (2000) Improvement of health outcomes after continued implementation of a clinical pathway for radical nephrectomy. *World Journal of Urology*, 18, 417–421.
114. Chang, P. L., Lee, S. H., Hsieh, M. L., Huang, S. T., Tsui, K. H. & Lai, R. H. (2002) Improvement of practice performance in urological surgery via clinical pathway implementation. *World Journal of Urology*, 20, 213–218.
115. Holtzman, J., Bjerke, T. & Kane, R. (1998) The effects of clinical pathways for renal transplant on patient outcomes and length of stay. *Medical Care*, 36, 826–834.
116. Leibman, B. D., Dillioglulugil, O., Abbas, F., Tanli, S., Kattan, M. W. & Scardino, P. T. (1998) Impact of a clinical pathway for radical retropubic prostatectomy. *Urology*, 52, 94–99.
117. Katon, W. J., Von Korff, M., Lin, E. H., Simon, G., Ludman, E., Russo, J., Ciechanowski, P., Walker, E. & Bush, T. (2004) The pathway study: a randomized trial of collaborative care in patients with diabetes and depression. *Archives of General Psychiatry*, 61, 1042–1049.
118. Kazui, H., Hashimoto, M., Nakano, Y., *et al.* (2004) Effectiveness of a clinical pathway for the diagnosis and treatment of dementia and for the education of families. *International Journal of Geriatric Psychiatry*, 19, 892–897.
119. Ilag, L. L., Kronick, S., Ernst, R. D., Grondin, L., Alaniz, C., Liu, L. & Herman, W. H. (2005) Impact of a critical pathway on inpatient management of diabetic ketoacidosis. *Diabetes Research and Clinical Practice*, 62, 23–32.
120. Browne, G. J., Giles, H., McCaskill, M. E., Fasher, B. J. & Lam, L. T. (2001) The benefits of using clinical pathways for managing acute paediatric illness in an emergency department. *Journal of Quality in Clinical Practice*, 21, 3–55.
121. Jones, M. L. & Smyth, K. A. (1999) Outcomes for high-risk neonates in a managed care clinical system. *Nursing Case Management*, 4, 71–76.
122. Pestian, J. P., Derkay, C. S. & Ritter, C. (1998) Outpatient tonsillectomy and adenoidectomy clinical pathways: an evaluative study. *American Journal of Otolaryngology*, 19, 45–49.
123. Dzwierzynski, W. W., Spitz, K., Hartz, A., Guse, C. & Larson, D. L. (1998) Improvement in resource utilization after development of a clinical pathway for patients with pressure ulcers. *Plastic and Reconstructive Surgery*, 102, 2006–2011.
124. Taylor, K. O., Goudie, C. M. & Muller, M. J. (2004) Evaluation of a pediatric scald burn clinical pathway. *Journal of Burn Care and Rehabilitation*, 25, 256–261.
125. Chen, A. Y., Callender, D., Mansyur, C., Reyna, K. M., Limitone, E. & Goepfert, H. (2000) The impact of clinical pathways on the practice of head and neck oncologic surgery: the University of Texas M. D. Anderson Cancer Center Experience. *Archives of Otolaryngology-Head and Neck Surgery*, 126, 322–326.
126. Husbands, J. M., Randal, S. W., Karpati, R. L., Weinstein, G. S., Chalian, A. A., Goldberg, A. N., Thaler, E. R. & Wolf, R. F. (1999) Clinical care pathways: decreasing resource utilization in head and neck surgical patients. *Otolaryngology and Head and Neck Surgery*, 121, 755–759.
127. Philbin, E. F., Rocco, T. A., Lindenmuth, N. W., Ulrich, K., McCall, M. & Jenkins, P. L. (2000) The results of a randomized trial of a quality improvement intervention in the care of patients with heart failure. The MISCHF Study Investigators. *American Journal of Medicine*, 109, 443–449.
128. Wild, D., Nawaz, H., Chan, W. & Katz, D. L. (2004) Effects of interdisciplinary rounds on length of stay in a telemetry unit. *Journal of Public Health Management and Practice*, 10, 63–69.
129. Ayestas, A. L., Diaz, E. & Kirtland, S. (1995) Clinical pathways: improving patient education and influencing readmission rates. *Journal of Healthcare Quality*, 17, 17–25.
130. Zielhuis, G. A., Straatman, H., Van't Hof-Grootenboer, A. E., Vanlier, H. J. J., Rach, G. H. & Van den Broek, P. (1990) The choice of a balanced allocation method for a clinical-trial in otitis-media with effusion. *Statistics in Medicine*, 9, 237–246.