University of Groningen

HRT use in 2001 and 2004 in The Netherlands

de Jong-van den Berg, L.T.W.; Faber, A.; van den Berg, Paulus

Published in:
Maturitas

DOI:
[10.1016/j.maturitas.2005.10.010](https://doi.org/10.1016/j.maturitas.2005.10.010)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2006

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

de Jong-van den Berg, L. T. W., Faber, A., & van den Berg, P. (2006). HRT use in 2001 and 2004 in The Netherlands: A world of difference. *Maturitas*, 54(2), 193-197.
<https://doi.org/10.1016/j.maturitas.2005.10.010>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

HRT use in 2001 and 2004 in The Netherlands— A world of difference

Lolkje T.W. de Jong-van den Berg*, Adrienne Faber, Paul B. van den Berg

*Groningen University Institute for Drug Exploration (GUIDE), University of Groningen, Department of Social Pharmacy,
Pharmacoepidemiology and Pharmacotherapy, Antonius Deusinglaan 1, 9713 AV Groningen, The Netherlands*

Received 5 August 2005; received in revised form 13 October 2005; accepted 14 October 2005

Abstract

Objective: Did the publication of the Women Health Initiative (WHI) trial in 2002 and the Million Women Study (MWS) in 2003 lead to changes in prescription rates of hormone replacement therapy (HRT). Therefore, we compare the prescribing of HRT in 2004 (after) with that of 2001 (before the publications) in The Netherlands.

Method: Community pharmacy dispensing data from a population of approximately 500,000 patients in The Netherlands. Women aged 40–74 years to whom at least one HRT prescription was dispensed in 2001 or 2004 were included. Annual prevalences of HRT in 2001 and 2004 and the percentage change (2004 versus 2001) were calculated for overall HRT (excluding vaginal products) and per HRT category (combined estrogens and progestagens, estrogens only, tibolon and vaginal preparations) and age category.

Results: In 2001, 5.64% of the women aged 40–74 used HRT and this percentage declined to 2.39 in 2004. The use of vaginal products among these women did not change, 1.76% in 2001 and 1.65% in 2004. The percentage change was highest in the opposed HRT group (66% decrease) and in women aged 50–54 (64.4% decrease). In 2004, compared with 2001, the proportion of long-term users (>3 year) increased with 12.7%.

Conclusions: In The Netherlands, after publication of the WHI study and the MWS the prescribing of HRT fell dramatically whereas the prescribing of vaginal products did not change. Future patterns of HRT use should be monitored to know whether this decrease will be sustained.

© 2005 Elsevier Ireland Ltd. All rights reserved.

Keywords: Hormone replacement therapy (HRT); Change in prescribing; Low potency estrogens/vaginal products; Opposed HRT; Unopposed HRT; Tibolon

1. Introduction

In The Netherlands, hormone replacement therapy (HRT) is licensed for the alleviation of climacterial symptoms and the prevention of osteoporosis, however in daily practice, it was mainly prescribed for treatment

DOI of the original article: [10.1016/j.maturitas.2005.08.002](https://doi.org/10.1016/j.maturitas.2005.08.002).

* Corresponding author. Tel.: +31 50 3633330.

E-mail address: l.t.w.de.jong-van.den.berg@rug.nl
(L.T.W. de Jong-van den Berg).

of vasomotor symptoms. HRT preparations used for this purpose are either combined estrogens and progestagens (opposed) or estrogens alone (unopposed) or tibolon. Vaginal applications, containing estriol or dienestrel, are licensed for the relief of urogenital symptoms.

Before 2002, HRT had been regarded as an intervention with great potential benefits in terms of cardiovascular disease prevention and osteoporosis treatment [1–3]. The publications of the Women Health Initiative (WHI) trial results [4–6] and the results of the observational Million Women Study (MWS) [7] reported increased rates of breast cancer, coronary heart diseases, stroke, dementia and venous thromboembolism and decreased rates of hip fractures and colorectal cancer in postmenopausal women using long-term HRT. Both studies led to much attention and debate in the medical and the lay press.

As a consequence, shortly after the publication of the MWS the Dutch associations of gynaecologists and general practitioners clearly stated that HRT should be prescribed only in women with severe vasomotoric complaints in the lowest effective dose and for a short period. According to the general practitioners this period should not exceed 6 months and the recommendations of the gynecologists stated that the treatment should be evaluated every 6–12 months [8,9]. The European Medicines Agency for Drug Safety (EMA) stated in a position paper that for the initiation and continuation of HRT, the minimum effective dose for the shortest duration should be used. And in all cases, a careful appraisal of the risks and benefits should be undertaken at least annually and HRT should only be continued as long as benefits outweighs risks [10].

The aim of this report is to compare the use of HRT in The Netherlands in 2004, after publication of the WHI trial and the MWS results, with the use in 2001, the year before these publications. In addition, we compared the duration of use among HRT-users before and after publication of both studies.

2. Methods

This study was performed with the InterAction Database (IADB), which contains prescription drug dispensing data from community pharmacies in the northern and eastern part of The Netherlands. The

IADB covers all prescriptions from an estimated population of approximately 500,000 since 1999 [11,12]. Each prescription record contains among others name of the drug, ATC-code, date of dispensing and amount dispensed. Each patient has a unique, though anonymous identifier. Date of birth and gender of patients are available. Due to a high patient–pharmacy commitment in The Netherlands and sophisticated pharmacy software, the medication records for each patient are virtually complete [13]. This database comprises all prescriptions, regardless of insurance or reimbursement status, apart from drugs dispensed during hospitalisations. Note that almost all HRT preparations are fully reimbursed. Only for transdermal oestrogen/progestagen preparations a patient's copayment is required.

2.1. Study population and design

All women aged 40–74 years, to whom at least one HRT prescription was dispensed in 2001 or in 2004 were selected from the IADB. Women who received HRT prescription were classified into four categories: (I) those who received conjugated estrogens or estradiol and progestagens either in a fixed combination or separately (opposed), (II) those who only received estrogens (conjugated estrogens or estradiol) (unopposed), (III) women who got tibolon and (IV) women who received low potency estrogens (estriol or dienestrel) mainly a vaginal application for vulvar vaginal atrophy. Tibolon is a synthetic steroid with weak estrogenic, progestogenic and androgenic properties [14]. Prevalence of HRT prescribing was estimated per year (2001 and 2004) and was defined as the number of women (aged 40–74) to whom any HRT prescription was dispensed per 100 women in the population covered by the IADB. Annual prevalence was stratified per HRT category and 5-years age categories. The percentage change, defined as: [(prevalence 2001 minus prevalence 2004) divided by prevalence 2001 multiplied by 100], was calculated and stratified per HRT category.

For calculating the duration of use, we selected in 2001 as well as in 2004 all HRT-users with at least 3-year medication history in the database before the first prescription in 2001 or 2004. Of these HRT-users we calculated the differences in proportion (2004 compared to 2001) in relation to duration of use, being: less

than 1 year, between 1 and 3 years, and more than 3-year use.

For comparing prevalences of HRT use in 2004 with 2001 we used Pearson Chi-square.

3. Results

In 2001, of the 91,873 women aged 40–74, 5182 (5.64%) received at least a HRT preparation, licenced for symptomatic treatment of vasomotor symptoms or the prevention of osteoporosis and 1618 (1.67%) received at least a vaginal low oestrogen product for vaginal atrophy. In 2004, these prevalences were 2.39 and 1.65, respectively. The prevalence in the different age categories of these two groups is shown in Fig. 1. A dramatic decrease in the prescribing of HRT is seen in all age categories, being the highest among the 50–54 (64.4% decrease) and the lowest among women aged 65–69 (40.3% decrease). For the low potency estrogens the prevalence of use did not change between 2001 and 2004. These products, aimed for vaginal atrophy, are prescribed to older women as can be seen in the figure.

Table 1 presents the percentage change in HRT use for overall HRT (all categories, excluding vaginal products), and per HRT category; combined estrogens and progestagens, estrogens only, tibolone and vaginal preparations. Between 2004 and 2001, the overall HRT prescribing decreased with 57.6%, whereas the % change for the low-potency estrogens is much lower (−6.3%) and not significantly different ($p=0.055$). Among the HRT-users the prescribing of opposed

HRT decreased most (66.0%), while the prevalence of tibolone changed from 0.71 in 2001 to 0.45 in 2004 (36.6% decrease).

Table 2 shows the differences in proportion (2004 compared with 2001) among HRT-users in relation to duration of use. Although the absolute numbers of HRT-users dropped from 4194 in 2001 to 2130 in 2004, the proportion of long-term users (>3 year) increased with 12.7% in 2004 compared with 2001. The changes in proportions of women who used HRT less than 1 year, and those who used it between 1 and 3 year were lower, −8 and −4.7%, respectively.

4. Discussion

Between 2001 and 2004, the prevalence of HRT use declined from 5.64 to 2.39% among women aged 40–74, whereas the use of vaginal products containing low potency estrogens did not change at all (1.76 and 1.65%, respectively). The decline in use was most pronounced among women who were prescribed combined estrogens and progestagens (65% decrease) and estrogens only (54.2% decrease). The proportion of long-term users among HRT-users increased with 12.7% in 2004 compared with 2001.

Before the release of the publications of WHI and MWS the prescribing of HRT (including vaginal applications) in The Netherlands among women aged 45–69 was 9.2% [15] and this is much lower than prevalences reported in the UK and USA. Data from the UK show a prevalence of 27.7% in 1998 among women aged 45–64 [16] and in the US these percentages vary between 38 and 50% [17,18]. In contrast with the US and the UK, HRT treatment for the prevention of cardiovascular disease and osteoporosis with consequent long-term treatment duration was never prevalent in The Netherlands. Dutch prescribers have always been very conservative in prescribing HRT, resulting in 2004 in a very low prevalence of use: 2.39 among 40–74 year olds and 2.95 in women aged 45–69.

A decline in HRT use is also evaluated with prescription claims data in the US [18,19]. Hersh et al. showed a 37% decrease of HRT prescriptions in 2003 compared with 1999 but in another study in a lower educated population (Medicaid program) the decline was less pronounced [19]. In a survey among HRT-users in New Zealand 40% stopped the use of HRT [20]. Our results

Fig. 1. Prevalences of HRT prescribing per age category in 2001 and 2004 for general HRT (combined, estrogen only and tibolone, excluding vaginal HRT) and for vaginal HRT prescribing.

Table 1

Prevalences of HRT in women aged 40–74 per HRT category and the % change in 2004 compared with 2001

	Prevalence among women aged 40–74 years			
	2001 (<i>n</i> = 91873), <i>N</i> (prevalence)	2004 (<i>n</i> = 98401), <i>N</i> (prevalence)	Change (%) ^b	<i>p</i> -Value ^a
All HRT (excluding vaginal)	5182 (5.64)	2351 (2.39)	–57.62	<0.001
• Opposed HRT	2559 (2.79)	939 (0.95)	–65.95	<0.001
• Oestrogens only	2069 (2.25)	1018 (1.03)	–54.22	<0.001
• Tibolon	652 (0.71)	440 (0.45)	–36.62	<0.001
Low-potency estrogens	1618 (1.76)	1621 (1.65)	–6.25	0.055

^a *p*-Value indicates whether prevalences differ between 2001 and 2004 using Pearson Chi-square.^b % Change: $\frac{\text{prevalence 2001} - \text{prevalence 2004}}{\text{prevalence 2004}}$.

Table 2

Number of HRT-users in women aged 40–74 according to the duration of use (<1, 1–3 and >3 year) in 2001 and 2004, and the difference in proportion in 2004 compared with 2001

Duration of use	Number of HRT (excluding vaginal HRT) users (40–74 year) (at least 3 years data available)		
	2001 <i>N</i> (%)	2004 <i>N</i> (%)	Difference in proportion 2004 vs. 2001 (%)
<1 year	1179 (28.1)	429 (20.1)	–8.0
Between 1 and 3 years	850 (20.3)	332 (15.6)	–4.7
>3 years	2165 (51.6)	1369 (64.3)	+12.7
Overall	4194 (100)	2130 (100)	

show a decline of 57.6% among women aged 40–74 who received HRT (excluding vaginal preparations). The different percentage decline seen in the studies can be explained by the method used, the age groups involved and the definition of the HRT. As is shown in this study, the decline among the different age categories varied from 64% (50–54 years) to 40% (65–69 years). Studies among different age groups will reveal different percentages change. We also illustrated that the use of vaginal estrogen products, used for vaginal atrophy, did change only marginally (–6%), whereas the decline of opposed HRT in the same age category (40–74 years) declined drastically (66%). Combining both types of HRT will consequently result in lower percentages change.

Although the Dutch recommendations stated that HRT should be used only in women with severe complaints and during of short time, we see in 2004 among HRT-users a greater proportion of long-term users (>3 year) 64.3% compared with 51.6% in 2001, whereas the proportion of short-term users (<1 year and between 1–3 years) declined. One explanation may relate to the beneficial effects in women with severe menopausal complaints. A recent Dutch survey showed that short-

term considerations prevail in deciding to continue or stop the use of HRT for women who used the medication for a longer time [21].

In summary, we described a 57.6% decrease in the prevalence of HRT-users in 2004 compared with 2001. This 57.6% decrease in a population with an existing low prevalence of HRT use resulted in 2004 in an overall prevalence of less than 3%. The prevalence of vaginal low-potency estrogens did not change in the same population. We will monitor future patterns of HRT use to know whether this decrease will be sustained.

References

- [1] Grodstein F, Stampfer MJ, Manson JE, et al. Postmenopausal estrogen and progestin use and the risk of cardiovascular disease. *N Engl J Med* 1996;335:453–61.
- [2] Pan CX. Hormone replacement therapy for secondary prevention of coronary heart disease. *JAMA* 1999;281:794.
- [3] Barrett-Connor E, Grady D. Hormone replacement therapy, heart disease, and other considerations. *Annu Rev Pub Health* 1998;19:55–72.
- [4] Writing group for the Women's Health Initiative Investigators. Risks and benefits of estrogen plus progestin in

- healthy postmenopausal women. Principal results from the Women's Health Initiative randomized controlled trial. *JAMA* 2002;288:321–33.
- [5] Shumaker SA, Legault C, Rapp SR, et al. Estrogen plus progestin and the incidence of dementia and mild cognitive impairment in postmenopausal women. The Women's Health Initiative Memory study: a randomized controlled trial. *JAMA* 2003;289:2651–62.
- [6] Writing group for the Women's Health Initiative Investigators. Effects of conjugated equine estrogen in postmenopausal women with hysterectomy. *JAMA* 2004;291:1701–12.
- [7] Million Women Study Collaborators. Breast cancer and hormone-replacement therapy in the Million Women Study. *Lancet* 2003;362:419–27.
- [8] Dutch College of General Practitioners (website: <http://nhg.artsennet.nl>, accessed on November 8, 2004).
- [9] Dutch Society of Obstetrics and Gynaecology (Press release on website: www.nvog.nl, accessed on November 8, 2004).
- [10] European Medicines Agency, Evaluation of Medicines for Human Use. Guidelines for hormone replacement therapy (on website: www.emea.eu.int, accessed on October, 2005).
- [11] Tobi H, van den Berg PB, de Jong-van den Berg LTW. The interaction database: synergy of science and practice in pharmacy. In: Brause RW, Hanisch E, editors. *Medical data analysis*. Berlin: Springer-Verlag; 2000. p. 206–11.
- [12] Schirm E, Monster TBM, de Vries R, van den Berg PB, de Jong-van den Berg LTW, Tobi H. How to estimate the population that is covered by community pharmacies? An evaluation of two methods using drug utilisation information. *Pharmacoepidemiol Drug Saf* 2004;13:173–9.
- [13] Leufkens HGM, Urquhart J. Automated pharmacy record linkage in The Netherlands. In: Strom BL, editor. *Pharmacoepidemiology*. 3rd ed. 2000. p. 347–60.
- [14] Albertazzi P, Di Micco R, Zanardi E. Tibolone: a review. *Maturitas* 1998;30:295–305.
- [15] Tobi H, van den Berg PB, Brouwers JRB, de Jong-van den Berg LTW. Hormone replacement therapy in peri- and postmenopausal women: more than half use HRT longer than 1 year. *Ned Tijdschr Geneesk* 2003;147:1853–5 (in Dutch).
- [16] Bromley SE, de Vries CS, Farmer RDT. Utilisation of hormone replacement therapy in the United Kingdom: a descriptive study using the GRPD. *BJOG* 2004;111:369–76.
- [17] Keating NL, Cleary PD, Rossi AS, Zaslavsky AM, Aynian JZ. Use of hormone replacement therapy by postmenopausal women in the United States. *Ann Intern Med* 1999;130:545–53.
- [18] Hersh AL, Ste MJ, Stafford RS. National use of postmenopausal hormone therapy. Annual trends and response to recent evidence. *JAMA* 2004;291:47–53.
- [19] Lawton B, Rose S, McLeod D, Dowell A. Changes in use of hormone replacement therapy after the report from the Women's Health Initiative: cross sectional survey of users. *BMJ* 2003;327:845–6.
- [20] Hillman JJ, Zuckerman IH, Lee E. The impact of the WHI on hormone replacement therapy in a medicais program. *J Women's Health* 2004;13:986–92.
- [21] Carton D, Snelder V, Moerman C, Verbeek-Heida P. Why do women continue their long-term use of postmenopausal hormone replacement therapy (HRT): a qualitative study. *Huisarts Wet* 2004;47:497–501 (in Dutch).