University of Groningen

Concerns of terminally ill cancer patients in their home environment

Rolf, C.A.M.; Pool, G.

Published in:
Palliative Medicine

DOI:
[10.1191/0269216306pm1155xx](https://doi.org/10.1191/0269216306pm1155xx)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2006

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Rolf, C. A. M., & Pool, G. (2006). Concerns of terminally ill cancer patients in their home environment. *Palliative Medicine*, 20(4), 481-482. <https://doi.org/10.1191/0269216306pm1155xx>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Letter to the Editor

Concerns of terminally ill cancer patients in their home environment

The concerns of Dutch terminally ill cancer patients who stayed in their home environment were explored, and in this 'Letter to the editor', we communicate some of our findings. In order to relate the actual *physical state* of the patients to their *concerns*, we constructed a relatively short self-report questionnaire of 19 items (partly open questions, partly using a Visual-analogue scale 0–10), including the WHO-functional scale. A total of 280 general practices in the northern part of the country were enlisted to approach patients with a life-expectancy of less than six months. The average general practice in the Netherlands takes care of about three to four terminally ill patients per year.¹ Therefore, during the recruitment period of seven months, about 500 patients were expected to meet our inclusion criteria. In principle, every Dutch inhabitant has access to basic home care. In actuality, a total of 51 patients were included (by 34 GPs); 34 questionnaires were used for analysis (respondents: 14 male (mean age: 63.9, SD: 9.2); 20 female (mean age: 67.5, SD: 10.5); median time since diagnosis: 14 months; 28 patients (82%) lived with a partner). Six patients died before completion of the questionnaire, 11 questionnaires were incomplete.

Concerning the functional state, about half of the 34 patients was partially/totally confined to bed and partially/totally disabled. A majority, 26 patients (76%) suffered from two or more physical problems. Pain was most frequently mentioned ($n=23$, 61%), followed by fatigue ($n=13$, 39%). A majority of the patients reported *concerns* about the further *physical* course of the disease ($n=22$, 60%), as well as *non-physical* concerns ($n=23$, 68%, eg, concerns about partner, children, losing control, loss of independence, not being able to say good-bye). Patients with a WHO-functional state of 3–4 reported significantly more concerns (physical as well as non-physical) than patients with a score of 0–2 ($P=0.04$). Almost half of the patients ($n=16$, 47%) reported a threat to their dignity. Important aspects concerning 'maintenance of dignity' were: being able to say goodbye; dying at home; dying without symptoms; dying in peace; being clear of mind; being able to control the end (by euthanasia). Interestingly, ten patients (29%) stated they had *no* concerns at all (physical or non-physical). Furthermore, the majority ($n=28$, 82%) reported *not* being afraid of dying.

Of course, the results of our study permit only limited conclusions. Although many patients were approached, only 10% of the potential population ($n=51$) was included. This confirms the structural difficulty of including terminal patients in a study.² However, this study evokes several considerations.

A high level of physical complaints in the terminally ill, as reported here and by others,^{3–5} may indicate that, for many patients, symptoms are not under *optimal* (medical) control. Another notable point from our study is that not dying as such, but the *uncertainty* about the dying process was related to the severity of actual physical symptoms. This was also mentioned in other recent studies.^{6,7} (Although Heaven and Maguire,⁸ reported that loss of *independence* seems to be the major concern in terminally ill, we think this may be due to the fact that their study was performed in a *hospice* population, where symptom control may be more optimal than in home situations, and the threat of losing control and independence may be greater. A wish for euthanasia (mentioned a few times by patients), or the wish to die at home, can be viewed as an attempt to maintain personal control. Of course, cultural aspects may influence patients' perceptions on suffering, dying, death and euthanasia.)

In conclusion, we believe our exploratory study stresses the high value of symptom control. A lack of (feelings of) symptom-control may *evoke* concerns regarding the course of the disease, as well as concerning the process of dying. Adequate symptom management may not only alleviate current suffering, but also *prevent* physical and non-physical concerns in the future.

Acknowledgements

We thank DT Sleijfer, MD, PhD for his critical comments, and the Comprehensive Cancer Centre North Netherlands (IKN), for facilitating the execution of this study.

CAM Rolf

Buorren 27, EL Tzummarum

G Pool

Northern Centre for Healthcare Research,

AV Groningen

References

- 1 Coebergh JWW. Epidemiology of cancer in the Netherlands. *Bijblijven* 2000; **16**: 7–19 (in Dutch).
- 2 McWhinney IR, Bass MJ, Donner A. Evaluation of a palliative care service. *BMJ* 1994; **309**: 1340–42.
- 3 Coyle N, Adelhardt J, Foley KM, Portenoy RK. Character of terminal illness in the advanced cancer patient: pain and other symptoms during the last four weeks of life. *J Pain Symptom Manage* 1990; **5**: 83–93.
- 4 Schuit KW, Sleijfer DT, Meijler WJ, *et al.* Symptoms and functional status of patients with disseminated cancer visiting outpatient departments. *J Pain Symptom Manage* 1998; **16**: 290–97.
- 5 Chochinov HM, Tataryn D, Clinch JJ, Dudgeon D. Will to live in the terminally ill. *Lancet* 1999; **354**: 816–19.
- 6 Chochinov HM, Hack T, Hassard T, Kristjanson LJ, McClement S, Harlos M. Dignity in the terminally ill: a cross-sectional, cohort study. *Lancet* 2002; **360**: 2026–30.
- 7 Goldsteen M, Proot I, Huijter Abu-Saad H, Widdershoven G. Good dying. Research project funded by the Ministry of Health, COPZ, University of Maastricht, 2001 (in Dutch).
- 8 Heaven CM, Maguire P. The relationship between patients' concerns and psychological distress in a hospice setting. *Psychooncology* 1998; **7**: 502–507.

Copyright of *Palliative Medicine* is the property of Arnold Publishers and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.