University of Groningen

Catalytic Oxidation of Homoallyl alcohols to α -Alkoxytetrahydrofurans by a Pd-Nitro Complex and Molecular Oxygen

Meulemans, Tom M.; Kiers, Niklaas H.; Leeuwen, Piet W.N.M. van; Feringa, Bernard

Published in:
Tetrahedron Letters

DOI:
[10.1016/0040-4039\(94\)85079-8](https://doi.org/10.1016/0040-4039(94)85079-8)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1994

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Meulemans, T. M., Kiers, N. H., Leeuwen, P. W. N. M. V., & Feringa, B. (1994). Catalytic Oxidation of Homoallyl alcohols to α -Alkoxytetrahydrofurans by a Pd-Nitro Complex and Molecular Oxygen. *Tetrahedron Letters*, 35(3). [https://doi.org/10.1016/0040-4039\(94\)85079-8](https://doi.org/10.1016/0040-4039(94)85079-8)

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

0040-4039(93)E0245-F

Catalytic Oxidation of Homoallylcohols to α -Alkoxytetrahydrofurans by a Pd-Nitro Complex and Molecular Oxygen

Tom M. Meulemans, Niklaas H. Kiers, Ben L. Feringa*
and Piet W.N.M. van Leeuwen†

Department of Organic and Molecular Inorganic Chemistry,
Groningen Center of Catalysis and Synthesis, University of Groningen,
Nijenborgh 4, 9747 AG Groningen, The Netherlands

† Koninklijke/Shell Laboratorium, Amsterdam(Shell Research B.V.), PO Box 3003, 1003 AA Amsterdam, The Netherlands

Abstract: Various homoallylcohols were highly regioselective oxidized with molecular oxygen at the terminal carbon to afford cyclic acetals (α -alkoxytetrahydrofurans) using $\text{PdNO}_2\text{Cl}(\text{CH}_3\text{CN})_2$ as a catalyst in the presence of CuCl_2 and t-butanol or isopropanol.

Catalytic oxidation of substituted olefins with molecular oxygen has attracted considerable attention¹ in order to avoid oxygen atom donors like hypochlorite, iodosylbenzene and peroxides and to have direct access to epoxides², alcohols and phenols³, aldehydes⁴ and ketones⁵. Considerable activity is devoted to mimicking oxygenases using O_2 as oxidant.^{6,7} Furthermore the indirect use of oxygen in Wacker type oxidations with PdCl_2 of olefins to ketones (scheme 1, route a) has extensively been studied.⁸ More recently palladium(II) catalyzed intramolecular cyclizations of hydroxyalkenes provide an attractive means of olefin functionalization.⁹ For instance the oxidative formation of cyclic acetals, which are good precursors for lactones¹⁰, the cyclization of a diol as a short way to both natural and unnatural Frontalin and the formation of aminoalcohols¹² illustrate nicely their potential in the synthesis of natural products. Control of the stereoselectivity might be achieved by the presence of stereogenic centers in the hydroxy alkenes.

Earlier^{4,13} we reported the use of a Pd-nitro complex capable of converting α -alkenes into aldehydes (60-70 % selectivity) or methyl ketones (> 90 % selectivity) by using molecular oxygen.⁴ Now we wish to report the effective oxidation of 1-alken-4-ols to α -alkoxytetrahydrofurans (up to 100 % selectivity and high yields) by $\text{PdNO}_2\text{Cl}(\text{CH}_3\text{CN})_2$ in the presence of CuCl_2 , O_2 in t-butanol (t-BuOH) or isopropanol (IPA), (scheme 1, path b). In addition the first selective oxidation of homoallylcohols without the necessity of directing substituents at the allylic position is described.¹⁰

Scheme 1 a. Wacker type oxidation to methylketones
b. oxidative cyclization to alkoxytetrahydrofurans

It appears that competing Wacker type oxidation to methylketones can be largely avoided.

The oxidation reactions were performed at 30°C using an in situ prepared Palladium-nitro catalyst under an O₂ atmosphere (Scheme 1, route b). The results of the oxidation reactions are given in table 1.

TABLE: 1^a

Entry	Homoallyl alcohol 4			Solvent (R ₄ OH)	React. time. (h)	Product 5 GC Yield/% (isolated Yield/%)
	R ₁	R ₂	R ₃			
1	H	H	H	IPA	3	26 ^b (≥ 70%)
2	H	H	H	t-BuOH	16	94 ^b (30)
3	H	CH ₃	CH ₃	IPA	2	100 ^b (50)
4	H	CH ₃	CH ₃	t-BuOH	2	100 ^b (80)
5	H	CH ₃	CH ₃ CH ₂	IPA	100	95 (55)
6	H	CH ₃	CH ₃ CH ₂	t-BuOH	40	100 (70)
7	H	Phenyl	Phenyl	IPA	16	80 (70)
8	H	Phenyl	Phenyl	t-BuOH	4	75 (32)
9	CH ₃	CH ₃	CH ₃	IPA	1	99 (60)
10	CH ₃	CH ₃	CH ₃	t-BuOH	2	96 (70)
11				IPA	1	56 (40)
12				t-BuOH	2.5	66 (60)

a: In a typical procedure the catalyst was made in situ by dissolving 5 mol% PdNO₂Cl(CH₃CN)₂, 20 mol% CuCl₂ in IPA or t-BuOH and stirred under oxygen at 55°C for 2 hours. After cooling the reaction mixture to 30°C, 1 g. of substrate together with an internal standard (isooctane) was added. The reaction was monitored with GC. Most reactions were completed within 4 hours. Work up was done by filtration (Al₂O₃) of the catalyst followed by extraction with pentane/water and molecular distillation under reduced pressure. All products showed IR, ¹H, ¹³C-NMR, MS data in accordance with the structures; 2-t-butoxytetrahydrofuran was prepared independently¹⁴. b: during isolation the product partly dimerizes to bisdihydrofuran **6**.

Most oxidations are highly selective at the terminal carbon. Thus the oxidation of 3-buten-1-ol in t-BuOH yields α -t-butoxytetrahydrofuran and all tertiary homoallyl alcohols are oxidized to α -alkoxytetrahydrofurans. It should be noted that, except for t-homoallyl alcohols, selectivities are higher in t-butanol as solvent. (c.f. ref. 13 for the t-butanol effect) When the carbon chain of the hydroxyalkene is lengthened by one carbon there is no selective oxidation at the terminal carbon. The formation of a six membered ring is less favourable than the formation of a five membered ring. Therefore, 2-methyl-5-hexen-2-ol and 2-allylphenol are not oxidized to α -alkoxy pyrans but only partly converted into the corresponding furans (table 1, entries 11, 12).

The catalyst is still active after one run. The activity of the catalyst was investigated in several subsequent runs using 3,4-dimethyl-1-penten-4-ol as the substrate. Typically 1 g. of substrate was added to the catalytic system (3 mol % of catalyst) and after complete conversion another gram of substrate was added and this was repeated once more. The observed conversions were 100% and 95% after the second and third run respectively showing that the catalyst is still active. However, small amounts of unidentified side products were formed after more than three runs. For comparison a number of oxidation reactions with $\text{PdCl}_2(\text{CH}_3\text{CN})_2$, CuCl_2 in t-BuOH using molecular oxygen were performed. A selective oxidation of 4-methyl-1-penten-4-ol to 5,5-dimethyl-2-t-butoxytetrahydrofuran was observed but the oxidation of 3,4-dimethyl-1-penten-4-ol was very slow and did not give complete conversion.

Nokami and co-workers¹⁰ found that a substituent on the allylic position (R_1 = alkyl, alkoxy, alkoxycarbonyl, sulfonyl) is necessary to achieve selective oxidation (scheme 1, path b) using a Wacker type system (PdCl_2 , p -benzoquinone in wet DMF). The absence of a substituent on the allylic position (**4**, R_1 =H) afforded a methylketone (scheme 1, path a). Inomata and co-workers¹⁵ found that a coordinating group is necessary to afford selective oxidation, i.e. a tosyl group was used on the allylic position (**4**, R_1 =Ts) with a PdCl_2 , CuCl_2 , N,N,N',N' -tetramethylurea, ethyl acetate based system in methanol under nitrogen to oxidize homoallyl alcohols to cyclic acetals.

Our results clearly show that a substituent at the allylic position or a "directing group" is not required for the highly selective oxidation of homoallylic alcohols to α -alkoxytetrahydrofurans with O_2 as the oxidant using a Palladium-nitro catalyst.

In a further extension the oxidative cyclization of allyl-substituted- α -hydroxyesters (**4**, $\text{R}_3=\text{CO}_2\text{Et}$) was investigated. The results of this investigation are summarized in table 2. The starting materials are readily available by allylation of α -hydroxyesters.¹⁶

TABLE 2^a

Entry	Homoallyl alcohol ^b			Solvent	React. time. (h)	Product ⁵ Yield (%)
	R^1	R^2	R^3			
1	H	CH_3	COOEt	IPA	2	60
2 ^c	H	CH_3	COOEt	t-BuOH	18	63
3 ^d	H	Phenyl	COOEt	IPA	2	70
4	H	Phenyl	COOEt	t-BuOH	20	>90

a: Typical procedure, see footnote table 1. b: racemate. c: 1 mol% of catalyst. d: 4 mol % of catalyst.

All homoallyl esters are oxidized to the α -alkoxytetrahydrofurans and only small amounts of methylketones are formed. However, the reactions in t-BuOH are significantly slower, but also more selective compared to the reactions in IPA. Ester substituted α -alkoxytetrahydrofurans are attractive precursors for substituted lactones and carbohydrate derivatives.

The oxidative cyclization by a palladium-nitro based catalyst proved to be an attractive synthetic reaction because: (1) less catalyst is required compared to related Wacker systems, (2) a very high selectivity and (3) good yields for a broad range of substrates are obtained. The availability of enzymatic methods for the preparation of optically active allylsubstituted α -hydroxyesters¹⁶ makes this catalytic oxidation reaction particularly attractive for the asymmetric synthesis of substituted tetrahydrofurans.

Acknowledgement: This work was supported by the Dutch Organization for Scientific Research (NWO) with financial support by the Stichting Technische Wetenschappen. We are grateful to Prof. R.M. Kellogg and Dr. H. Moorlag for samples of allyl-substituted α -hydroxy esters.

REFERENCES

1. Mimoun, H. *"Metal Complexes in Oxidations"* in Comprehensive Coordination Chemistry, ed. Wilkinson, G., Pergamon Press, vol.6, 317, **1987**.
2. Groves, J. T., in *"Metal ion activation of dioxygen"* ed. Spiro, T. G., Wiley, New York, **1980**.
3. Karlin, K. D.; Hayes, J. C.; Gultneh, Y.; Cruse, R. W.; Mc Kown, J. W.; Hutchinson, J. P.; Zubieta, J., *J.Am.Chem.Soc.* **1984**, 106, 2121.; Tyeklar, Z.; Karlin, K. D., *Acc.Chem.Res.* **1989**, 22, 241.; Gelling, O. J.; Feringa, B. L., *J.Am.Chem.Soc.* **1990**, 112, 7599.
4. Feringa, B. L., *J.Chem.Soc.Chem.Comm.* **1986**, 909.
5. Sheldon, R. A.; Kochi, J. K., *"Metal-catalyzed oxidations of Organic Compounds"* Academic Press, New York, **1981**.
6. Hayaishi, O. ed., *"Oxygenases"*, Academic Press, New York, **1976**.
7. Feringa, B. L., in *"Bioinorganic Chemistry of Copper"*, Karlin, K. D. ed., Chapman & Hall, **1993**, ch. 24.
8. Henry, P. M., *"Palladium catalyzed Oxidation of Hydrocarbons"*, D. Reidel, Boston, **1980**.
9. Hosokawa, T.; Miyagi, S.; Murahashi, S.; Sonoda, A., *J.Chem.Soc.Chem.Comm.* **1978**, 687.
10. Nokami, J.; Ogawa, H.; Miyamoto, S.; Mandai, T.; Wakabayashi, S.; Tsuji, J., *Tetrahedron Lett.* **1988**, 29, 5181.
11. Hosokawa, T.; Makabe, Y.; Shinohara, T.; Murahashi, S., *Chem.Lett.* **1985**, 1529. Hosokawa, T.; Nakajima, F.; Iwasa, S.; Murahashi, S., *Chem.Lett.* **1990**, 1387.
12. Benthem, R. A. T. M. van; Hiemstra, H.; Speckamp, W. M., *J.Org.Chem.* **1992**, 57, 6083.
13. Kiers, N. H.; Feringa, B. L.; Leeuwen, P. W. N. M. van, *Tetrahedron Lett.* **1992**, 2403. Kiers, N. H.; Feringa, B. L.; Kooiman, H.; Spek, A. L.; Leeuwen, P. W. N. M. van, *J.Chem.Soc.Chem.Comm.* **1992**, 1169.
14. Kruse, C. G.; Jonker, F. L.; Dert, V.; Gen, A. van de, *Recl.Trav.Chim.Pays-Bas* **1979**, 98, 371.
15. Igarashi, S.; Haruta, Y.; Ozawa, M.; Nishide, Y.; Kinoshita, H.; Inomata, K., *Chem.Lett.* **1989**, 737.
16. Moorlag, H.; Kellogg, R. M., *Tetrahedron Asymm.* **1991**, 2, 705.