University of Groningen

Synthesis of γ -Methoxy Carboxylic Acids via Nucleophilic Additions to Oxycarbenium Ions derived from 5-Methoxy-2(3H)-dihydrofuranones.

Oeveren, Arjan van; Feringa, Bernard

Published in:
Tetrahedron Letters

DOI:
[10.1016/S0040-4039\(00\)74427-3](https://doi.org/10.1016/S0040-4039(00)74427-3)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1994

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Oeveren, A. V., & Feringa, B. (1994). Synthesis of γ -Methoxy Carboxylic Acids via Nucleophilic Additions to Oxycarbenium Ions derived from 5-Methoxy-2(3H)-dihydrofuranones. *Tetrahedron Letters*, 35(45).
[https://doi.org/10.1016/S0040-4039\(00\)74427-3](https://doi.org/10.1016/S0040-4039(00)74427-3)

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

0040-4039(94)01777-8

Synthesis of γ -Methoxy Carboxylic Acids via Nucleophilic Additions to Oxycarbenium Ions derived from 5-Methoxy-2(3H)-dihydrofuranones.

Arjan van Oeveren and Ben L. Feringa*

Department of Organic and Molecular Inorganic Chemistry, Groningen Center for Catalysis and Synthesis,
University of Groningen, Nijenborgh 4, 9747 AG, Groningen, The Netherlands.

Abstract: The Lewis acid induced addition of silylated nucleophiles to 5-methoxy-2(3H)-dihydrofuranones was studied. The methodology gives access to α,β -substituted- γ -alkoxy carboxylic acids with modest to high levels of acyclic diastereocontrol in the addition reaction.

α,β -Substituted- γ -alkoxy carboxylic acids are important building blocks in natural products synthesis and as a consequence the stereoselective preparation of this type of compounds has drawn a lot of attention.^{1,2} Some derivatives have shown to be hydroxy dipeptide isosteres³ while structurally related 5-substituted γ -lactones and several 4,5-disubstituted γ -lactones are natural products by themselves.⁴

As part of our research program towards developing new methodology based on 5-alkoxy-2(3H)-dihydrofuranones **1**⁵ we have investigated the chemistry of oxycarbenium ions **2** or **3** derived from these lactones (scheme 1). The Lewis acid mediated nucleophilic additions to oxycarbenium ions derived from cyclic O,O-acetals has been extensively studied.⁶ In most cases however the acetals used so far have a general structure with both acetal oxygen atoms being part of a ring. In the case of mixed acyloxy- alkoxy acetals, treatment with a Lewis acid leads to the formation of the alkoxy substituted oxycarbenium ion due to the better leaving group ability of the acyloxy group and products derived from the acyloxy-substituted oxycarbenium ion are not found.^{7,8}

scheme 1

In the case of 5-alkoxy-2(3H)-dihydrofuranones **1** there is also a mixed acyloxy- alkoxy acetal moiety present, but in contrast only the acyloxy moiety is part of the ring. Upon treatment with strong Lewis acids 5-alkoxy-2(3H)-dihydrofuranones can form two types of oxycarbenium ions **2** or **3** as depicted in scheme 1. Quenching of these intermediates with a nucleophile might lead to two possible adducts **4** or **5** depending on the intermediate oxycarbenium ion.

We report here preliminary results of the TiCl_4 induced nucleophilic addition of silylated nucleophiles to oxycarbenium ions derived from (3,4-disubstituted)-5-methoxy-2(3H)dihydrofuranones **1**. The synthesis of substrates

1 has been described by us previously⁹ and is outlined in scheme 2. A large variety of substituents can be readily introduced at the 3 and 4 positions of **1** by tandem Michael additions-enolate alkylations or cycloadditions to 5-methoxy-2(5H)-furanone **6**.

scheme 2

The addition reactions of allyltrimethylsilane **9** and 1-trimethylsilyloxyhexene **10** to 5-methoxy-2(3H)dihydrofuranones **7** or **8** were examined (scheme 3). The oxycarbenium ions were generated in all cases by treatment with an excess of TiCl_4 as the Lewis acid. The results are summarized in table 1.¹⁰ Alkyl or hydrogen

scheme 3

Table 1: Additions of Silylated Nucleophiles to Various Substituted 5-Methoxy-2(3H)-dihydrofuranones **7** and **8**.

Entry	substrate	R ¹	R ²	nucleophile	T (°C)	product	Yield ^a %	syn:anti ^b
1	7a	H	H	9	-70	11a	74	-
2	7a	H	H	10	-70	13	59	50:50
3	7b	Me	H	9	-70	11b	87	65:35
4	7c	Et	CH ₂ Ph	9	-70	11c	86	85:15
5	8	CH ₂ C(Me)=C(Me)CH ₂	H	9	-70	12	85	c
6	7d	C(Me) ₂ CO ₂ Et	Me	9	0	11d	53	c
7	7e	CPh(CO ₂ Et) ₂	H	9	23	11e	34 ^d	<5:>95

^a Yields are of isolated pure products after chromatography

^b Syn:anti ratio determined by ¹³C NMR

^c Only one diastereomer is observed

^d >60 % starting material was recovered as a mixture of cis and trans furanone **7e**

substituted substrates (**7a,b,c,8**, entry 1,2,3,4,5) gave a very clean addition reaction at a temperature of -70°C. The substrates containing an ester group in the substituent R¹ (**7d,e**, entry 6,7) did not react at this temperature, but gave the addition reaction at more elevated temperatures, although in modest yields. This lack of reactivity might be due to stabilization of the oxycarbenium ion by the adjacent ester oxygen atom.

It should be noted that all products are derived from the intermediate oxycarbenium ion **3**, as no lactones **4**

were detected (scheme 1). The yields of these addition reactions to various alkyl substituted lactones **7a,b,c**, **8** were all consistently high, demonstrating the versatility of this method for the synthesis of γ -methoxy (alkoxy) substituted carboxylic acids.

Diastereoselectivities of the nucleophilic additions to oxycarbenium ions derived from lactones **1** depend strongly on the substituents at C₄ (see table 1). The diastereomeric ratio could be easily determined by ¹³C NMR, while the syn or anti configuration could be assigned by comparison of the ¹³C NMR data with those reported by Kunz and co-workers for **11b**.² The relative configuration of adduct **11e** was unequivocally established by a single crystal X-ray analysis.¹¹

In the case of adduct **11b** (entry 3) a mixture of diastereomers was obtained in a syn:anti ratio of 65:35. To obtain further information on the mechanism of the reaction and on the stereodirecting effect of the C₄ substituent (R¹) in **7b** both cis- and trans-substituted 5-methoxy-4-methyl-2(3H)-dihydrofuranone **7b** were tested under the same reaction conditions. The syn:anti ratio of the adducts was independent of the starting material used: cis-lactone **7b** gave the same ratio of diastereomers of the product **11b** as the trans-lactone **7b**. When 5-methoxy-4-methyl-2(3H)-dihydrofuranone **7b** was treated with TiCl₄ in CH₂Cl₂ at 0°C for 2 min. and subsequently quenched with water the product was a mixture of cis- and trans-substituted lactones **7b** in a ratio of 33:67. These results are a strong indication

scheme 4

for the direct formation of the oxycarbenium ion **3** (schemes 1 and 4), which is in equilibrium with the ring closed furanone **7**. After the initial formation of the oxycarbenium ion **3**, it reacts with the nucleophile (either intra- or intermolecular, to **7** or **11** respectively) to give a mixture of diastereomeric products, mainly determined by the size of the R¹-substituent in the substrate. The major diastereomer is the one that is predicted by employing the Felkin-Anh model.¹² The C₄-ethyl substituted substrate **7c** with a somewhat greater steric demand than the methyl substituted **7b** gives preferentially syn adduct **11c** (entry 4). Though not unambiguously determined in this case, the Felkin Ahn model predicts the syn diastereomer as the main product (scheme 4, intermediate **14**). Substrate **8** contains a rigid cyclohexyl structure. The intermediate oxycarbenium ion exists essentially in one conformation and the addition product is formed as a single diastereomer.

The substrates with an ester functional group at C₄, **7d** and **7e**, gave the intermolecular addition products **11** in a lower yield and only at higher temperatures (>0°C). Much to our surprise the nucleophilic addition took place in a completely diastereocontrolled fashion. The oxycarbenium ion derived from trans-**7e** is however formed at temperatures below -50°C. This can be concluded from the reaction of **7e** with more than 2 equivalents of TiCl₄ in

CH_2Cl_2 at -60°C . After aqueous work up a mixture of the starting material and one product was obtained. This compound was characterized by single crystal X-ray analysis¹¹ and by NMR as cis-4-substituted-5-methoxy-2(3H)-dihydrofuranone **7e**. This means that epimerization must have taken place at the acetal center. This epimerization can be explained by the formation of the oxycarbenium ion **3** as depicted in scheme 4. The oxycarbenium ion which is initially formed can undergo a ringclosure reaction to furnish a mixture of cis- and trans-substituted lactones **7e**. The ratio between cis- and trans-substituted furanone **7e** seems to be determined by the size of the substituent R^1 , but also the Lewis acid used has an effect on the equilibrium. While TiCl_4 gives cis and trans furanone **7e** in a ratio of 8:1, the use of SnCl_4 results in a cis-trans ratio of 1:17 and these reactions are reversible. This difference in ratio is not clear at present, as cis **7e** is expected from the Felkin Ahn model (scheme 4), but might be explained by the greater oxophilicity of titanium giving a strong chelate with the ester carbonyl oxygen atoms of R^1 , inhibiting the stabilization of the oxycarbenium ion by the ester carbonyl oxygen atom. The ester carbonyl oxygen atoms of the R^1 substituent are probably not blocked by SnCl_4 , allowing initial complexation to the oxycarbenium ion. Ring closure will in this case be a $\text{S}_{\text{N}}2$ type nucleophilic displacement, resulting predominantly in the trans substituted furanone **7e**.

It should be noted that in case there is a phenyl group in the substrate (entry 7), it does not act as a nucleophile to give a cyclization reaction.¹³

It can be concluded that the Lewis acid mediated addition reaction of silylated nucleophiles to oxycarbenium ions derived from 5-methoxy-2(3H)dihydrofuranones **1** is a versatile method of synthesizing a variety of γ -methoxy substituted carboxylic acids. The diastereoselectivity in the addition reactions to the forementioned oxycarbenium ions is strongly depending on the substituents at the 4-position of the substrates and both high syn and anti selectivities have been found. The stereochemical outcome of the nucleophilic addition to the oxycarbenium ions can be rationalized by the Felkin-Anh model.

Acknowledgment This work was supported by the Netherlands Foundation of Chemical Research (SON-NWO).

References and notes

1. a) Yamamoto, Y.; Chouan, Y.; Nishii, S.; Ibuka, T.; Kitahara, H. *J. Am. Chem. Soc.* **1992**, *114*, 7652. b) Hanessian S.; Sumi, K. *Synthesis* **1991**, 1083.
2. Kunz, T.; Janowitz, A.; Reißig, H-U. *Chem. Ber.* **1989**, *122*, 2165 and references cited.
3. Baker, W.R.; Pratt, J.K. *Tetrahedron* **1993**, *49*, 8739.
4. a) Takahata, H.; Uchida, Y.; Momose, T. *Tetrahedron Lett.* **1994**, *35*, 4123. b) Ebata, T.; Matsumoto, K.; Yoshikoshi, H.; Koschi, K.; Kawakami, H.; Okano, K.; Matsushita, H. *Heterocycles* **1993**, *36*, 1017. c) Nagao, Y.; Dai, W-M.; Ochiai, M.; Shiro, M. *J. Org. Chem.* **1989**, *54*, 5211. d) Ortuño, R.M.; Mercé, R.; Font, J. *Tetrahedron Lett.* **1986**, *27*, 2519.
5. For a review see: Feringa, B.L.; De Jong, J.C. *Bull. Soc. Chim. Belg.* **1992**, *101*, 627.
6. a) Johnson, W.S.; Harben, C.A.; Stipanovic, R.D. *J. Am. Chem. Soc.* **1968**, *90*, 5279. b) Denmark, S.E.; Almstead, N.G. *J. Org. Chem.* **1991**, *56*, 6458. c) Alexakis, A.; Mangeney, P. *Tetrahedron: Asymm.*, **1990**, *1*, 477.
7. a) Mashraqui, S.H.; Kellogg, R.M. *J. Org. Chem.* **1984**, *49*, 2513. b) Seebach, D.; Imwinkelried, R. *Helv. Chim. Acta* **1987**, *70*, 448.
8. For an example of type **2**, see: Jäger, V.; Kuhn, W.; Schubert, U. *Tetrahedron Lett.* **1986**, *27*, 2583.
9. a) Feringa, B.L.; De Lange, B.; Jansen, J.F.G.A.; De Jong, J.C.; Lubben, M.; Faber, W.S.; Schudde, E.P. *Pure and Appl. Chem.* **1992**, *64*, 1865. b) Van Oeveren, A.; Jansen, J.F.G.A.; Feringa, B.L. *J. Org. Chem.*, in press.
10. A typical experimental procedure is as follows: To a solution of the methoxy furanone **7** (1 mmol) and allyltrimethyl silane (3 mmol) in 5 ml CH_2Cl_2 was added at -70°C a solution of TiCl_4 in CH_2Cl_2 (1 ml, 1.5M). The reaction mixture was stirred at this temperature for 1 h and poured into a mixture of 10 ml water and 20 ml ether. The layers were separated and the water layer extracted with 10 ml ether. The combined organic layers were washed with brine (10 ml) and dried. Evaporation of the solvent gave the crude product which was purified by chromatography over silica gel.
11. Single crystal X-ray analyses were performed by Auke Meetsma from the crystallographic department of the University of Groningen; data will be published separately.
12. See: a) Eliel, E.L. *Asymmetric Synthesis*, Morrison, J.D. (Ed.), Acad. Press, New York: Vol. 2, 1983; pp. 125-155. b) Ahn, N.T. *Top. Curr. Chem.* **1980**, *88*, 145. c) Cherest, M.; Felkin, H.; Prudent, N. *Tetrahedron Lett.* **1968**, 2199.
13. There are examples known of the nucleophilic addition of a phenyl group to oxycarbenium ions, i.e. Mohler, D.L.; Thompson, D.W. *Tetrahedron Lett.* **1987**, *28*, 2567.

(Received in UK 11 July 1994; revised 5 September 1994; accepted 9 September 1994)