University of Groningen

Characterization and functional expression in Escherichia coli of the sodium/proton/glutamate symport proteins of Bacillus stearothermophilus and Bacillus caldovenax

TOLNER, B; POOLMAN, B; KONINGS, WN

Published in:
Molecular Microbiology

DOI:
[10.1111/j.1365-2958.1992.tb01464.x](https://doi.org/10.1111/j.1365-2958.1992.tb01464.x)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1992

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

TOLNER, B., POOLMAN, B., & KONINGS, WN. (1992). Characterization and functional expression in Escherichia coli of the sodium/proton/glutamate symport proteins of Bacillus stearothermophilus and Bacillus caldovenax. *Molecular Microbiology*, 6(19), 2845-2856. <https://doi.org/10.1111/j.1365-2958.1992.tb01464.x>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Characterization and functional expression in *Escherichia coli* of the sodium/proton/glutamate symport proteins of *Bacillus stearothermophilus* and *Bacillus caldotenax*

Berend Tolner, Bert Poolman and Wil N. Konings*

Department of Microbiology, University of Groningen, Kerklaan 30, NL-9751 NN Haren, The Netherlands.

Summary

The genes encoding the $\text{Na}^+/\text{H}^+/\text{L}$ -glutamate symport proteins of the thermophilic organisms *Bacillus stearothermophilus* (*gluT_{BS}*) and *Bacillus caldotenax* (*gluT_{BC}*) were cloned by complementation of *Escherichia coli* JC5412 for growth on glutamate as sole source of carbon, energy and nitrogen. The nucleotide sequences of the *gluT_{BS}* and *gluT_{BC}* genes were determined. In both cases the translated sequences corresponded with proteins of 421 amino acid residues (96.7% amino acid identity between GluT_{BS} and GluT_{BC}). Putative promoter, terminator and ribosome-binding-site sequences were found in the flanking regions. These expression signals were functional in *E. coli*. The hydropathy profiles indicate that the proteins are hydrophobic and could form 12 membrane-spanning regions. The Na^+/H^+ coupled L-glutamate symport proteins GluT_{BS} and GluT_{BC} are homologous to the strictly H^+ coupled L-glutamate transport protein of *E. coli* K-12 (overall 57.2% identity). Functional expression of glutamate transport activity was demonstrated by uptake of glutamate in whole cells and membrane vesicles. In accordance with previous observations (de Vrij *et al.*, 1989; Heyne *et al.*, 1991), glutamate uptake was driven by the electrochemical gradients of sodium ions and protons.

Introduction

To date, two types of L-glutamate transport mechanisms have been reported for thermophilic bacteria. In *Bacillus stearothermophilus* L-glutamate (or L-aspartate) transport proceeds via a sodium/proton-symport mechanism with a 1:1:1 stoichiometry (de Vrij *et al.*, 1989; 1990; Heyne *et al.*, 1991). In *Clostridium fervidus* an electrogenic sodium

symport mechanism with a stoichiometry of 2 has been identified (Speelmans *et al.*, 1989). In the mesophilic organism *Escherichia coli* three L-glutamate transport systems have been identified: (i) a binding-protein-dependent, sodium-independent, glutamate–aspartate system (inhibited by cysteate); (ii) a binding-protein-independent, sodium-independent, glutamate–aspartate system (inhibited by β -hydroxyaspartate and cysteate; and (iii) a binding-protein-independent, sodium-dependent, glutamate-specific system (inhibited by α -methylglutamate) (Halpern *et al.*, 1973; Miner and Frank, 1974; Schellenberg and Furlong, 1977). Genes encoding the sodium-motive and proton-motive transport systems, designated *gluS* and *gluP*, respectively, have been cloned (Deguchi *et al.*, 1989; Kalman *et al.*, 1991; Wallace *et al.*, 1990) and their nucleotide sequences have been reported (Deguchi *et al.*, 1990; Kalman *et al.*, 1991; Wallace *et al.*, 1990). Recently the reported sequence of the *gluP* gene of *E. coli* has been corrected (Tolner *et al.*, 1992).

The mechanism of energy coupling to glutamate transport in *B. stearothermophilus* has been described (de Vrij *et al.*, 1989; Heyne *et al.*, 1991). To elucidate the molecular properties of the sodium/proton/L-glutamate-symport transport system of *B. stearothermophilus* in more detail, a strategy was devised to clone the gene encoding the glutamate transport protein. This strategy is based on the complementation of an *E. coli* K-12 strain for growth on glutamate as sole source of energy, nitrogen and carbon. *E. coli* K-12 strains do not grow in media containing glutamate as sole source of energy, nitrogen and carbon because of an insufficient capacity to accumulate glutamate (Halpern and Lupo, 1965). Another thermophilic bacillus is *Bacillus caldotenax*, which has a higher optimum temperature of growth (70 versus 63°C of *B. stearothermophilus*), and can grow much faster than *B. stearothermophilus* on glutamate as sole source of energy, nitrogen and carbon (t_d : 30 min versus 5 h for *B. stearothermophilus*). The L-glutamate transport gene of *B. caldotenax* has also been isolated and characterized.

In this paper we report the cloning of the genes encoding the $\text{Na}^+/\text{H}^+/\text{L}$ -glutamate symport proteins of *B. stearothermophilus* (*gluT_{BS}*) and *B. caldotenax* (*gluT_{BC}*), their nucleotide sequence, deduced amino acid sequence

Fig. 1. Expression of the *gltT* genes of *B. stearotheophilus* and *B. caldodenax* in minicells of *E. coli* P678-54. Proteins were labelled in the presence of [35 S]-methionine (>1000 Ci mmol $^{-1}$) and 100 μ M IPTG, and separated on a 15% SDS/PAA gel. Lanes 1-4: P678-54 containing pUC18 (vector control), pGBT231 (*GltT_{Bc}*), pGBT102 (*GltT_{Bs}*), and pGBT112 (*GltT_{Bs}* expressed in the opposite direction from the *lac* promoter), respectively. Molecular size markers (in kDa) are indicated. Solid arrow: glutamate transport proteins. Open arrow: product of the ampicillin-resistance gene.

and deduced hydropathy profile. We conclude that *GltT_{Bs}* and *GltT_{Bc}* are homologous. Furthermore, these proteins are homologous to the H $^{+}$ /glutamate symport protein of *E. coli* K-12.

Results

Cloning of the glutamate transport genes of *B. stearotheophilus* and *B. caldodenax*

The *gltT_{Bs}* and *gltT_{Bc}* genes were cloned using the strategy outlined in the *Experimental procedures*. In the case of *gltT_{Bs}* 61 Glu $^{+}$ transformants able to grow on M9G plates (supplemented with carbenicillin and IPTG) were collected after 48 h. The cells were grown in liquid media

and their plasmid content was analysed with respect to insert size. All transformants did harbour plasmid pKK223-3, with inserts ranging from 2.5 to 7 kb in length. A total of 30 of these plasmids conferred a Glu $^{+}$ phenotype on *E. coli* JC5412 upon retransformation. One transformant harbouring pGBT38 (insert 2.5 kb) was used to perform uptake experiments in whole cells. In these cells, sodium-stimulated glutamate transport activity was significantly higher than in cells harbouring plasmid pKK223-3 (data not shown). To obtain the smallest insert that allowed JC5412 to grow on M9G, subclones of pGBT38 were constructed in pUC18. The two smallest hybrid plasmids which resulted in a Glu $^{+}$ phenotype of JC5412, were pGBT102 and pGBT112. Both plasmids contained a 1537 bp *EcoRI* fragment of pGBT38 but in opposite orientation. Since the *gltT_{Bs}* gene in pGBT112 is expressed in the opposite direction of the *lac* promoter, the *gltT_{Bs}* promoter may have been cloned along with the *gltT_{Bs}* gene.

The *gltT_{Bc}* gene was cloned essentially as described above for the *gltT_{Bs}* gene, and was located on a 1535 bp *EcoRI* fragment (pGBT231). The *gltT_{Bc}* gene could also be expressed independently of its orientation relative to the *lac* promoter of pUC18.

Expression of the glutamate transport genes of *B. stearotheophilus* and *B. caldodenax*

In the minicell-producing strain *E. coli* P678-54, in which pGBT102 and pGBT112 were used to express *GltT_{Bs}*, one additional protein band with an apparent molecular mass of 33 kDa was found which was not present in a control strain containing pUC18 (Fig. 1).

Uptake of L-glutamate and L-aspartate by whole cells (strain *E. coli* JC5412) harbouring pGBT102 (*GltT_{Bs}*) was several-fold higher than in cells harbouring pUC18 (Fig. 2). The initial rate of uptake and steady-state level of

Fig. 2. Sodium-ion dependent uptake of L-glutamate (A) and L-aspartate (B) by cells expressing the glutamate transport protein of *B. stearotheophilus*. Uptake by cells harbouring plasmid pUC18 (vector control; $\square$, $\blacksquare$) and pGBT102 (*GltT_{Bs}*; $\circ$, $\bullet$) were compared. Concentrated cells were diluted to a final concentration of 0.75 mg protein per ml into 50 mM potassium phosphate, pH 6.0, 5 mM MgSO $_4$, and 10 mM glucose buffer, with ($\bullet$, $\blacksquare$) or without ($\circ$, $\square$) 20 mM NaCl. After 1 min of incubation, L-[14 C]-glutamate (1.75 μ M) or L-[14 C]-aspartate (2.23 μ M) was added and transport assays were further handled as described in the *Experimental procedures*.

Fig. 3. L-glutamate counterflow activity by membrane vesicles of *E. coli* (JC5412(pUC18 or pGBT102)). Counterflow by membrane vesicles prepared from *E. coli* harbouring either pUC18 (vector control; □) or pGBT102 (GltT_{BS}; O). Membrane vesicles loaded with 1 mM L-glutamate were diluted 100-fold into 50 mM potassium phosphate, pH 6.0, 5 mM MgSO₄ and 3.5 μM L-[¹⁴C]-glutamate. The transport reaction was stopped at different time intervals as indicated in the *Experimental procedures*.

accumulation of L-glutamate increased significantly upon the addition of 20 mM NaCl (Fig. 2). Similar observations were made for (sodium)-proton motive force driven L-glutamate uptake in membrane vesicles derived from strain JC5412 harbouring pGBT102 (data not shown). Membrane vesicles derived from strain JC5412 harbouring pGBT102 also showed significantly higher L-glutamate counterflow activity than membrane vesicles derived from strain JC5412 harbouring pUC18 (Fig. 3).

The kinetic parameters (apparent K_m and V_{max}) of L-glutamate uptake in membrane vesicles of *B. caldovenax* and membrane vesicles of *E. coli* JC5412 expressing GltT_{BS} or GltT_{BC} were determined (Table 1). Uptake of

glutamate mediated by GltT_{BS} and GltT_{BC} yielded lower V_{max} and K_m values when the proteins were assayed in membrane vesicles of *E. coli* relative to membrane vesicles of *B. stearothermophilus* and *B. caldovenax*.

Nucleotide sequence and coding regions of the *EcoRI* fragments of pGBT102 and pGBT231

The sequencing strategies for the *B. stearothermophilus* and *B. caldovenax* glutamate transport genes are presented in Fig. 4, A and B, respectively. The sequence of the 1537 bp *EcoRI* fragment of *B. stearothermophilus* in pGBT102 is shown in Fig. 5A. Between positions 110 and 1373 an open reading frame (ORF) of 1263 bp is found. The deduced polypeptide contains 421 amino acid residues, corresponding with a molecular mass of 45 469 Da. The nucleotide sequence of the 1535 bp *EcoRI* insert

Fig. 4. Sequencing strategy for the *EcoRI* fragments containing the *gltT_{BS}* gene of *B. stearothermophilus* (pGBT102) (A) and the *gltT_{BC}* gene of *B. caldovenax* (pGBT231) (B). Part of the vector sequence (open box), the cloned fragment (shaded box) and the position and direction of transcription of the *gltT_{BS/BC}* (large arrow above the sequence) are shown. The regions sequenced are indicated by black arrows. Symbols: E, Hd, R, S and Hc represent *EcoRI*, *HindIII*, *RsaI*, *Sau3A* and *HincII* restriction endonuclease, respectively.

Table 1. Apparent kinetic parameters for glutamate transport^a by membrane vesicles of *B. stearothermophilus*, *B. caldovenax*, *E. coli* JC5412(pGBT102) and JC5412(pGBT231).

Vesicles derived from:	K_m^{app} (μM)	V_{max} (nmol mg protein ⁻¹ min ⁻¹)
<i>B. stearothermophilus</i> (GltT _{BS})	4.7 ^b	11.4 ^b
<i>B. caldovenax</i> (GltT _{BC})	2.9	17.4
<i>E. coli</i> JC5412(pGBT102) (GltT _{BS})	31.8	4.8
<i>E. coli</i> JC5412(pGBT231) (GltT _{BC})	25.1	6.2

a. Uptake experiments were performed by diluting membrane vesicles 100-fold in 50 mM potassium phosphate, pH 6.0, 5 mM MgSO₄, 40 mM glucose and 500 μM NaCl. After 3 min of preincubation PQQ 920 μM was added. After another min of incubation L-[¹⁴C]-glutamate (1.75 μM) was added and transport assays were further handled as described in the *Experimental procedures*.

b. Data taken from Heyne *et al.* (1991).

A ⁻¹⁰
 GAATTCAATGACAATATATGAAATATTCCTGACATAACGAAATATGATAGTAGAATA
 -35
 GTTTGTCTCTTTTGTCTATAAAATGTCATAAAAAGAAAGGGCGGAATGATGAGAGAAAAT
 START
 M R K I
 TGGATTAGCTTGGCAAAATTTTATTTGGTCTCATTTCTAGGATTTATCGTGGAGCCATTTT
 G L A W Q I F I G L I L G I I V G A I F
 TTACGAAATCCGAAGGTGGCACATTATTACAGCCTATTGGAGATATTTTCTTCGTTT
 Y G N P K V A T Y L Q P I G D I F L R L
 AATCAAAATGATGTCATTCGATTTGTTATTTCTAGCCTGTAGTTGGAGTCGCCACGCT
 I K M I V I P I V I S L V V G V A S V
 TGGGATTGGAAGCTTGGAAAATTAGCGGGCAAAACGATTTATTTCTCGAGATTAT
 G D L K K L G K L G G K T I I Y F E I I
 CACAACGATCGGATTGTCGCGTTTATTGGCAGCGAATATTTTTCAGCCAGGACCGG
 T T I A I V V G L L A A N I F Q P G T G
 CGTTAATATGAATCATAGAAAACCGATATTCAAAGCTATGTTGATACAAACGA
 V N M K S L E K T D I Q S Y V D T T N E
 AGTCAGCATCATTCGATGGGAAACTTTTGTAAATATGTTCCAAAAAATATTTTGA
 V Q H H S M V E T F V N I V P K N I F E
 ATCGTTAACCAAGGGATATGCTCGCATCATTTCTCTCTCTTATGTTTCGGTTTAGG
 S L T K G D M L P I I F F S V M F G L G
 AGTAGCGGATTTGGCGAAAAGGAGCCAGTTCTTCAATTTTCCAAAGGTACAGCAGA
 V A A I G E K G K P V L Q F F Q G T A E
 AGCGATGTTTATGTAACAAACCAATTAAGAAGTTTGGCCCGTTTCGGCGTTTTCGGT
 A M F Y V T N Q I M K F A P F G V F A L
 GATTGGTGTAACCGTTTCTAAGTTTGGGGTAGAGTCGCTTATTCGCTCAGCAAGCTCGT
 I G V T V S K F G V E S L I P L S K L V

60 CATGTGTTTACGCCAACGATGGTGTTCCTTTATCTTTGTCTGCTGGCGGTGTGCTAA 840
 I V V Y A T M V F F I F V V L G G V A K
 900 GTTATTGGCATAAAATATTTTTCATATATATAAAATTTTGAAGATGAGTTAAATCTTGC
 L F G I N I F H I I K I L K D E L I L A
 960 TTATAGTACAGCAAGTTTCGGAACCGTCTTCCGAAAAATATGAGAAAAATGGAGAATTT
 Y S T A S S E T V L P K I M E K M E N E
 1020 CGGTGTCGCAAAAGCGATTACATCCTTTGTCTCATCCGACAGGGTATCTTTTAACTAGA
 G C P K A I T S F V I P T G Y S F N L D
 1080 CGGTCTACGTTATATCAGGCGTTGGCGCCATTTTATCGCGCAGTTGTACGGTATTGA
 G S T L Y Q A L A A I F I A Q L Y G I D
 1140 CATGCCGATTTCTCAACAAATCTCGCTTTTGTCTTGTGTTAAATGTTGACTTCGAAAGGAAT
 M P I S O Q I S L L L V L M V T S K G I
 1200 CGCTGGGTGCGGGTGTATCCTTTCTCGTCTGCTGTACGTTAGGCACGGTTGGGAT
 A G V P G V S F V V L L A T L G T V G I
 1260 TCCGATAGAAGGATTAGCATTTATCGCTGGAATCGACCGTATTTTAGATATGCGCGCAC
 P I E G L A F I A G I D R I L D M A R T
 1320 AGCAGTGAATGTTATTTGGCACTCGTTAGCAGCGATCATTTATGTCAAAAATGGGAAGCCA
 A V N V I G N S L A A I I M S K W E G Q
 1380 ATATAACGAAGAAAAGGAAAAACAATACATCGCGCAATTCAGCAAAAGTGCATAAATGCTT
 Y N E E K G K Q Y I A Q L Q Q S A - --
 1440 GAAAAAGCTGCTCTAGAAAAGGACAGCTTTTTCAGTATAAGGACTTATTTTTCCTT
 -----> <-----
 1500 CCACGTAGTATAGTATAATTGTTGCTGAATAAGTTGTAAATGAGATGCTTTGGCTAAGAG
 1537 TACATAATACATAGGAGGAAGAACCATGCACGAATTC

B	GAATCAATGACAATATTATATGAATATCTTTGACATAACGAAAAATATAATAGTAGAATA -35 -10 START GTTTGTCTCTTTTGTCTAAAAATGCTGTAATAAGGGGGCGCAATCAATGAGAAAAATC M R K I G L A W O I F I G L I L G I I V G A I F GGGTAGCTTGGCAAAATTTTCATGCTGCTCATTTTGGGATATCGTGGGAGCCATTTT TATGGGAATCCGAAGTTGCCGCTATTACAGCCGATGGAGATATTTCTCCGTTA Y G N P K V A A Y L Q P I G D I F L R L ATAAAAATGATGCTCAATCCGATGTTATTTCTAGCCTGTGCTGAGTCGCCAGCGTT I K M I V I P I V I S S L V V G V A S V GGGGATTTGAAAAAGCTTGGAAAAATAGCGGCAAAACGATTTATTTTTCAGATTATC G D L K K L G K L G G K T I I Y F E I I ACAAGCATCGGATGCTCGCTGCTTTATTTGGCGGCAATATTTTTCAGCCAGGGCGGC T T I A I V V G L L A A N I F Q P G A G GTTAATATGAATCATAGAAAAACGGATATTCAAAGCTATGTTGATACAAACAGAA V N M K S L E K T D I Q S Y V D T T N E GTGCAGCATCATCGATGGTAGAAACATTTGCAATATGTTCCGAAAAACATTTTGAA V Q H H S M V E T F V N I V P K N I F E TCGTTATCCACAGGAGACATGCTCCGATCATTTTCTCTGTTATGTTCCGTTTAGGA S L S T G D M L P I I F F S V M F G L G GTAGCGCGATGGCGAAAAAGGAGCCAGTCTTCAATTTTCCAAAGGCACACAGAA V A A I G E K G K P V L Q F F Q G T A E GCAATGTTTATGTAACAAACCAATATGAAGTTTGGCGGCTTCCGTTGTTGCGCTT A M F Y V T N Q I M K F A P F G V F A L ATCGGTGTAAACGTTTAAATTTGGAGTAGAGTCGCTTATTTCCGTCAGCAAGCTCGTC I G V T V S K F G V E S L I P L S K L V	ATTGTTGTTTACGCAACGATGCTGTTCTTTATCTTTGCGAGTCTTGGCGGTGTTGCTAAG I V V Y A T M L F F I F A V L G G V A K 60 TTATTTGGTATAAATATTTTTCATATTTATAAAAAATTTTGAAGAATCAATTAATCTTGCT L F G I N I F H I I K I L K D E L I L A 900 TATAGTACAGCAAGTTCCGAAACCGTTCTTCCAAAGAAATCAATGATAAATGGAATTTT Y S T A S S E T V L P R I M D K M E K F 960 GGTCGCCAAAAGCGATTACATCTTTGTCTATCCCGACAGGTATTCCTTTAACTTAGAC G C P K A I T S F V I P T G Y S F N L D 1020 GGATCTACGTTATATACGGCGTTGGTCCCATTTTATCGCGCAGTTGTACGGTATGAC G S T L Y Q A L A A I F I A Q L Y G I D 1080 ATGTCGGTTTCTCAACAAATTTTCGCTTTTGTCTGTCTTAATGGTGAATTCGAAGGAATC M S V S Q Q I S L L L V L M V T S K G I 1140 GCTGGGTGCCAGGTGATCCTTTGCTGCTGTCTGTCTACATTAGGCACGGTTGGGATT A G V P G V S F V V L L A T L G T V G I 1200 CCGGTAGAAGGATTAGCATTTATCGGGGATCGACCGCATTTTATAGATATGGCAGCACA P V E G L A F I A G I D R I L D M A R T 1260 GCAGTCAATGTTATTCGCAACTCATTAGCAGCGATCATTTATGTCAAAATGGGAAGGCCAA A V N V I G N S L A A I I M S K W E G Q 1320 TATAACGAAGAAAAAGAAAAACAATATCTTTCAGAGTTGCAGCAAAAGTCATAATGATCA Y N E E K G K Q Y L A E L Q Q S A - STOP 1380 AAAAGTGTCCCGCTAGAGAGACGGCTTTTTCAGTGTAAAAAATTTATTTTCTCTGC -----> <----- 1440 AACGTAGTATAGTATAAATTTTCTGCTGAATAAGTTGTAACGGGATGCTTGGCTAAATA 1500 CATAAATGCATAGGAGAGACAAATGCACGAATTC 1535
----------	---	---

Fig. 5. Nucleotide sequence of the *EcoRI* fragments containing the *gluT_{bs}* gene of *B. stearothermophilus* (pGBT102) (A) and the *gluT_{bs}* gene of *B. caldolenax* (pGBT231) (B). The start and stop codons, putative promoter(-35/-10), possible ribosome-binding site (RBS) and possible terminator sequences (→←) are indicated. The amino acid sequence deduced from the DNA sequences of the *gluT_{bs}* and *gluT_{bs}* genes are shown below the DNA sequence. The 12 possible membrane-spanning regions are underlined and in bold face. These sequence data appear in the EMBL/GenBank/DBJ Nucleotide Sequence Data Libraries under the accession numbers M86508 (*gluT_{bs}*) and M86509 (*gluT_{bs}*).

Protein	% Amino Acid Identity (Similarity) With:			
	GlIT _{BS}	GlIT _{BC}	GlTP _{Eck-12}	GlTS _{Eck-12/EcB}
GlIT _{BS}	100			
GlIT _{BC}	96.7 (2.1)	100		
GlTP _{Eck-12}	60.1 (17.3)	60.3 (16.9)	100	
GlTS _{Eck-12/EcB}	8.5 (6.0)	9.5 (6.5)	12.5 (8.5)	100

		# S	
GltT _{Bs}	MRKI--GLAWQIFIGLILGIIVGAI--FYGNPK---VATYLPQIGDIFLR	43	
GltT _{Bc}	MRKI--GLAWQIFIGLILGIIVGAI--FYGNPK---VAAYLPQIGDIFLR	43	
GltP _{Ec}	MKNIKFSLAWQILFAMVLGILLGSYLHYHSDSRDWLVNLLSPAGDIFIH	50	
	.. .*****.....*..* *..* * ..*..*		
	S S SS		
GltT _{Bs}	LIKMIVIPIVISSLVVGASVGDLLKGLGKGTIIYFEIITITIAIVVGL	93	
GltT _{Bc}	LIKMIVIPIVISSLVVGASVGDLLKGLGKGTIIYFEIITITIAIVVGL	93	
GltP _{Ec}	LIKMIIVPIVISTLVVGIAVGDAKQLGRIGAKTIIYFEVITTVAILGI	100	
	*****.*****.*****.*.*** *..*..* .*****.*****.*****.		
	#		
GltT _{Bs}	LAANIFQPGTGVNMKSLEKTDIQSYVDTTNEVQH--HSMVETFFVNIVPKN	141	
GltT _{Bc}	LAANIFQPGAGVNMKSLEKTDIQSYVDTTNEVQH--HSMVETFFVNIVPKN	141	
GltP _{Ec}	TLANVFQPGAGVDMSQLATVDISKYQSTTEAVQSSSHGIMGTILSLVPTN	150	
	..***.*..* *..*..* *..*..*..*..*..*		
	##		
GltT _{Bs}	IFESLTKGDMLPPIIFFSVMFGLGVAAGKEGK-PVLQFFQGTAEAMFYVT	190	
GltT _{Bc}	IFESLSTGDMLPPIIFFSVMFGLGVAAGKEGK-PVLQFFQGTAEAMFYVT	190	
GltP _{Ec}	IVASMAKGEMLPIIFFSVLFGLGLSSLPATHREPLVTVFRSISSETMFKVT	200	
	* *****.*****. *..* *..*..*		
	# #		
GltT _{Bs}	NQIMKFAFPGVFALIGVTVSKFGVESLIPLSKLVIVVYATMVFFIFVVLG	240	
GltT _{Bc}	NQIMKFAFPGVFALIGVTVSKFGVESLIPLSKLVIVVYATMLFFIFAVLG	240	
GltP _{Ec}	HMVMRYAPVGVFALIAVTVANFGFSSLWPLAKLVLLVHFALFFALVVLG	250	
	 ....* *****.*****. *..*..*..*..*..*		
	# # #		
GltT _{Bs}	GVAKLFGINIFHIIKILKDELILAYSTASSETVLPKIMEKMENFGCPKAI	290	
GltT _{Bc}	GVAKLFGINIFHIIKILKDELILAYSTASSETVLPKIMEKMENFGCPKAI	290	
GltP _{Ec}	IVARLGLSVWILIRILKDELILAYSTASSESVLPRIIEKMEAYGAPVSI	300	
	.. *..*..* *****.*****.*****.*..*..*..*..*		
	##		
GltT _{Bs}	TSFVIPTGYSFNLGSLTYQALAAIFIAQLYGIDMPISQQISLLLVLMVT	340	
GltT _{Bc}	TSFVIPTGYSFNLGSLTYQALAAIFIAQLYGIDMSVSQQISLLLVLMVT	340	
GltP _{Ec}	TSFVVPTGYSFNLGSLTYQSIAAIFIAQLYGIDLSIWQEIIILVLTLMVT	350	
	****.*****.*****.*****.*****. *..* *..*..*..*		
	#		
GltT _{Bs}	SKGIAGVPGVSFVLLATLGTGVIPIEGLAFIAGIDRILDMARTAVNVIG	390	
GltT _{Bc}	SKGIAGVPGVSFVLLATLGTGVIPIEGLAFIAGIDRILDMARTAVNVIG	390	
GltP _{Ec}	SKGIAGVPGVSFVLLATLGSVGIPLEGLAFIAGVDRILDMARTALNVVG	400	
	*****.*****.*****.*****.*****.*****.*****.*****.		
	# #		
GltT _{Bs}	NSLAAIIMSKWEGQYNEEKQKQY----IAQLQQA--	421	
GltT _{Bc}	NSLAAIIMSKWEGQYNEEKQKQY----LAELQQA--	421	
GltP _{Ec}	NALAVLVIAKWEHKFDRKALAYEREVLGKFDKTADQ	437	
	....*..* *..* *..*..*..*..*		

Fig. 6. Multiple alignment of the deduced amino acid sequences of the GltT_{Bs}, GltT_{Bc} and GltP_{EcK-12} glutamate transport proteins. The best fit was achieved by introducing gaps in order to maximize the identity score. The overall identity was 57.2%. Identical residues and conserved substitutions are indicated by asterisks and full points, respectively. Symbols: S, residues involved in the putative Na⁺ recognition or binding motif; #, mismatches between the GltT_{Bs} and GltT_{Bc} proteins. GltP_{EcK-12} sequence was taken from Tolner *et al.* (1992).

B. stearothermophilus (de Vrij *et al.*, 1989; Heyne *et al.*, 1991) and therefore suggest that the gene encoding the previously described sodium/proton/glutamate symport protein of *B. stearothermophilus* (de Vrij *et al.*, 1989; Heyne *et al.*, 1991) has been cloned.

The GltT_{Bs} and GltT_{Bc} proteins both consist of 421 amino acid residues, corresponding with molecular masses of 45 469 and 45 345 Da, respectively. These values are higher than the apparent molecular masses of

33 000 Da estimated from SDS-polyacrylamide gel electrophoresis (SDS-PAGE) (Fig. 1). However, aberrant electrophoretic behaviour is often observed for integral membrane proteins (e.g. Büchel *et al.*, 1980; Deguchi *et al.*, 1990; Ehrling *et al.*, 1980; Nakao *et al.*, 1987; Poolman *et al.*, 1989; Van der Rest *et al.*, 1990; Wallace *et al.*, 1990; Yazyu *et al.*, 1984), and is probably explained by increased binding of sodium dodecyl sulphate due to the hydrophobic nature of the proteins.

Fig. 7. Hydropathy profiles of the GltT_{Bs} (A), GltT_{Bc} (B) and GltP_{Eck-12} (C) glutamate transport proteins. The hydropathy was calculated according to the method of Eisenberg *et al.* (1984) with a window of 21 amino acids. The positions of the 12 putative membrane-spanning segments are indicated by solid bars. GltP_{Eck-12} sequence was taken from Tolner *et al.* (1992).

Transcription of *gltT_{Bs}* is probably initiated from the promoter indicated in Fig. 5A. The -35 and -10 promoter regions, and the spacing of 17 bp between the two regions, is similar to that of consensus promoter sequences (-35: TTGACA; -10: TATAAT; spacing 16–18, 17–19 bp) that are recognized by the σ^{43} and σ^{70} factors of the holoenzyme form of RNA polymerase of *B. subtilis* (Helmann and Chamberlin, 1988; Moran *et al.*, 1982) and *E. coli* (Hawley and McClure, 1983; Helmann and Chamberlin, 1988), respectively. This suggests the

existence of a sigma factor in *B. stearothermophilus*, which is similar to σ^{43} in *B. subtilis* and σ^{70} in *E. coli*, that are involved in transcription of genes for housekeeping functions. A putative ribosome-binding site (RBS) is located at proper distance (4 bp) from the translation initiation codon (Fig. 5A). The RBS shows extensive complementarity to the 3' end of *B. stearothermophilus* 16S rRNA (Douthwaite *et al.*, 1983). The stop codon (TAA at position 1373–1375) is followed by an inverted repeat (ΔG° -114.2 kJ mol⁻¹, calculated according to Tinoco *et al.*, 1973) with features of a putative rho-independent transcription terminator sequence (Rosenberg and Court, 1979). The -35 and -10 promoter regions of *gltT_{Bc}* are identical to those of *gltT_{Bs}*. The putative ribosomal binding site (Fig. 5B), however, shows major differences and is probably much weaker than the one upstream of *gltT_{Bs}* of *B. stearothermophilus*. Also, the transcription terminator sequence is much weaker when compared with the one in the *B. stearothermophilus* glutamate transport gene (ΔG° -75.6 kJ mol⁻¹, calculated according to Tinoco *et al.*, 1973). The differences in the expression signals of *gltT_{Bs}*

Table 4. Codon usage of the glutamate transport gene of *B. stearothermophilus* (Bs) and *B. caldovenax* (Bc).

Codon	Amino Acid	No. of times used		Codon	Amino Acid	No. of times used	
		Bs	(Bc)			Bs	(Bc)
TTT	Phe	19	(20)	TAT	Tyr	8	(10)
TTC	Phe	9	(8)	TAC	Tyr	4	(2)
TTA	Leu	17	(17)	TAA	---	1	(1)
TTG	Leu	7	(8)	TAG	---	0	(0)
CTT	Leu	11	(12)	CAT	His	3	(3)
CTC	Leu	3	(4)	CAC	His	0	(0)
CTA	Leu	1	(0)	CAA	Gln	11	(10)
CTG	Leu	3	(3)	CAG	Gln	6	(6)
ATT	Ile	40	(36)	AAT	Asn	8	(6)
ATC	Ile	13	(14)	AAC	Asn	5	(6)
ATA	Ile	3	(3)	AAA	Lys	18	(18)
ATG	MET	15	(15)	AAG	Lys	8	(7)
GTT	Val	17	(16)	GAT	Asp	7	(7)
GTC	Val	8	(9)	GAC	Asp	3	(4)
GTA	Val	6	(7)	GAA	Glu	11	(13)
GTG	Val	11	(10)	GAG	Glu	5	(3)
TCT	Ser	6	(5)	TGT	Cys	1	(0)
TCC	Ser	2	(5)	TGC	Cys	0	(1)
TCA	Ser	2	(3)	TGA	---	0	(0)
TCG	Ser	7	(6)	TGG	Trp	2	(2)
CCT	Pro	1	(0)	CGT	Arg	2	(1)
CCC	Pro	0	(0)	CGC	Arg	1	(2)
CCA	Pro	4	(5)	CGA	Arg	0	(0)
CCG	Pro	10	(9)	CGG	Arg	0	(0)
ACT	Thr	3	(1)	AGT	Ser	3	(3)
ACC	Thr	5	(2)	AGC	Ser	4	(4)
ACA	Thr	9	(12)	AGA	Arg	1	(2)
ACG	Thr	6	(6)	AGG	Arg	0	(0)
GCT	Ala	6	(7)	GGT	Gly	10	(10)
GCC	Ala	4	(5)	GGC	Gly	10	(10)
GCA	Ala	8	(11)	GGA	Gly	12	(11)
GCG	Ala	14	(12)	GGG	Gly	8	(9)

and $gltT_{Bc}$ are not reflected in the expression levels of the proteins both in *E. coli* and the thermophilic bacilli (Table 1). Since the expression levels of $GltT$ in *B. stearothermophilus* and *B. caldodenax* are similar, the observed differences in growth rates on glutamate as sole carbon, energy and nitrogen source cannot be explained at the level of transport. The presence of putative promoter and transcription termination sequences flanking the glutamate transport genes of *B. stearothermophilus* and *B. caldodenax* suggests that both genes are transcribed as single cistronic messages.

The apparent K_m and V_{max} values for L-glutamate transport in membrane vesicles derived from *B. stearothermophilus* and *B. caldodenax* are very similar (Table 1). However, *B. caldodenax* can grow approximately 10-fold faster than *B. stearothermophilus* in media with 50 mM glutamate as sole source of energy, carbon and nitrogen (data not shown). Therefore it is unlikely that the $GltT_{Bs}$ transport protein is limiting the growth of *B. stearothermophilus* on glutamate as sole source of energy, carbon and nitrogen.

The deduced amino acid sequence of the Na^+/H^+ /glutamate symport proteins of *B. stearothermophilus* and *B. caldodenax* were initially compared with the H^+ /glutamate symport protein of *E. coli* K-12 (Wallace *et al.*, 1990). This did reveal regions of homology while other regions differed completely. By translating the nucleotide sequence of $gltP_{EcK-12}$ in different reading frames, and by comparing the translated sequences with those of $GltT_{Bs}$ and $GltT_{Bc}$ it became apparent that the sequence divergence between $GltP_{EcK-12}$ and $GltT_{Bs}$ and $GltT_{Bc}$ was probably caused by sequencing errors, i.e. base substitutions, deletions and insertions, in the L-glutamate transport gene of *E. coli*. The $gltP_{EcK-12}$ sequence, as published by Wallace *et al.* (1990), was therefore resequenced and revised (Tolner *et al.*, 1992). Sequence comparisons revealed extensive similarity between the $GltT_{Bs}$ and $GltT_{Bc}$ and the revised $GltP_{EcK-12}$ sequence (Fig. 6 and Table 3).

A conserved amino acid sequence has been proposed to be involved in Na^+ recognition or binding (SOB-motif $\text{---G---A---L---GR---}$) (Deguchi *et al.*, 1990). This sequence was also found in the $GltT_{Bs}$ and $GltT_{Bc}$ proteins, except that Arg was replaced for Lys ($\text{---G}_{38}\text{---A}_{62}\text{---L}_{67}\text{---G}_{71}\text{K}_{72}\text{---}$; see also Fig. 6). However, apart from one mismatch, the SOB motif can also be found in the $GltP_{Ec}$ protein ($\text{---G}_{45}\text{---A}_{69}\text{---A}_{74}\text{---G}_{78}\text{R}_{79}\text{---}$; see also Fig. 6). If indeed this SOB motif is essential for Na^+ -binding this mismatch could explain the inability of $GltP_{Ec}$ to use Na^+ as coupling ion despite the extensive similarity between $GltT_{Bs}$ and $GltP_{Ec}$. On the other hand, the SOB motif of $GltT_{Bs}$ and $GltT_{Bc}$ is located in a short hydrophilic region and might be involved in retention of the three-dimensional shape of these proteins, as is proposed for

this region in the 'consensus glucose transport protein' (Henderson, 1990).

Experimental procedures

Bacterial strains, plasmids and growth conditions

The bacterial strains, plasmids and phages used in this study are listed in Table 5. *B. stearothermophilus* and *B. caldodenax* were grown at 63 and 70°C, respectively, with vigorous aeration in a medium containing 2% (w/v) tryptone, 1% (w/v) yeast extract and 170 mM NaCl, and adjusted to pH 7.0. For growth experiments, mineral medium of pH 7.0 was used, containing 1 ml of trace element solution (Vishniac and Santer, 1957) per litre of medium, 34 mM Na_2HPO_4 , 22 mM KH_2PO_4 , 10 mM NaCl, 1 mM $MgSO_4$, 0.1 mM $CaCl_2$, and 50 mM L-glutamate as sole source of energy, nitrogen and carbon. *E. coli* strains were grown at 37°C with vigorous aeration in LB, M9, M9G (M9 in which ammonium-chloride was replaced by L-glutamate at a final concentration of 10 mM) or M9CA medium (Sambrook *et al.*, 1989). The mineral media were supplemented with essential nutrients as indicated by the auxotrophic markers. When needed, carbenicillin and IPTG were added to a final concentration of 100 $\mu\text{g ml}^{-1}$ and 100 μM , respectively.

DNA manipulations

Mini- and large-scale preparations of plasmid DNA were obtained by the alkaline lysis method (Birnboim and Doly, 1979; Ish-Horowicz and Burke, 1981). Chromosomal DNA was isolated essentially as described previously (Leenhouts *et al.*, 1990), except that mutanolysine was omitted. The strains were transformed by the rubidium chloride (Sambrook *et al.*, 1989) or by the electrotransformation (Dower *et al.*, 1988) method. Other DNA techniques were performed as described previously (Sambrook *et al.*, 1989).

Cloning of the glutamate transport gene

The strategy for cloning the $gltT_{Bs}$ and $gltT_{Bc}$ genes is based on the complementation of *E. coli* K-12 strain JC5412, which does not grow on glutamate as sole source of energy, nitrogen and carbon. Partially *EcoRI*-, *HindIII*-, *PstI*- or *Sau3A*-digested chromosomal DNA of *B. stearothermophilus* or *B. caldodenax* was fractionated by polyacrylamide gel (6% w/v) electrophoresis. Fragments of 2 to 20 kb were electroeluted from the gel and ligated into linearized and dephosphorylated pKK223-3. The resulting hybrid plasmids were used to transform *E. coli* JC5412 by electrotransformation. Transformants able to grow on M9G plates (supplemented with carbenicillin and IPTG) were analysed with respect to their plasmid content. Purified plasmids were used to retransform *E. coli* JC5412 in order to distinguish between Glu^+ revertants and true transformants. Transformants again were selected on M9G plates.

Sequence determination of the glutamate transport genes

The nucleotide sequences of both strands of the *EcoRI* fragment of pGBT102 and pGBT231, or subclones derived thereof

Table 5. Bacterial strains, plasmids and phages used.

Bacterial strain, plasmid or phage	Relevant characteristics	Source/Reference
Bacterium		
<i>B. stearotheophilus</i>		ATCC7954
<i>B. caldopenax</i>		Heinen and Heinen (1972)
<i>E. coli</i>		
JM101	$\Delta(lac-proAB)$ (F^+ $lacI^q$ $\Delta M15$)	Yanisch-Perron <i>et al.</i> (1985)
JC5412	doesn't grow on L-glutamate as sole carbon, nitrogen and energy source	Willets and Clark (1969)
P678-54	Minicell-producing	Adler <i>et al.</i> (1967)
Plasmid		
pUC18	Ap ^R	Yanisch-Perron <i>et al.</i> (1985)
pKK223-3	Ap ^R , expression vector	Pharmacia
pGBT38	pKK223-3, carrying <i>gltT</i> of <i>B. stearotheophilus</i> on a 2500 bp <i>EcoRI</i> – <i>EcoRI</i> fragment	This work
pGBT102	pUC18, carrying <i>gltT</i> of <i>B. stearotheophilus</i> on a 1537 bp <i>EcoRI</i> – <i>EcoRI</i> fragment	This work
pGBT112	pUC18, carrying <i>gltT</i> of <i>B. stearotheophilus</i> on a 1537 bp <i>EcoRI</i> – <i>EcoRI</i> fragment (in reverse orientation relative to pGBT102)	This work
pGBT231	pUC18, carrying <i>gltT</i> of <i>B. caldopenax</i> on a 1535 bp <i>EcoRI</i> – <i>EcoRI</i> fragment	This work
Phage		
M13mp18/19		Yanisch-Perron <i>et al.</i> (1985)
Ap ^R , ampicillin-resistant.		

in pUC18 or M13mp18/19 (*RsaI*, *Sau3A*, *HpaI*, *HindIII* or *HincII* fragments), were determined by using the dideoxy-chain termination method (Sanger *et al.*, 1977). A T7 sequencing kit (Pharmacia) was used in sequencing either single- or double-stranded DNA. MICROGENIE (Release 5.0, Beckman, Palo Alto, Cal., USA) and PCGENE (release 6.26, Genofit) were used for computer-assisted sequence analysis.

Transport assays with whole cells

Cells (15 ml) of strain JC5412 harbouring plasmid pUC18, pGBT102, or pGBT231, grown for 10 h in LB (supplemented with carbenicillin and IPTG), were harvested, washed three times in 50 mM potassium phosphate, pH 6.0, and 5 mM MgSO₄ and resuspended to a final A₆₆₀ of approximately 300 in the same buffer. Uptake of L-[¹⁴C]-glutamate or L-[¹⁴C]-aspartate was assayed at 37°C, upon 100-fold dilution of the cells into buffer consisting of 50 mM potassium phosphate, pH 6.0, 5 mM MgSO₄, and 10 mM glucose, with or without the addition of 20 mM NaCl or 20 mM KCl (see legends for details). This mixture was incubated for 1 min at 37°C under continuous aeration. To initiate the uptake experiment L-[¹⁴C]-glutamate or L-[¹⁴C]-aspartate was added to a final concentration of 1.75 or 2.23 μ M, respectively. The uptake reactions were terminated by adding a 20-fold excess of ice-cold 0.1 M potassium chloride, followed by immediate filtration over cellulose nitrate filters (0.45 μ m, pore size). The filters were washed once with 2 ml of ice-cold potassium chloride.

Transport assays with membrane vesicles

For transport studies in membrane vesicles, cells of strain JC5412 harbouring plasmid pUC18, pGBT102, or pGBT231, were grown to an A₆₆₀ of 0.7 in LB (supplemented with carbenicillin and IPTG). Cells were harvested and membrane vesicles were isolated as described previously (Kaback, 1971). Membranes were finally resuspended to 15 mg protein per ml in 50 mM potassium phosphate, pH 6.0, and stored in liquid nitrogen.

Modes of transport

Counterflow activity. Membrane vesicles were washed twice with 50 mM potassium phosphate, pH 6.0, 5 mM MgSO₄ and resuspended in the same buffer supplemented with 1 mM L-glutamate. After 2 h of incubation at room temperature, membrane vesicles were pelleted by centrifugation and resuspended to 20 mg protein per ml in the same buffer. Counterflow was initiated by diluting membrane vesicles 100-fold with buffer consisting of 50 mM potassium phosphate, pH 6.0, 5 mM MgSO₄ and 3.5 μ M L-[¹⁴C]-glutamate. The reaction was terminated as described for whole cells.

Sodium/proton motive force driven uptake. The electron donor system 2,7,9, tricarboxy-1-H-pyrrolo-(2,3)-quinoline-4,5-dione (PQQ)/glucose was used to generate a Δp (van Schie *et al.*, 1985). Membrane vesicles were diluted 100-fold with buffer consisting of 50 mM potassium phosphate, pH 6.0, 5 mM

MgSO₄, and 40 mM glucose. After 3 min of preincubation the electron mediator PQQ was added to a final concentration of 20 µM. To initiate the uptake experiment, L-[¹⁴C]-glutamate or L-[¹⁴C]-aspartate was added after another minute of incubation to a final concentration of 1.75 and 2.23 µM, respectively. Further handling was the same as described for whole cells. All transport assays were carried out at 37°C.

Determination of kinetic parameters

The kinetic parameters for transport, apparent K_m and V_{max} , were estimated from the uptake of labelled amino acid in the first 10 s. Results were analysed by Eadie-Hofstee Plots.

Minicells

Minicells of strain P678-54 were purified in three subsequent sucrose gradient centrifugations (Maeger *et al.*, 1977). The *in vivo* labelled ([³⁵S]-methionine) proteins were resolved by 15% (w/v) (SDS-PAGE) and visualized by autoradiography.

Protein determination

Protein was determined by the method of Lowry *et al.* (1951) using bovine serum albumin as standard.

Nomenclature

In order to discriminate between the proton/glutamate and the sodium/glutamate transport proteins of *E. coli* the gene designations *gltP* and *gltS* are used. The L-glutamate transport systems of *B. stearothermophilus* and *B. caldolenax* translocate glutamate in symport with sodium ions and protons. For the gene encoding these proteins the designation *gltT* was used. To discriminate between the genes and proteins the subscripts Bs, Bc and Ec (B or K-12) were added, for *B. stearothermophilus*, *B. caldolenax* and *E. coli* (B or K-12), respectively.

References

- Adler, H.I., Fischer, W.D., Cohen, A., and Hardigree, A.A. (1967) Miniature *Escherichia coli* cells deficient in DNA. *Proc Natl Acad Sci USA* **57**: 321–326.
- Birnboim, H.C., and Doly, J. (1979) A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucl Acids Res* **7**: 1513–1523.
- Büchel, D.E., Gronenborn, B., and Müller-Hill, B. (1980) Sequence of the lactose permease gene. *Nature* **283**: 541–545.
- Deguchi, Y., Yamato, I., and Anraku, Y. (1989) Molecular cloning of *gltS* and *gltP*, which encode glutamate carriers of *Escherichia coli* B. *J Bacteriol* **171**: 1314–1319.
- Deguchi, Y., Yamato, I., and Anraku, Y. (1990) Nucleotide sequence of *gltS*, the Na⁺/glutamate symport carrier gene of *Escherichia coli* B. *J Biol Chem* **265**: 21704–21708.
- de Vrij, W., Bulthuis, R.A., van Iwaarden, P.R., and Konings, W.N. (1989) Mechanism of L-glutamate transport in membrane vesicles from *Bacillus stearothermophilus*. *J Bacteriol* **171**: 1118–1125.
- de Vrij, W., Speelmans, G., Heyne, R.I.R., and Konings, W.N. (1990) Energy transduction and amino acid transport in thermophilic aerobic and fermentative bacteria. *FEMS Microbiol Revs* **75**: 183–200.
- Douthwaite S., Christensen, A., and Garrett, R.A. (1983) Higher order structure in the 3'-minor domain of small sub-unit ribosomal RNAs from a gram negative bacterium, a gram positive bacterium and a eukaryote. *J Mol Biol* **169**: 249–279.
- Dower W.J., Miller, J.F., and Ragsdale, C.W. (1988) High efficiency transformation of *E. coli* by high voltage electroporation. *Nucl Acids Res* **16**: 6127–6145.
- Ehring, R., Beyreuther, K., Wright, J.K., and Overath, P. (1980) *In vitro* and *in vivo* products of *E. coli* lactose permease gene are identical. *Nature* **283**: 537–540.
- Eisenberg, D., Schwarz, E., Komarony, M., and Wall, R. (1984) Analysis of membrane and surface protein sequences with the hydrophobic moment plot. *J Mol Biol* **179**: 125–142.
- Halpern, Y.S., and Lupo, M. (1965) Glutamate transport in wild-type and mutant strains of *Escherichia coli*. *J Bacteriol* **90**: 1288–1295.
- Halpern, Y.S., Barash, H., Dover, S., and Druck, K. (1973) Sodium and potassium requirements for active transport of glutamate by *Escherichia coli* K12. *J Bacteriol* **114**: 53–58.
- Hawley, D.K., and McClure, W.R. (1983) Compilation and analysis of *Escherichia coli* promoter DNA sequences. *Nucl Acids Res* **1**: 2237–55.
- Heinen, U.J., and Heinen, W. (1972) Characteristics and properties of a caldo-active bacterium producing extracellular enzymes and two related strains. *Arch Mikrobiol* **82**: 1–23.
- Helmann, J.D., and Chamberlin, M.J. (1988) Structure and function of bacterial sigma factors. *Annu Rev Biochem* **57**: 839–872.
- Henderson, P.J.F. (1990) The homologous glucose transport proteins of prokaryotes and eukaryotes. *Res Microbiol* **141**: 316–328.
- Heyne, R.I.R., de Vrij, W., Crielard, W., and Konings, W.N. (1991) Sodium ion dependent amino acid transport in membrane vesicles of *Bacillus stearothermophilus*. *J Bacteriol* **173**: 791–800.
- Ish-Horowicz, D., and Burke, F.J. (1981) Rapid and efficient cosmid cloning. *Nucl Acids Res* **9**: 2989–2999.
- Kaback, H.R. (1971) Bacterial membranes. *Meth Enzymol* **22**: 99–120.
- Kalman, M., Gentry, D.R., and Cashel, M. (1991) Characterization of the *Escherichia coli* K12 *gltS* glutamate permease gene. *Mol Gen Genet* **225**: 379–386.
- Leenhouts, K.J., Kok, J., and Venema, G. (1990) Stability of integrated plasmids in the chromosome of *Lactococcus lactis*. *Appl Environ Microbiol* **56**: 2726–2735.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L., and Randall, R.J. (1951) Protein measurement with the Folin phenol reagent. *J Biol Chem* **193**: 265–275.
- Maeger, R.B., Tait, R.C., Betlach, M.C., and Boyer, H.W. (1977) Protein expression in *Escherichia coli* minicells by recombinant plasmids. *Cell* **10**: 521–536.
- Miner, K.M., and Frank, L. (1974) Sodium-stimulated glutamate transport in osmotically shocked cells and membrane vesicles of *Escherichia coli*. *J Bacteriol* **117**: 1093–1098.
- Moran, C.P., Lang, N., LeGrice, S.F.J., Lee, G., Stephens, M., Sonenshein, A.L., Pero, J., and Losick, R. (1982) Nucleotide sequences that signal the initiation of transcription and translation in *Bacillus subtilis*. *Mol Gen Genet* **186**: 339–346.
- Nakao, T., Yamato, I., and Anraku, Y. (1987) Nucleotide sequence of *putP*, the proline carrier gene of *Escherichia coli* K12. *Mol Gen Genet* **208**: 70–75.

- Poolman, B., Royer, T.J., Mainzer, S.E., and Schmidt, B.F. (1989) Lactose transport system of *Streptococcus thermophilus*: a hybrid protein with homology to the melibiose carrier and enzyme III of phosphoenolpyruvate-dependent phosphotransferase systems. *J Bacteriol* **171**: 244–253.
- Rosenberg, M., and Court, D. (1979) Regulatory sequences involved in the promotion and termination of RNA transcription. *Annu Rev Genet* **13**: 319–53.
- Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*. 2nd ed. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press.
- Sanger, F., Nicklen, S., and Coulson, A.R. (1977) DNA sequencing with chain terminating inhibitors. *Proc Natl Acad Sci USA* **74**: 5463–5467.
- Schellenberg, G.D., and Furlong, C.E. (1977) Resolution of the multiplicity of the glutamate and aspartate transport systems of *Escherichia coli*. *J Biol Chem* **252**: 9055–9064.
- Speelmans G., de Vrij, W. and Konings, W.N. (1989) Characterization of amino acid transport in membrane vesicles from the thermophilic fermentative bacterium *Clostridium fervidus*. *J Bacteriol* **171**: 3788–3795.
- Tinoco, I., Borer, P.N., Dengler, B., Levine, M.K., Uhlenbeck, O.C., Crothers D.M., and Gralla, J. (1973) Improved estimation of secondary structure in ribonucleic acids. *Nature* **246**: 40–41.
- Tolner, B., Poolman, B., Wallace, B., and Konings, W.N. (1992) Revised nucleotide sequence of the *glpP* gene, which encodes the proton-glutamate-aspartate transport protein of *Escherichia coli* K-12. *J Bacteriol* **174**: 2391–2393.
- Van der Rest, M.E., Schwarz, E., Oesterheld, D., and Konings, W.N. (1990) DNA Sequence of a citrate carrier of *Klebsiella pneumoniae*. *Eur J Biochem* **189**: 401–407.
- van Schie, B.J., Hellingwerf, K.J., van Dijken, J.P., Elferink, M.G.L., van Dijk, J.M., Kuenen, J.G., and Konings, W.N. (1985) Energy transduction by electron transfer via a pyrrolo-quinoline quinone-dependent glucose dehydrogenase in *Escherichia coli*, *Pseudomonas aeruginosa*, and *Acinetobacter calcoaceticus* (var. *lwoffii*). *J Bacteriol* **163**: 493–499.
- Vishniac, W., and Santer, M. (1957) The thiobacilli. *Bact Rev* **21**: 195–213.
- Wallace, B., Yang, Y., Hong, J., and Lum, D. (1990) Cloning and sequencing of a gene encoding a glutamate and aspartate carrier of *Escherichia coli* K12. *J Bacteriol* **172**: 3214–3220.
- Willetts, N.S., and Clark, A.J. (1969) Characteristics of some multiply recombination-deficient strains of *Escherichia coli*. *J Bacteriol* **100**: 231–239.
- Yanisch-Perron, C., Vieira, J., and Messing, J. (1985) Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13mp18 and pUC19 vectors. *Gene* **33**: 103–119.
- Yazyu, H., Niiya, S.S., Shimamoto, T., Kanazawa, H., Futai, M., and Tsuchiya, T. (1984) Nucleotide sequence of the *melB* gene and characteristics of deduced amino acid sequence of the melibiose carrier in *Escherichia coli*. *J Biol Chem* **259**: 4320–4326.