University of Groningen

Lithium, an inhibitor of cAMP-induced inositol 1,4,5-trisphosphate accumulation in Dictyostelium discoideum, inhibits activation of guanine-nucleotide-binding regulatory proteins, reduces activation of adenylylcyclase, but potentiates activation of guanyl

Peters, Dorien J.M.; Snaar-Jagalska, B. Ewa; Haastert, Peter J.M. van; Schaap, Pauline

Published in:
European Journal of Biochemistry

DOI:
[10.1111/j.1432-1033.1992.tb17289.x](https://doi.org/10.1111/j.1432-1033.1992.tb17289.x)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1992

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Peters, D. J. M., Snaar-Jagalska, B. E., Haastert, P. J. M. V., & Schaap, P. (1992). Lithium, an inhibitor of cAMP-induced inositol 1,4,5-trisphosphate accumulation in Dictyostelium discoideum, inhibits activation of guanine-nucleotide-binding regulatory proteins, reduces activation of adenylylcyclase, but potentiates activation of guanyl. *European Journal of Biochemistry*, 209(1), 299-304. <https://doi.org/10.1111/j.1432-1033.1992.tb17289.x>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Lithium, an inhibitor of cAMP-induced inositol 1,4,5-trisphosphate accumulation in *Dictyostelium discoideum*, inhibits activation of guanine-nucleotide-binding regulatory proteins, reduces activation of adenylylcyclase, but potentiates activation of guanylyl cyclase by cAMP

Dorien J. M. PETERS¹, B. Ewa SNAAR-JAGALSKA¹, Peter J. M. VAN HAASTERT² and Pauline SCHAAP¹

¹ Cell Biology and Genetics, Department of Biology, University of Leiden, The Netherlands

² Department of Biochemistry, University of Groningen, The Netherlands

(Received June 10, 1992) — EJB 92 0815

Li^+ drastically alters pattern formation in *Dictyostelium* by inhibiting cAMP-induced prespore-gene expression and promoting cAMP-induced prestalk-gene expression. We reported previously that Li^+ inhibits inositol monophosphatases in this organism and strongly reduces basal and cAMP-stimulated inositol 1,4,5-trisphosphate levels. We show here that Li^+ also reduces cAMP-induced accumulation of cAMP, but promotes cAMP-induced accumulation of cGMP. This effect is not due to inhibition of cGMP hydrolysis or inhibition of adaptation and may therefore reflect stimulation of guanylyl-cyclase activation. Li^+ does not affect the binding of cAMP to surface receptors but interferes with the interaction between receptors and guanine-nucleotide-binding regulatory (G) proteins. These effects are complex; in the absence of Mg^{2+} , Li^+ increases guanosine 5'-[γ -thio]triphosphate (GTP[S])-binding activity to similar levels as 1 mM Mg^{2+} . However, while Mg^{2+} potentiates cAMP-induced stimulation of GTP[S]-binding activity, Li^+ effectively inhibits stimulation. Li^+ also inhibits cAMP-stimulated, but not basal high-affinity GTP-ase activity, indicating an inhibitory effect on cAMP-induced activation of G-proteins.

Our data suggest that in addition to inositolphosphate metabolism, the activation of G-proteins may be a second biochemical target for Li^+ effects on pattern formation and signal transduction in *Dictyostelium*.

Lithium strongly affects processes as diverse as human behaviour and embryonic development, but its effects are nevertheless quite selective. Lithium can control manic depressive psychosis at serum levels around 1 mM, without major peripheral side effects. During amphibian development, lithium acts primarily at the early cleavage stages and dorsalizes the response of animal cells to mesoderm-inducing signals (Backstrom, 1954; Breckenridge et al., 1987; Kao and Elinson, 1989).

Both therapeutic and teratogenic effects of Li^+ may result from inhibition by Li^+ of inositol monophosphatases

(Hallcher and Sherman, 1980; Delvaux et al., 1987), which impairs inositol synthesis, resulting in reduced levels of inositolphospholipids available for inositol 1,4,5-trisphosphate [$\text{Ins}(1,4,5)\text{P}_3$]-mediated signalling (Berridge et al., 1989). The therapeutic effects of Li^+ have also been attributed to direct or guanine-nucleotide-binding-regulatory-protein-(G-protein)-mediated inhibitory effects of Li^+ on adenylylcyclase activation (Ebstein et al., 1980; Mørk and Geissler, 1987; Avissar et al., 1988) and to Li^+ induced potentiation of neuropeptide-gene expression (Dobner et al., 1988).

Li^+ -induced inositol depletion accounts for the selectivity of the effects of Li^+ , since it will specifically affect those systems, such as the brain and the early embryo, which do not have ready access to inositol from the blood circulation (Berridge et al., 1989). This hypothesis is supported by observations that inositol levels as well as agonist-induced formation of $\text{Ins}(1,4,5)\text{P}_3$ and inositol 1,3,4,5-tetrakisphosphate [$\text{Ins}(1,3,4,5)\text{P}_4$] are reduced in the brains of Li^+ -treated rats (Allison and Stewart, 1971; Batty and Nahorski, 1985; Whitworth and Kendall, 1988; Godfrey et al., 1989; Casebolt and Jope, 1989); it was furthermore shown that inositol can abolish the teratogenic effects of Li^+ in amphibians (Busa and Gimlich, 1989). However, recent data showed that guanine-

Correspondence to P. Schaap, Department of Biology, University of Leiden, Kaiserstraat 63, NL-2311 GP Leiden, The Netherlands

Abbreviations. GTP[S], guanosine 5'-[γ -thio]triphosphate; ATP[S], adenosine 5'-[γ -thio]triphosphate; App[NH]p, adenosine 5'-[β , γ -imino]triphosphate; cAMP[S], adenosine 3',5'-[thio]phosphate, Sp-isomer; 2'-deoxy-cAMP, 2'-deoxy-adenosine 3',5'-monophosphate; $\text{Ins}(1,4,5)\text{P}_3$, inositol 1,4,5-trisphosphate; $\text{Ins}(1,3,4,5)\text{P}_4$, inositol 1,3,4,5-tetrakisphosphate; G protein, guanine-nucleotide-binding regulatory protein; G_s protein, guanine-nucleotide-binding regulatory protein activating adenylylcyclase.

Enzymes. Adenylylcyclase (EC 4.6.1.1); guanylyl cyclase (EC 4.6.1.2); phospholipase C (EC 3.1.4.3); cGMP phosphodiesterase (EC 3.1.4.35); creatine kinase (EC 2.7.3.2).

nucleotide-binding regulatory proteins activating adenylyl cyclase (G_s proteins) in brain are more sensitive to inhibition by Li^+ than their homologues in heart, which may provide an alternative explanation for the specific sensitivity of the brain to Li^+ (Schreiber et al., 1990).

During development of the primitive eukaryote *Dictyostelium discoideum*, Li^+ perturbs pattern formation by inhibiting expression of prespore genes and promoting expression of a subclass of prestalk genes (Maeda, 1970; Van Lookeren Campagne et al., 1988a; Peters et al., 1989). Both classes of genes are regulated by extracellular cAMP (Kay, 1982; Mehdy et al., 1983; Wang et al., 1988). cAMP-signal processing is initiated by the binding of cAMP to cell surface receptors. This interaction induces G-protein-mediated activation of adenylyl cyclase and phospholipase C, as well as the activation of guanylyl cyclase (Mato and Malchow, 1978; Theibert and Devreotes, 1986; Europe-Finner and Newell, 1987; Van Haastert et al., 1987, 1989). Li^+ ions inhibit inositol monophosphatases in *Dictyostelium* (Van Lookeren Campagne et al., 1988b) and decrease basal and cAMP-stimulated $Ins(1,4,5)P_3$ levels (Peters et al., 1989). It was suggested that effects of Li^+ on pattern formation and inositol phosphate signalling were in *Dictyostelium* also due to inositol depletion (Peters et al., 1989; Berridge et al., 1989).

We here report the effects of Li^+ on the other aspects of cAMP-signal processing. The most striking result is a pronounced inhibitory effect of Li^+ on cAMP-induced activation of G-proteins, suggesting a second target for Li^+ effects on gene expression and signal transduction.

MATERIALS AND METHODS

Materials

Creatine phosphate, creatine kinase, GTP[S], adenosine 5'-[β , γ -imino]triphosphate (App[NH]p), adenosine 5'-[γ -thio]triphosphate (ATP[S]) and adenosine 3',5'-[thio]phosphate, Sp-isomer (cAMP[S]) were obtained from Boehringer Mannheim. [2,8- 3H]cAMP, [^{35}S]GTP[S], cGMP-radioimmunoassay kits and $Ins(1,4,5)P_3$ assay kits were from Amersham, [γ - ^{32}P]GTP was from New England Nuclear, and 2'-deoxyadenosine 3',5'-monophosphate (2'-deoxy-cAMP) and dithiothreitol were obtained from Sigma.

Culture conditions

D. discoideum strain NC4 and *stm* F mutant NP368 (Ross and Newell, 1981) were grown in association with *Escherichia coli* 281 on glucose/peptone agar. Cells were freed from bacteria by repeated washing with 10 mM Na/K phosphate (phosphate buffer), pH 6.5, and subsequently resuspended in this buffer and shaken at 150 rpm at 22°C, or distributed on phosphate-buffered agar and incubated at 22°C.

Assays for [^{35}S]GTP[S] binding, GTPase and [2,8- 3H]cAMP-binding activity in cell membranes

Preparation of membranes

Cells starved for 5 h on phosphate-buffered agar were resuspended in ice-cold lysis buffer (0.5 mM EDTA, 250 mM sucrose, 40 mM Hepes, pH 7.7) to a density of 10^8 cells/ml and forced through a nuclepore filter (pore size, 3 μ m). Lysates were centrifuged at $10000 \times g$ for 5 min; pellets were washed and resuspended to a density of 2×10^8 cells/ml in

either phosphate buffer (GTP[S] and cAMP binding) or 50 mM triethanolamine/HCl, pH 7.4 (GTPase activity).

[^{35}S]GTP[S] binding

Binding of [^{35}S]GTP[S] was measured in a total volume of 100 μ l phosphate buffer containing 50 μ l membranes, 0.1 nM [^{35}S]GTP[S], 1 mM ATP and variable additions of cAMP, LiCl and $MgCl_2$. After 30 min incubation at 0°C, samples were centrifuged for 3 min at $10000 \times g$ and 4°C. The supernatant was aspirated and the radioactivity of the pellet was determined. Assay controls were obtained by including 0.1 mM GTP in the incubation mixture (Snaar-Jagalska et al., 1988a).

[2,8- 3H]cAMP binding

Binding of [2,8- 3H]cAMP to membranes was measured at 20°C in a total volume of 100 μ l, containing 50 μ l membranes, 2 nM [2,8- 3H]cAMP and variable additions of GTP[S] and LiCl. After 75 s, membranes were centrifuged through silicon oil and the radioactivity of the pellet was measured. Control values were obtained by including 1 mM cAMP in the incubation mixture.

GTPase activity

The GTPase reaction mixture, containing 37 nM [γ - ^{32}P]GTP (0.1 μ Ci/assay), 2 mM $MgCl_2$, 0.1 mM EGTA, 0.2 mM App[NH]p, 0.1 mM ATP[S], 10 mM dithiothreitol, 5 mM creatine phosphate, 0.4 mg/ml creatine kinase and 2 mg/ml bovine serum albumin in 50 mM triethanolamine/HCl, pH 7.4, was incubated for 5 min at 25°C. The reaction was started by adding 30 μ l membranes to 70 μ l reaction mixture and terminated after 3 min by adding 0.5 ml 5% (mass/vol.) activated charcoal in 50 mM sodium phosphate, pH 2.0. Samples were centrifuged for 5 min at $10000 \times g$ and 4°C and the amount of $^{32}PO_4^{3-}$ in the supernatant was determined. Low-affinity GTPase was determined in the presence of 100 μ M GTP and high-affinity GTPase in the presence of 0.01 μ M GTP. Stimulation of high-affinity GTPase was measured by adding 100 μ M cAMP to the reaction mixture (Snaar-Jagalska et al., 1988b).

Assays for cAMP and cGMP accumulation

To measure cAMP relay, 27- μ l aliquots of suspension of 4×10^7 cells/ml were stimulated with 3 μ l 50 μ M 2'-deoxy-cAMP, 50 mM dithiothreitol. After 0 min or 3 min of incubation at 20°C, the reaction was terminated by addition of 30 μ l 3.5% perchloric acid. cAMP levels in the neutralized extracts were measured by means of a cAMP isotope-dilution assay.

To measure cAMP-stimulated cGMP levels 27- μ l aliquots of 8×10^7 cells/ml were stimulated with 3 μ l 1 μ M cAMP or 2 μ M cAMP[S], at 20°C. The reaction was terminated by adding 30 μ l 3.5% perchloric acid. cGMP levels in the neutralized extracts were measured by means of a cGMP radioimmunoassay.

RESULTS

Effects of Li^+ on cAMP-receptor/G-protein interactions

cAMP-signal transduction is initiated by binding of cAMP to cell surface receptors and activation of one or more G

Fig. 1. Effects of Li⁺ on Mg²⁺ and cAMP stimulation of GTP[S] binding. [³⁵S]GTP[S] binding to membranes of aggregation-competent *Dictyostelium* cells was measured in the presence of 0, 1 or 10 mM MgCl₂ and the indicated concentrations of LiCl. (—) No cAMP; (+) plus 100 μM cAMP. Results are expressed as percentage of binding measured in the absence of cAMP, Li⁺ or Mg²⁺ (0.06 pmol/mg protein). Values are means ± SEM, *n* = 3.

proteins. Receptor mediated activation of G proteins results in increased GTP binding and increased GTPase activity (Cassel and Selinger, 1976, 1978; Snaar-Jagalska et al., 1988a, b). Activated G proteins, however, reduce the affinity of agonist receptors (Ross et al., 1977; Van Haastert, 1984). Mg²⁺ increase GTP[S] binding activity, by causing dissociation of GDP from the G protein, which makes the GTP-binding site on the α subunit accessible to GTP, or its slowly hydrolysed derivative GTP[S] (Higashima et al., 1987).

We first measured effects of LiCl on cAMP-activated and Mg²⁺-activated [³⁵S]-GTP[S] binding to membranes of aggregation-competent *Dictyostelium* cells. Fig. 1 shows that 1 mM and 10 mM Mg²⁺ cause twofold and fourfold increases of [³⁵S]GTP[S] binding, respectively. cAMP stimulation of GTP[S] binding is low in the absence of Mg²⁺ and optimal at 1 mM Mg²⁺. At 10 mM Mg²⁺, further stimulation by cAMP becomes less pronounced. Remarkably, Li⁺ (2–10 mM) induces an increase in [³⁵S]GTP[S] binding comparable to that induced by 1 mM Mg²⁺. However, in contrast to 1 mM Mg²⁺, which potentiates cAMP stimulation of [³⁵S]GTP[S] binding, Li⁺ almost completely inhibits stimulation by cAMP.

Fig. 2 shows the effect of Li⁺ on cAMP-stimulated GTPase activity. *Dictyostelium* membranes exhibit both high-affinity and low-affinity GTPase activity. cAMP stimulates only high-affinity GTPase, by increasing the affinity for GTP (Snaar-Jagalska et al., 1988b). Basal low-affinity and high-affinity GTPase activities were not affected by Li⁺. However, 10 mM Li⁺ completely inhibited cAMP stimulation of high-affinity GTPase activity; half-maximal inhibition was achieved by about 1 mM Li⁺. No effects of other monovalent cations such as Na⁺ and K⁺ on GTP[S] binding or GTPase activity were observed (data not shown).

Fig. 3 shows the effect of LiCl on [2,8-³H]cAMP-binding to membranes. LiCl concentrations up to 10 mM did not significantly affect [2,8-³H]cAMP-binding activity. GTP[S] induced an 80% reduction of [2,8-³H]cAMP binding, which was also not altered by Li⁺.

Effects of Li⁺ on activation of second messenger systems

We showed previously that Li⁺ inhibits cAMP-induced production of the intracellular messenger Ins(1,4,5)P₃ (Peters

Fig. 2. Effects of Li⁺ on GTPase activity. Total, high-affinity and low-affinity GTPase activities were measured in membranes in the presence of 2 mM Mg²⁺ and the indicated concentrations of Li⁺. Total GTPase represents the sum of high-affinity and low-affinity GTPase. To measure low-affinity (aff.) or high-affinity GTPase, substrate concentrations were 100 μM or 0.01 μM GTP, respectively. High-affinity GTPase was determined in the presence or absence of 100 μM cAMP. Results are expressed as percentage of total GTPase activity measured in the absence of Li⁺ (3.84 pmol · min⁻¹ · mg protein⁻¹). Values are means ± SEM, *n* = 3.

Fig. 3. Effects of Li⁺ and GTP[S] on [³H]cAMP binding. The binding of 2 nM [³H]cAMP binding to membranes of aggregation-competent cells was measured in the presence or absence of 100 μM GTP[S] and the indicated LiCl concentrations. Data are expressed as percentage of [³H]cAMP binding in the absence of GTP[S] and LiCl. Values are means ± SEM, *n* = 3.

et al., 1989). The second messengers cAMP and cGMP are also formed upon cAMP stimulation. Fig. 4 shows the effects of LiCl on cAMP accumulation induced by the receptor agonist 2'-deoxy-cAMP. Saturating agonist concentrations induce a 10-fold increase of basal cAMP levels, which reach a maximum at about 4 min. LiCl causes a 50% reduction of this increase. A half-maximal reduction was obtained by about 2 mM LiCl (Fig. 4B). No effects of LiCl on basal cAMP levels were observed.

Fig. 5 shows the effects of Li⁺ on the cAMP-induced cGMP response. In this case, Li⁺ appears to potentiate cAMP-stimulated accumulation of cGMP. Basal cGMP levels are not significantly affected by LiCl. Agonist-induced accumulation of cGMP is a complex response which consists of activation of guanylyl cyclase, rapid degradation of cGMP by

Fig. 4. Effects of Li^+ on agonist-induced cAMP accumulation. Aggregation-competent cells were resuspended in phosphate buffer at 4×10^7 cells/ml, incubated for 15 min with LiCl and stimulated with $5 \mu\text{M}$ 2'-deoxy-cAMP and 5 mM dithiothreitol. (A) Time course of agonist-induced cAMP accumulation in the presence of (○) 0, (△) 3 or (□) 10 mM LiCl. (B) Dose dependence of the effect of LiCl on cAMP accumulation, measured after 3 min of stimulation with agonist. Data are expressed as percentage of cAMP levels accumulated after 3 min of stimulation in the absence of LiCl (31 ± 12 pmol/ 10^7 cells). Values are means $\pm$ SEM, $n = 3$.

a cGMP-stimulated cGMP phosphodiesterase (Bulgakov and Van Haastert, 1982), sensory adaptation and inhibition of guanylyl-cyclase activity (Van Haastert and van der Heijden, 1983).

We analysed the level at which Li^+ interferes with the cGMP response. Possible effects on cGMP degradation were determined in *stim* F mutant NP368, which lacks cGMP phosphodiesterase (Ross and Newell, 1981). Fig. 6A shows that instead of being reduced, the stimulatory effect of LiCl on the cGMP response is much more pronounced in the *stim* F mutant than in wild-type cells, indicating that the effect of Li^+ is not due to inhibition of cGMP phosphodiesterase. The comparatively small effect of Li^+ on the cGMP response in wild-type cells is probably due to the fact that cGMP is very rapidly degraded.

Li^+ could also increase the cGMP response by inhibiting sensory adaptation. To test this possibility, cells were stimulated with the slowly hydrolysable cAMP-derivative cAMP[S], at a non-saturating concentration. After 30 s, cells were restimulated with a saturating stimulus of 100 nM cAMP. Fig. 6B shows that the second cGMP response to 100 nM cAMP is considerably reduced, compared to the response to 100 nM cAMP which is not preceded by a non-saturating stimulus. The difference in responsiveness to a first or second stimulus is due to adaptation of the cells. Both first and second responses are higher in the presence of 2 mM or 10 mM LiCl, but when background cGMP levels ($t = 30$ s) are subtracted, the relative decrease is about the same in the presence or absence of LiCl. This indicates that LiCl does not increase cGMP levels by inhibiting adaptation and we conclude that Li^+ stimulates activation of guanylyl cyclase.

DISCUSSION

Pronounced effects of Li^+ were observed on cAMP-induced activation of GTP-binding proteins. Li^+ inhibits cAMP

Fig. 5. Effects of Li^+ on cAMP-induced cGMP accumulation. Aggregation-competent NC-4 cells were incubated for 15 min with different LiCl concentrations. (A) cGMP accumulation in cells treated with 0 (○) or 10 mM (□) LiCl measured at different time intervals after stimulation with 100 nM cAMP. (B) Dose dependence of the effect of LiCl on cGMP accumulation measured at 0 s (○) and 10 s (●) after stimulation with 100 nM cAMP. Data are expressed as percentage of the amount of cGMP accumulated after 10 s of cAMP stimulation in the absence of LiCl (14 ± 4 pmol/ 10^7 cells). Values are means $\pm$ SEM, $n = 3-5$.

stimulation of GTP[S]-binding activity and high affinity GTPase activity. The effects of Li^+ on GTP[S] binding are complex; basal and cAMP-stimulated GTP[S]-binding activity increase in the presence of Mg^{2+} . Li^+ induces a moderate increase in GTP[S]-binding activity, comparable to that of 1 mM Mg^{2+} , but completely prevents stimulation by cAMP.

Avissar et al. (1988, 1991) reported that in rat cortex, Li^+ inhibits stimulation of GTP binding by both cholinergic and adrenergic agonists. In their system, Li^+ also inhibits the guanine-5'-[β , γ -imino]triphosphate-induced reduction in receptor affinity, but does not inhibit agonist stimulated GTPase activity. This is rather unexpected, since both GTP binding and GTP hydrolysis are determined by Mg^{2+} -induced and agonist-induced GDP dissociation (Gilman, 1987). The effects of Li^+ in rat cortex can be abolished by 1–5 mM Mg^{2+} , which suggested that Li^+ competes with Mg^{2+} for binding to the low-affinity Mg^{2+} site, which controls GDP/GTP exchange (Higashima et al., 1987).

In *Dictyostelium*, Li^+ inhibits agonist stimulation of GTP[S] binding as well as GTPase activity, but does not affect receptor affinity. The effects of Li^+ on agonist-stimulated GTP[S] binding cannot be abolished by Mg^{2+} at concentrations up to 10 mM. Higher Mg^{2+} concentrations could not be tested, because agonist stimulation of GTP[S] binding then becomes negligible. The low-affinity Mg^{2+} site may also be the target for Li^+ in *Dictyostelium*, and a speculative explanation for both the stimulatory and inhibitory effects of this ion on GTP[S] binding, could be that it partially mimics Mg^{2+} -induced GDP/GTP exchange, but prevents further activation of the G protein, for instance by blocking $\beta\gamma$ subunit dissociation. The major conclusion, relevant to the present study is however, that G proteins are evidently a putative

Fig. 6. Effects of Li^+ on cGMP phosphodiesterase or cGMP adaptation. (A) Aggregation-competent cells of *stm F* mutant NP368 were incubated for 15 min with 0 (○), 2 (△) or 10 (□) mM LiCl and subsequently stimulated with 100 nM cAMP. cGMP levels were measured at the indicated times. Values are means $\pm$ SEM, $n = 2$ (100% $\approx 90 \pm 6$ pmol cGMP/ 10^7 cells). (B) Aggregation-competent NC4 cells were incubated for 15 min with 0, 2 or 10 mM LiCl . Cells were stimulated for 10 s with a saturating stimulus of 100 nM cAMP to induce a maximal cGMP response. To measure adaptation, cells were first stimulated at $t = 0$ s with a suboptimal stimulus of 200 nM cAMP[S] (comparable with about 2 nM cAMP) and restimulated at 30 s with 100 nM cAMP. cGMP levels were measured after 0, 10, 30 and 40 s of stimulation. Data are expressed as percentage of cGMP levels reached after 10 s stimulation with 100 nM cAMP in the absence of LiCl . Values are means $\pm$ SEM.

target for the effects of Li^+ on signalling and differentiation in *Dictyostelium*.

It is at present not clear, which G proteins are inhibited by Li^+ . Eight different G-protein α subunits have been cloned in *Dictyostelium* (Pupillo et al., 1989; Hadwiger et al., 1991; Wu and Devreotes, 1991), but a function for only one G protein has been indicated. One of the first cloned G proteins, $\text{G}\alpha 2$, probably mediates cAMP-induced phospholipase-C activation (Kesbeke et al., 1988; Snaar-Jagalska et al., 1988c; Kumagai et al., 1989). Since Li^+ effectively inhibits cAMP-induced $\text{Ins}(1,4,5)\text{P}_3$ production (Peters et al., 1989), $\text{G}\alpha 2$ is a possible target for Li^+ .

cAMP-stimulated GTP[S] binding and GTPase in aggregative cells are partially inhibited by pertussis toxin and may represent an inhibitory G protein controlling adenylylcyclase activity, since pertussis toxin abolishes GTP[S]-induced inhibition of this enzyme (Van Haastert et al., 1987; Snaar-Jagalska et al., 1988a, b). Li^+ reduces adenylylcyclase activation, and it is therefore less likely that this pertussis-toxin-sensitive G protein is inhibited by Li^+ .

The G_s protein for *Dictyostelium* adenylylcyclase has not yet been identified, but could be a putative Li^+ target. If so, we would expect Li^+ inhibition of adenylylcyclase activation to be more severe than the observed 50% reduction at 10 mM LiCl , which saturates inhibition of G-protein activation.

Involvement of G proteins in guanylyl-cyclase activation has not yet been unequivocally demonstrated (Janssens et al., 1989). Remarkably, Li^+ stimulates cAMP-induced accumulation of cGMP. This effect is not due to inhibition of cGMP phosphodiesterase, because in *stm F* mutant NP368, which lacks cGMP phosphodiesterase, the stimulatory effect is much more pronounced than in wild-type cells. Apparently, hydrolysis of cGMP in wild-type cells largely overrules the

stimulatory effect of LiCl . Since Li^+ does also not inhibit cAMP-induced adaptation of guanylyl cyclase, it most likely stimulates activation of the enzyme.

The observation that Li^+ inhibits $\text{Ins}(1,4,5)\text{P}_3$ accumulation, but stimulates guanylyl-cyclase activation contradicts an earlier hypothesis that guanylyl cyclase is activated via the $\text{Ins}(1,4,5)\text{P}_3/\text{Ca}^{2+}$ pathway (Europe-Finner and Newell, 1985, 1986; Small et al., 1986; Snaar-Jagalska et al., 1988c). Further evidence against this mechanism are observations that submicromolar Ca^{2+} concentrations completely inhibit guanylyl-cyclase activity *in vitro* (Janssens et al., 1989) and that a partial cAMP antagonist, which cannot increase $\text{Ins}(1,4,5)\text{P}_3$ levels, nevertheless induces normal cGMP accumulation (Peters et al., 1991). In contrast to this hypothesis, it seems plausible that the cAMP-induced $\text{Ins}(1,4,5)\text{P}_3/\text{Ca}^{2+}$ increase actually serves to switch off the guanylyl-cyclase enzyme. The stimulatory effect of LiCl on guanylyl-cyclase activation could then be due to its inhibitory effect on $\text{Ins}(1,4,5)\text{P}_3/\text{Ca}^{2+}$ accumulation.

Relevance for Li^+ effects on gene expression

Li^+ effectively inhibits cAMP induction of prespore-gene expression, but potentiates cAMP induction of prestalk-gene and aggregative-gene expression (Van Lookeren Campagne et al., 1988b; Peters et al., 1989). cAMP-induced expression of these genes is mediated by surface cAMP receptors (Schaap and Van Driel, 1985; Oyama and Blumberg, 1986; Haribabu and Dottin, 1986; Mann and Firtel, 1987) and does not require adenylylcyclase activation (Schaap et al., 1986; Mann and Firtel, 1987). However, intracellular messengers controlling gene regulation have not yet been identified.

Li^+ inhibition of prespore-gene expression correlates with Li^+ inhibition of $\text{Ins}(1,4,5)\text{P}_3$ accumulation, suggesting that $\text{Ins}(1,4,5)\text{P}_3$ may mediate prespore-gene expression. This is supported by observations that introduction of $\text{Ins}(1,4,5)\text{P}_3$ and diacylglycerol in permeabilized cells enhances expression of prespore genes (Ginsburg and Kimmel, 1989).

Li^+ stimulates cAMP-induced prestalk-gene and aggregative-gene expression. The cAMP-induced production of cGMP is the only second-messenger response found so far, which is also stimulated by Li^+ . This suggests that intracellular cGMP may mediate induction of prestalk-gene and aggregative-gene expression. However, the stimulatory effect of Li^+ on the cGMP response is small, and further evidence is required to support involvement of cGMP in gene regulation. Since Li^+ abolishes cAMP-induced $\text{Ins}(1,4,5)\text{P}_3$ production, this response is most likely not involved in positive regulation of prestalk-gene and aggregative-gene expression.

It was suggested earlier that Li^+ effects on prespore-gene expression and inositolphosphate signalling result from inhibition of inositolphosphatases by Li^+ , causing depletion of inositol (Van Lookeren Campagne et al., 1988a; Peters et al., 1989; Berridge et al., 1989). We recently found that inositol up to 0.3 M cannot abolish Li^+ inhibition of prespore-gene expression (Brandt, R. and Schaap, P., unpublished results) and now consider it more likely that Li^+ inhibition of G-protein activation causes its effects on pattern formation.

We thank Raymond Brandt for excellent technical assistance. This work was supported by the Foundation for Biological Research (BION), which is subsidized by the Netherlands Organisation for Scientific Research (NWO).

REFERENCES

- Allison, J. H. & Stewart, M. A. (1971) *Nature New Biol.* 233, 267–268.
- Avissar, S., Schreiber, G., Danon, A. & Belmaker, R. H. (1988) *Nature* 331, 440–442.
- Avissar, S., Murphy, D. L. & Schreiber, G. (1991) *Biochem. Pharmacol.* 41, 171–175.
- Backstrom, S. (1954) *Ark. Zool.* 6, 527–536.
- Batty, I. & Nahorski, S. R. (1985) *J. Neurochem.* 45, 1514–1521.
- Berridge, M. J., Downes, P. C. & Hanley, M. R. (1989) *Cell* 59, 411–419.
- Breckenridge, L. J., Warren, R. L. & Warner, A. E. (1987) *Development* 99, 353–370.
- Bulgakov, R. & Van Haastert, P. J. M. (1982) *Biochim. Biophys. Acta* 756, 56–66.
- Busa, W. B. & Gimlich, R. L. (1989) *Dev. Biol.* 132, 315–324.
- Casebolt, T. L. & Jope, R. S. (1989) *Biol. Psychiatry* 25, 329–340.
- Cassel, D. & Selinger, Z. (1976) *Biochim. Biophys. Acta* 452, 538–551.
- Cassel, D. & Selinger, Z. (1978) *Proc. Natl Acad. Sci. USA* 75, 4155–4159.
- Delvaux, A., Dumont, J. E. & Erneux, C. (1987) *Biochem. Biophys. Res. Commun.* 145, 59–65.
- Dobner, P. R., Tischler, A. S., Lee, Y. C., Bloom, S. R. & Donahue, S. R. (1988) *J. Biol. Chem.* 263, 13983–13986.
- Ebstein, R. P., Hermoni, M. & Belmaker, R. H. (1980) *J. Pharmacol. Exp. Ther.* 213, 161–167.
- Europe-Finner, G. N. & Newell, P. C. (1985) *Biochem. Biophys. Res. Commun.* 130, 1115–1122.
- Europe-Finner, G. N. & Newell, P. C. (1987) *J. Cell. Sci.* 87, 221–229.
- Gilman, A. G. (1987) *Annu. Rev. Biochem.* 56, 615–649.
- Ginsburg, G. & Kimmel, A. R. (1989) *Proc. Natl Acad. Sci. USA* 86, 9332–9336.
- Godfrey, P. P., McClue, S. J., White, A. M., Wood, A. J. & Grahame-Smith, D. C. (1989) *J. Neurochem.* 52, 498–506.
- Hadwiger, J. A., Wilkie, T. M., Strathmann, M. & Firtel, R. A. (1991) *Proc. Natl Acad. Sci. USA* 88, 8213–8217.
- Hallcher, L. M. & Sherman, W. R. (1980) *J. Biol. Chem.* 255, 10896–10901.
- Haribabu, B. & Dottin, R. P. (1986) *Mol. Cell. Biol.* 6, 2402–2408.
- Higashijima, T., Ferguson, K. M., Sternweis, P. C., Smigel, M. D. & Gilman, A. G. (1987) *J. Biol. Chem.* 262, 762–766.
- Janssens, P. M. W., De Jong, C. C. C., Vink, A. A. & Van Haastert, P. J. M. (1989) *J. Biol. Chem.* 264, 4329–4335.
- Kao, K. R. & Elinson, R. P. (1989) *Dev. Biol.* 132, 81–90.
- Kay, R. R. (1982) *Proc. Natl Acad. Sci. USA* 79, 3228–3231.
- Kesbeke, F., Snaar-Jagalska, B. E. & Van Haastert, P. J. M. (1988) *J. Cell Biol.* 107, 521–528.
- Kumagai, A., Pupillo, M., Gundersen, R., Miake-Lye, R., Devreotes, P. N. & Firtel, R. A. (1989) *Cell* 57, 265–275.
- Maeda, Y. (1970) *Develop. Growth & Differ.* 12, 217–227.
- Mann, S. K. O. & Firtel, R. A. (1987) *Mol. Cell. Biol.* 7, 458–469.
- Mato, J. M. & Malchow, D. (1978) *FEBS Lett.* 90, 119–122.
- Mehdy, M. C., Ratner, D. & Firtel, R. A. (1983) *Cell* 32, 763–771.
- Mork, A. & Geisler, A. (1987) *Pharmacol. & Toxicol.* 60, 241–248.
- Oyama, M. & Blumberg, D. D. (1986) *Proc. Natl Acad. Sci. USA* 83, 4819–4823.
- Peters, D. J. M., Van Lookeren Campagne, M. M., Van Haastert, P. J. M., Spek, W. & Schaap, P. (1989) *J. Cell Sci.* 93, 205–210.
- Peters, D. J. M., Bominaar, A. A., Snaar-Jagalska, B. E., Brandt, R., Van Haastert, P. J. M., Ceccarelli, A., Williams, J. G. & Schaap, P. (1991) *Proc. Natl Acad. Sci. USA* 88, 9219–9223.
- Pupillo, M., Kumagai, A., Pitt, G. S., Firtel, R. A. & Devreotes, P. N. (1989) *Proc. Natl Acad. Sci. USA* 86, 4892–4896.
- Ross, E. M., Maguire, M. E., Sturgill, T. W., Biltonen, R. L. & Gilman, A. G. (1977) *J. Biol. Chem.* 252, 5761–5779.
- Ross, F. M. & Newell, P. C. (1981) *J. Gen. Microbiol.* 127, 339–350.
- Schaap, P. & Van Driel, R. (1985) *Exp. Cell Res.* 159, 388–398.
- Schaap, P. & Van Lookeren Campagne, M. M., Van Driel, R., Spek, W., Van Haastert, P. J. M. & Pinas, J. (1986) *Dev. Biol.* 118, 52–63.
- Schreiber, G., Avissar, S., Charanjit, S. A. & Murphy, D. L. (1990) *Neuropharmacology* 29, 1067–1071.
- Small, N. V., Europe-Finner, G. N. & Newell, P. C. (1986) *FEBS Lett.* 203, 11–14.
- Snaar-Jagalska, B. E., De Wit, R. J. W. & Van Haastert, P. J. M. (1988a) *FEBS Lett.* 232, 148–152.
- Snaar-Jagalska, B. E., Jakobs, K. H. & Van Haastert, P. J. M. (1988b) *FEBS Lett.* 236, 139–144.
- Snaar-Jagalska, B. E., Kesbeke, F. & Van Haastert, P. J. M. (1988c) *Dev. Genet.* 9, 215–226.
- Theibert, A. & Devreotes, P. (1986) *J. Biol. Chem.* 261, 15121–15125.
- Van Haastert, P. J. M. (1984) *Biochem. Biophys. Res. Commun.* 124, 597–604.
- Van Haastert, P. J. M. & Van der Heijden, P. R. (1983) *J. Cell Biol.* 96, 347–353.
- Van Haastert, P. J. M., Snaar-Jagalska, B. E. & Janssens, P. M. W. (1987) *Eur. J. Biochem.* 162, 251–258.
- Van Haastert, P. J. M., De Vries, M., Penning, L. C., Roovers, E., Van der Kaay, J., Erneux, C. & Van Lookeren Campagne, M. M. (1989) *Biochem. J.* 258, 577–586.
- Van Lookeren Campagne, M. M., Erneux, C., Van Eijk, R. & Van Haastert, P. J. M. (1988a) *Biochem. J.* 254, 343–350.
- Van Lookeren Campagne, M. M., Wang, M., Spek, W., Peters, D. & Schaap, P. (1988b) *Dev. Genet.* 9, 589–596.
- Wang, M., Van Driel, R. & Schaap, P. (1988) *Development* 103, 611–618.
- Whitworth, P. & Kendall, D. A. (1988) *J. Neurochem.* 51, 258–265.
- Wu, L. J. & Devreotes, P. N. (1991) *Biochem. Biophys. Res. Commun.* 179, 1141–1147.