University of Groningen

A POLYNUCLEAR COPPER(I) COMPLEX WITH A SINGLE HELICAL STRUCTURE

GELLING, OJ; VANBOLHUIS, F; FERINGA, BL

Published in:
Journal of the Chemical Society-Chemical Communications

DOI:
[10.1039/c39910000917](https://doi.org/10.1039/c39910000917)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1991

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

GELLING, OJ., VANBOLHUIS, F., & FERINGA, BL. (1991). A POLYNUCLEAR COPPER(I) COMPLEX WITH A SINGLE HELICAL STRUCTURE. *Journal of the Chemical Society-Chemical Communications*, 31(14), 917-919. <https://doi.org/10.1039/c39910000917>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

A Polynuclear Copper(I) Complex with a Single Helical Structure

O. J. Gelling, F. van Bolhuis and Ben L. Feringa*

Department of Organic Chemistry, University of Groningen, Nijenborgh 16, 9747 AG Groningen, The Netherlands

The synthesis and crystal and molecular structure of a single helical polynuclear Cu^I complex are described.

Molecular recognition by design and self-assembly of (supra)molecular structures are currently being extensively investigated and a crucial role for molecular helicity has been revealed.¹⁻⁴ Based on oligopyridines as ligands, spontaneous assembly of double helical polynuclear metal complexes has been demonstrated. Thus, ligands containing two to five

2,2'-bipyridine units separated by oxapropylene spacers form double helical Cu^I complexes^{3,5} and similar molecular topology was found in Ni^{II}, Cd^{II}, Pd^{II} and Cu^I complexes of various oligopyridines.^{4,6} A single stranded diruthenium(II) helical complex has recently been discovered by Constable, Tocher and co-workers.⁷

Scheme 1 THF = tetrahydrofuran

We report in this communication the synthesis and crystal and molecular structure of a single helical copper(I) coordination polymer. When the new ligand 2,6-bis[N-{2-(2-pyridyl)ethyl}formimidoyl]-1-methoxybenzene **1a** was allowed to react with two equivalents of Cu(MeCN)₄BF₄, dinuclear copper(I) complexes **2a-c** were formed.⁸ The formation of **2a-c** is strongly dependent upon the amount of MeCN present; heating of **2b** at reflux in CH₂Cl₂ for 10 min provided the acetonitrile-free and presumably two-coordinated dinuclear Cu^I complex **2c**. Addition of MeCN (4 equiv.) to a CHCl₃ solution of **2c** reconverted it into acetonitrile complex **2b** (80–90% yield).⁸ When, however, excess (≥10 equiv.) of MeCN was added to **2c** in CHCl₃ and the resulting solution allowed to crystallize at room temperature for a period of 50 h, white crystalline and orange crystalline material could be obtained in a 50 : 50 (±5) ratio. The white compound was Cu(MeCN)₄BF₄ **4**, whereas the orange compound showed a ligand-copper stoichiometry of 1 : 1 and a copper-nitrogen ratio of 1 : 2 indicating a mononuclear or polynuclear 1 : 1 complex. This complex, in solution, is very sensitive to molecular oxygen and when a 1 : 1 mixture of **3** and **4** in CHCl₃ is oxidized with O₂ the same oxygenated ligand⁸ **1b** is obtained as is the case with **2a-c**.⁹ These results indicate an intact bis-imine ligand in **3** and an equilibrium between **3+4** and **2a-c** in solution.

Bright-orange crystals of **3** suitable for X-ray analysis were obtained from CHCl₃-MeCN. The crystal and molecular

Fig. 1 (a) A view of the cation of **3** with adopted numbering scheme. Selected bond lengths (Å) and angles (°): Cu(1)–N(1) 2.097(5), Cu(1)–N(2) 2.032(5), Cu(1)–N(3)[†] 2.048(4), Cu(1)–N(4)[†] 2.073(5), Cu(1)–Cu(1)[†] 7.650(4); N(1)–Cu(1)–N(2) 97.0(2), N(1)–Cu(1)–N(3)[†] 107.4(2), N(1)–Cu(1)–N(4)[†] 109.6(2), N(2)–Cu(1)–N(3)[†] 131.2(2), N(2)–Cu(1)–N(4)[†] 111.9(2), N(3)[†]–Cu(1)–N(4)[†] 98.9(2). (b) Crystal structure of part of the coordination polymer **3**. (c) Schematic view of the single-helical arrangement (X = OMe).

structure of **3** is shown in Fig. 1.[†] Each Cu^I atom in **3** is four-coordinated with two pyridylethylimine bidentate units from two different ligand molecules **1a**, in a distorted tetrahedral geometry. A dihedral angle between the CuN(1)–N(2) and CuN(3)[†]N(4)[†] planes of 83.6° is observed. Upon

[†] Crystal data: C₂₄H₂₅N₄BCl₃F₄OCu, monoclinic, space group *P*2₁/*n*, *a* = 13.826(3), *b* = 9.938(2), *c* = 20.555(6) Å, β = 102.58(2)°, *U* = 2756.5 Å³, *Z* = 4, *D*_c = 1.55 g cm^{−3}, Mo-Kα radiation (λ = 0.71073 Å), μ(Mo-Kα) = 11.4 cm^{−1}.

Data were collected on a Nonius CAD4F diffractometer at 130 K in the range 1° ≤ θ ≤ 27°. 3618 Unique reflections with *I* ≥ 3σ(*I*) were used in the refinements. The structure was partly solved by direct methods; the remaining atoms including all the H-atoms were located in succeeding difference Fourier synthesis. Block-diagonal least-squares on *F*, with unit weights, converged to a final *R* = 0.059 and *R*_w = 0.065 using anisotropic thermal parameters for the non-hydrogen atoms. Atomic coordinates, bond lengths and angles, and thermal parameters have been deposited at the Cambridge Crystallographic Data Centre. See Notice to Authors, Issue No. 1.

coordination of Cu^{I} the ligand **1a** has formed a linear coordination polymer with two helical strains, lying next to each other in the infinite unit cell [Fig. 1(b)].

Furthermore, one BF_4^- anion and one molecule of CHCl_3 are present for each Cu^{I} unit. The copper–nitrogen distances are nearly the same (2.03–2.09 Å) and on the average slightly longer than the Cu–N distance in the three-coordinated Cu^{I} complex derived from **1c**⁹ as might be expected on the basis of steric repulsion. The Cu– N_{pyr} bonds (average 2.085 Å) are rather long compared to the Cu– N_{pyr} bond lengths generally found for four-coordinated Cu^{I} ions (2.0–2.05 Å)¹⁰ and those in helical (oligo-)bipyridine Cu^{I} complexes (average 2.02 Å).³ Not unexpectedly, the $\text{N}_{\text{imine}}\text{--Cu--N}_{\text{pyridine}}$ bond angles are relatively small (97.0, 98.9°) compared to the other four N–Cu–N angles (average 115.0°). The 1,3-bis-imine substituted anisole group bridges two Cu^{I} centres 7.650 Å apart; the Cu–Cu distance is considerably longer than the distance of 4.952 Å observed in the binuclear Cu^{I} complex derived from ligand **1c**.⁹

In complex **3** the Cu^{I} ions are an integral part of the polymer backbone and the two bidentate units of each ligand **1a** are twisted with respect to each other. Other polymers with backbone metals are known which form linear, planar or three-dimensional structures depending on the identity of the ligand and the coordination abilities of the metal ion.¹¹ As far as copper(I) coordination polymers are concerned, Reedijk and coworkers recently described a reaction of *N,N,N',N'*-tetrakis(pyrazol-1'-ylmethyl)-1,2-diaminoethane (tpde) with $\text{Cu}(\text{MeCN})_4\text{BF}_4$ which produced a white $\text{Cu}(\text{tpde})\text{BF}_4\text{MeCN}$ linear polymer. The Cu^{I} ions are coordinated by four pyrazole N-atoms in an almost tetrahedral arrangement.¹²

Fares and coworkers reported a polymeric cyano-2,2'-biquinoline- Cu^{I} complex containing both linear and tetrahedrally coordinated Cu^{I} in the chain.¹³ In the Cu^{I} coordination polymer **3** the two helices in the infinite unit cell are single stranded, one of which is left handed and the other right handed [schematically shown in Fig. 1(c)]. The ability of ligands such as **1a** to assemble in the presence of Cu^{I} ions to single stranded helical complexes provides flexible and synthetically easy accessible alternatives to double stranded

polypyridine Cu^{I} systems in approaches towards supramolecular recognition and electroactive systems. The fact that the methoxy substituent in the ligand points outside the helicate, thereby minimizing steric hindrance, allows attachment of functional units that arrange in a helical geometry. We are currently investigating these molecular arrangements.

Received, 3rd September 1990; Com. 0/03983G

References

- 1 J. M. Lehn, *Angew. Chem., Int. Ed. Engl.*, 1988, **27**, 89.
- 2 J. K. Barton, *J. Biomol. Struct. Dyn.*, 1983, **1**, 621; J. K. Barton, *Chem. Eng. News*, 1988, **66**, 30.
- 3 J. M. Lehn, A. Rigault, J. Siegel, J. Harrowfield, B. Chevrier and D. Moras, *Proc. Natl. Acad. Sci. USA*, 1987, **84**, 2565.
- 4 E. C. Constable, M. G. B. Drew and M. D. Ward, *J. Chem. Soc., Chem. Commun.*, 1987, 1600.
- 5 J. M. Lehn and A. Rigault, *Angew. Chem., Int. Ed. Engl.*, 1988, **27**, 1095.
- 6 E. C. Constable, M. D. Ward, M. G. B. Drew and G. A. Forsyth, *Polyhedron*, 1989, **8**, 2551; E. C. Constable, M. D. Ward and D. A. Tocher, *J. Am. Chem. Soc.*, 1990, **112**, 1256; E. C. Constable, S. M. Elder, J. Healy, M. D. Ward and D. A. Tocher, *J. Am. Chem. Soc.*, 1990, **112**, 4590.
- 7 C. J. Cathey, E. C. Constable, M. J. Hannon, D. A. Tocher and M. D. Ward, *J. Chem. Soc., Chem. Commun.*, 1990, 621.
- 8 O. J. Gelling and B. L. Feringa, *J. Am. Chem. Soc.*, 1990, **112**, 7599.
- 9 O. J. Gelling, F. van Bolhuis, A. Meetsma and B. L. Feringa, *J. Chem. Soc., Chem. Commun.*, 1988, 552.
- 10 K. D. Karlin, R. W. Cruse, Y. Gultneh, A. Faroog, J. C. Hayes and J. Zubietta, *J. Am. Chem. Soc.*, 1987, **109**, 2668, and references cited therein.
- 11 B. M. Foxman and S. W. Gersten in *Encyclopedia of Polymer Science and Engineering*, ed. J. I. Kroschwitz, 1985, Wiley, New York, vol. iv, p. 175; J. T. Wroblewski and D. B. Brown, *Inorg. Chem.*, 1979, **18**, 498; M. Hanack, *Chimia*, 1983, **37**, 238.
- 12 W. L. Driessen, F. B. Hulsbergen, J. Reedijk and G. C. Verschoor, *Transition Met. Chem.*, 1985, **10**, 390.
- 13 G. Dessy, V. Fares, P. Imperatori and G. O. Morpurgo, *J. Chem. Soc., Dalton Trans.*, 1985, 1285.