University of Groningen

Synthesis and Physicochemical Properties of [19,20-13C]-17 α -Ethinylestradiol

Kraan, G.P.B.; Drayer, N.M.; Kruizinga, W.H.; Vaalburg, W.; Hummelen, J.C.

Published in:
 Synthesis and Applications of Isotopically Labelled Compounds

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 1989

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Kraan, G. P. B., Drayer, N. M., Kruizinga, W. H., Vaalburg, W., & Hummelen, J. C. (1989). Synthesis and Physicochemical Properties of [19,20-13C]-17 α -Ethinylestradiol. In *Synthesis and Applications of Isotopically Labelled Compounds: Proceedings of the Third International Symposium, Innsbruck, Austria, 17 -21 July 1988* (pp. 705-708). Elsevier.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

SYNTHESIS AND PHYSICOCHEMICAL PROPERTIES OF [19, 20-¹³C]-17 α -ETHINYLESTRADIOL

G. P. B. KRAAN¹, N. M. DRAYER¹, J. KOEK², W. H. KRUIZINGA², W. VAALBURG² and J. C. HUMMELEN².

1. Department of Pediatrics, University of Groningen, Oostersingel 59, 9713 EZ Groningen (The Netherlands)
2. Department of Organic Chemistry, University of Groningen, Nijenborgh 16, 9747 AG, Groningen (The Netherlands)

SUMMARY

¹³C₂-17 α -ethinylestradiol (¹³C₂-EE₂) was synthesized from estrone and ¹³C₂-C₂H₂-gas to measure the metabolic clearance rate and the plasma concentration of 17 α -ethinylestradiol (EE₂) in tall girls, who are treated with high dosages of this estrogen. Interesting characteristics determined by (i) MS: the molecular ion m/z 298.183 is the only labelled ion in the spectrum, and by (ii) ¹H-NMR δ (ppm) 2.96 (dd, ¹J(¹³C-H)=249 Hz, ²J(¹³C-H)=49.6 Hz; 1H; H₂₀) and (iii) ¹³C-NMR δ 88.8 (dd, ¹J(¹³C-¹³C)=166 Hz, ²J(¹³C-H₂₀)=49.8 Hz; C₁₉); 80.2 (dd, ¹J(¹³C₁₇-¹³C₁₉)=73 Hz, ²J(¹³C₁₇-¹³C₂₀)=12.3 Hz; C₁₇) and 74.6 (dd, ¹J(¹³C-¹³C)=166 Hz, ¹J(¹³C-H)=249 Hz; C₂₀) confirm the synthesis and structure of ¹³C₂-EE₂.

INTRODUCTION

Administration of estrogens to healthy, (pre)pubertal tall girls will cause a decreased growth rate, accelerated sexual maturation, epiphyseal fusion and, as desired, a reduced final adult height (ref. 1).

In order to keep the potential complications of the estrogen therapy, existing of 2x 50 μ g 17 α -ethinylestradiol (EE₂) per os daily, as low as possible it would be necessary to measure carefully the individual concentration time relationship of EE₂ in plasma in relation to growth rate, skeletal maturation and other parameters. For this purpose we decided to synthesize ¹³C-labelled 17 α -ethinylestradiol (¹³C₂-EE₂) according to the previously published method (ref.2).

In this paper the synthesis and main physicochemical characteristics of ¹³C₂-EE₂ are described.

METHODS

Instrumentation

Centrifugal liquid chromatography was performed on a Hitachi CLC-5 instrument. A rotating disk of 3 mm silicagel 60-G (Merck, Darmstadt, G.) and 10 cm path length was eluted with chloroform (Merck, p.a.) at a rate of 8.3 ml. min⁻¹ at 480 rpm approximately. Mass spectra were recorded on a

AEI-MS-9 spectrometer. The infrared spectrum was recorded on a Unicam SP-200 spectrophotometer in a KBr pellet. Nuclear magnetic resonance spectra were recorded on a Varian-VXR-300 at 300 MHz for ^1H -NMR and at 75.48 MHz for ^{13}C -NMR in CD_3OD . ^1H -chemical shifts are reported in δ units (ppm) relative to TMS and ^{13}C -chemical shifts relative to CD_3OD .

Synthesis

Approximately 3 mmole $^{13}\text{C}_2\text{-C}_2\text{H}_2$ gas (MSD Isotopes) was condensed in a 250 ml three necked flask containing a mixture of 1.5 ml 1.5 N n-Buli/n-hexane and 15 ml distilled THF at -115°C . During warming up to 0°C the mixture was stirred. Then a solution of 1.1 mmole estrone (Aldrich) in 20 ml dry THF was added dropwise and the mixture was stirred during 30 minutes at 0°C and 15 minutes at 40°C . After cooling to room temperature a solution of 12 g NH_4Cl in 60 ml H_2O was added dropwise and the resulting mixture was stirred 30 minutes. The product was extracted with CHCl_3 (2x 100 ml) and the organic layer was dried on MgSO_4 , filtered and evaporated to dryness to yield 365 mg of crude steroid as a white powder. Spectroscopically pure $^{13}\text{C}_2\text{-EE}_2$ was obtained after chromatography on a Hitachi CLC-5 chromatograph using CHCl_3 as the eluent and monitoring at 254 nm. After two runs 0.67 mmole of pure $^{13}\text{C}_2\text{-EE}_2$ was obtained as a white mass.

RESULTS

Fig.1 shows the structure and the numbering of the carbon atoms of $^{13}\text{C}_2\text{-EE}_2$ (^{13}C indicated by an asterix).

The mass spectra of $^{13}\text{C}_2\text{-EE}_2$ (a) and EE_2 (b) are shown in fig. 2. Apart from very small fragments at $m/z > 250$ the only labelled fragment is the molecular ion of m/z 298. The exact mass of the molecular ion was measured and calculated to be 298.183 and 298.184, respectively.

Absorbance maxima of $^{13}\text{C}_2\text{-EE}_2$ in the infrared were recorded at 3350 (s), 2950 (s), 1620 (m), 1510 (m), 1460 (m), 1290 (m), 1240 (m) and 1025 (s) cm^{-1} .

Fig.1. [19,20- ^{13}C]-17 α -ethinyl-estra-1,3,5-triene-3,17 β -diol

Fig.2. Electron impact mass spectra (70eV) of $^{13}\text{C}_2\text{-EE}_2$ (a) and EE_2 (b)

$^1\text{H-NMR}$ of $^{13}\text{C}_2\text{-EE}_2$ was recorded in CD_3OD at 300 MHz. The chemical shifts are given in δ units (ppm) relative to TMS: 7.21(d, $J=8.4$ Hz; 1H; H_2); 6.72(d, $J=8.4$ Hz; 1H; H_1); 6.65(s, 1H; H_4); 2.96(dd, $^1J(^{13}\text{C-H})=249$ Hz, $^2J(^{13}\text{C-H})=49.6$ Hz; 1H; H_{20}); 2.9(m, 2H); 2.45(m, 2H); 2.33-1.8(m, 7H); 1.65-1.3(m, 4H); 1.00(s, 3H; H_8).

$^{13}\text{C-NMR}$ of $^{13}\text{C}_2\text{-EE}_2$ was recorded in CD_3OD at 75.48 MHz. The chemical shifts are given in δ units (ppm) relative to CD_3OD . The assignments of these shifts to the carbon atoms were derived from the APT spectrum and the coupled and uncoupled spectra. The uncoupled spectrum (fig. 3) shows strong resonance at 88.8(dd, $^1J(^{13}\text{C}-^{13}\text{C})=166$ Hz, $^2J(^{13}\text{C}-\text{H}_{20})=49.8$ Hz; C_{19}); 74.6(dd, $^1J(^{13}\text{C}-^{13}\text{C})=166$ Hz, $^1J(^{13}\text{C-H})=249$ Hz; C_{20}),

Fig . 3. ^{13}C -NMR spectrum of $^{13}\text{C}_2$ -EE₂ at 75.48MHz in CD_3OD .

The result of coupling between $^{13}\text{C}_{17}$ and $^{13}\text{C}_{19}$ and $^{13}\text{C}_{20}$ around 80 ppm is shown in the insert.

and weak resonance at 155.5(s; C_3), 138.7(s; C_5); 132.6(s; C_{10}); 127.1(d; C_1); 116.1(d; C_4); 113.7(d; C_2); 80.2(dd, $^1J(^{13}\text{C}_{17}-^{13}\text{C}_{19})=73\text{ Hz}$, $^2J(^{13}\text{C}_{17}-^{13}\text{C}_{20})=12.3\text{ Hz}$; C_{17}); 50.7(d; C_{14}); 48.1(s; C_{13}); 44.8(d; C_9); 40.9(d; C_8); 39.9(t; C_{16}); 133.9(t; C_{12}); 30.5(t; C_6); 28.5(t; C_7); 27.6(t; C_{11}); 23.6(t; C_{15}); 13.3(q; C_{18}).

CONCLUSION

From the characteristics of the steroid determined by IR, MS and NMR it can be concluded that indeed $[19,20-^{13}\text{C}]\text{-EE}_2$ is synthesized.

REFERENCES

- ¹ N.J. Charest and J.J. Van Wyk, Tall stature, in: S.S. Gellis and B.M. Kagan (Eds), Current Pediatric Therapy, W.B. Saunders Cy, Philadelphia, 1986, pp. 299-301.
- ² W. Vaalburg, S. Reiffers, E. Beerling et al., The preparation of carbon-11 labeled 17α -ethinylestradiol, J. Label Compds Radiopharm 12 (1977), 200-201.