University of Groningen

VENTROMEDIAL HYPOTHALAMIC REGULATION OF HORMONAL AND METABOLIC RESPONSES TO EXERCISE

Vissing, John; Wallace, Jo L.; Scheurink, Anton J.W.; Galbo, Henrik; Steffens, Anton B.

Published in:

The American Journal of Physiology - Regulatory, Integrative and Comparative Physiology

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

1989

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Vissing, J., Wallace, J. L., Scheurink, A. J. W., Galbo, H., & Steffens, A. B. (1989). VENTROMEDIAL HYPOTHALAMIC REGULATION OF HORMONAL AND METABOLIC RESPONSES TO EXERCISE. *The American Journal of Physiology - Regulatory, Integrative and Comparative Physiology*, 256(5), R1019-R1026.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

J. Vissing, J. L. Wallace, A. J. Scheurink, H. Galbo and A. B. Steffens
Am J Physiol Regulatory Integrative Comp Physiol 256:1019-1026, 1989.

You might find this additional information useful...

This article has been cited by 2 other HighWire hosted articles:

Delineation of responsive AVP-containing neurons to running stress in the hypothalamus

T. Saito and H. Soya

Am J Physiol Regulatory Integrative Comp Physiol, March 1, 2004; 286 (3): R484-R490.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Influence of muscle glycogen content on metabolic regulation

S. M. Weltan, A. N. Bosch, S. C. Dennis and T. D. Noakes

Am J Physiol Endocrinol Metab, January 1, 1998; 274 (1): E72-E82.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Medline items on this article's topics can be found at

<http://highwire.stanford.edu/lists/artbytopic.dtl>

on the following topics:

Biochemistry .. Glycogenolysis

Oncology .. Corticosterone

Medicine .. Insulin

Physiology .. Exertion

Medicine .. Fitness (Physical Activity)

Medicine .. Exercise

Additional material and information about *American Journal of Physiology - Regulatory, Integrative and Comparative Physiology* can be found at:

<http://www.the-aps.org/publications/ajpregu>

This information is current as of September 16, 2009 .

Ventromedial hypothalamic regulation of hormonal and metabolic responses to exercise

JOHN VISSING, JO L. WALLACE, ANTON J. W. SCHEURINK,
HENRIK GALBO, AND ANTON B. STEFFENS

*Department of Medical Physiology B, The Panum Institute, University of Copenhagen,
DK-2200 Copenhagen, Denmark; and Department of Animal Physiology,
State University of Groningen, 9700 AA Haren, The Netherlands*

VISSING, JOHN, JO L. WALLACE, ANTON J. W. SCHEURINK, HENRIK GALBO, AND ANTON B. STEFFENS. *Ventromedial hypothalamic regulation of hormonal and metabolic responses to exercise*. Am. J. Physiol. 256 (Regulatory Integrative Comp. Physiol. 25): R1019-R1026, 1989.—Recent studies have indicated a neural regulation of hormonal and metabolic responses to exercise. Studies on the ventromedial hypothalamus (VMH) suggest that the VMH might be involved in neural control of exercise metabolism. We therefore studied 25 rats with or without Marcain-anesthetized VMH (Marcain and control rats, respectively) at rest or during treadmill running (26 m/min). The rats had cannulas aimed bilaterally at the VMH and chronic catheters in the left jugular vein and right carotid artery. At rest, glucose turnover and plasma concentrations of glucose, lactate, glycerol, epinephrine, norepinephrine, insulin, and corticosterone remained stable in control rats, whereas in Marcain rats these parameters, except for unchanged insulin and glucose utilization, increased after Marcain administration along with increases in hepatic and muscular glycogenolysis. During exercise, glucose turnover, hepatic and muscular glycogenolysis, and plasma concentrations of glucose, lactate, glycerol, catecholamines, and corticosterone increased, and insulin decreased in both Marcain and control rats. However, during exercise, initial hepatic glucose production, plasma catecholamines, and subsequent plasma glucose concentrations and overall hepatic glycogenolysis were lower in Marcain compared with control rats. In conclusion, glucose mobilization is enhanced by the VMH during exercise. The results suggest that decreased and increased activity in sympathetic inhibitory and facilitating VMH areas, respectively, are involved in the VMH's response to exercise.

rats; ventromedial hypothalamus; catecholamines; blood glucose homeostasis; glucose turnover; insulin; glucagon; corticosterone

DURING EXERCISE, the traditional view has been that mobilization of extramuscular fuel stores is regulated exclusively by sensitive blood-borne metabolic feedback mechanisms, primarily relying on changes in plasma glucose concentrations (5, 10). More recently, however, evidence has been presented in favor of a neural feedforward regulation of substrate mobilization during exercise (7, 8, 11, 22, 25, 29–31). In accordance with neural regulation of substrate mobilization, glucose production increases faster than glucose uptake from plasma in fed running and swimming rats, giving rise to an increase in

plasma glucose concentration during exercise (22, 25, 29–31). Furthermore, when glucose availability is artificially increased or decreased in exercising rats, counterregulatory changes in hepatic glucose production do not compensate accurately for the inflicted changes in glucose availability (1, 29). These results in rats demonstrate inaccuracy of metabolic feedback systems in matching fuel mobilization to the energy requirements of working muscles. The studies provide indirect evidence for a neural feedforward regulation of substrate mobilization during exercise.

Based on previous findings concerning the ventromedial hypothalamus (VMH), we hypothesized in the present study that the VMH can be involved in a neural feedforward regulation of substrate mobilization during exercise. The ventromedial hypothalamus is recognized as an important mainly sympathetic center involved in the regulation of intermediate metabolism (24). Ablation of the VMH produces gross metabolic abnormalities such as hyperinsulinemia, hyperphagia, and obesity (9, 28, 35). Electrical or adrenergic stimulation of the VMH accelerates lipolysis (12, 24, 27), the release of glucagon, which accelerates glycogenolysis and gluconeogenesis, leading to hyperglycemia (6, 12, 23, 27). The numerous studies that have demonstrated these potentially functional properties of the VMH in the control of intermediate metabolism are supported by recent anatomic mapping studies that show connections between the VMH and the peripheral sympathetic nervous system (13).

We therefore studied the hormonal and metabolic responses to running in brain-cannulated conscious rats during bilateral anesthesia of the VMH. An impaired mobilization of substrates and hormones during exercise in VMH-anesthetized rats compared with controls would indicate an involvement of the VMH in the regulation of metabolism during exercise.

MATERIALS AND METHODS

Animal preparation. Twenty-five male Wistar rats weighing 273 ± 3 (SE) g were used for the experiments. The rats were kept individually at a room temperature of 23°C and with a light cycle from 6:00 A.M. to 6:00 P.M. One week before the experiment, all studied groups of rats had chronic brain cannulas implanted during Immobilon vet anesthesia (Pharmacia, Denmark, 0.01 ml/

100 g body wt sc). Atropine (0.05 mg/100 g body wt sc) was given to prevent bradycardia and bronchospasm. The implantation of brain cannulas was performed with the rats placed in a stereotaxic apparatus (Narashige, SR-6, SM-11 micromanipulator). The skin covering the skull was incised in the midline, and a 3×3 mm craniotomy was performed using a high-speed dental drill. Bleeding was negligible during these procedures, using Spongostan (Ferrosan, Denmark) for hemostasis. Sterile stainless steel cannulas (20.0 mm length, 0.3 mm OD, 0.1 mm ID) were stereotactically implanted bilaterally so that the tip of the cannulas were aimed at the dorsal aspect of the VMH. The lateral coordinate (± 0.7 mm) was found by using the sagittal sinus as zero. The dorsoventral coordinate (9.2 mm) was found by using the surface of the skull as zero. Bregma and the interaural line were used as points of reference for the determination of anteroposterior coordinates. When bregma-interaural distance in the rats differed from the atlas distance (18), the coordinates were adjusted. Anteroposterior coordinates were thus 5.8 ± 0.1 (SE) mm ($n = 25$) from the interaural line. The cannulas were fixed to the skull with a glue (Cyanolit, 3M a/s, Denmark) and dental acrylic. Sterile stainless steel obturators, 0.07 mm in diameter and with exactly the same length as the cannulas, were inserted into each cannula to prevent clotting of the lumen. The ends of the cannulas protruding from the dental acrylic were protected by a 21-gauge sleeve of stainless steel except for the top 3 mm where polyethylene caps closed the cannulas. The skin incision was closed with surgical sutures so that only the polyethylene caps covering the cannulas protruded from the skin.

Postoperatively, ampicillin (5 mg/100 g body wt sc) was given to prevent infections, and Revivon vet (Pharmacia, 0.01 ml/100 g body wt sc) was given as an Immobilon antidote to shorten time of anesthesia. The 2nd to the 5th day after the operation, the rats were accustomed to running on a rodent treadmill at a speed of 26 m/min for 20 min. The 6th day, 14–16 h before the experiment, the rats had polyethylene catheters inserted into the left common carotid artery and the right external jugular vein during Immobilon vet anesthesia, as previously described (30). Before the operation, the rats weighed 287 ± 3 g, which was not different from their weight (293 ± 3 g) before brain cannulation 6 days earlier. Postoperatively, the food was removed to avoid an intestinal contribution to endogenous glucose production during the experiments. To minimize hepatic glycogen depletion during this period, 2.5 g of fructose dissolved in 1.5 ml water was given postoperatively via a stomach tube. The rats lost 14 ± 1 g in body weight during this period.

Experimental protocol. The experiment was performed on the 7th day after brain cannulation. Polyethylene caps and obturators were removed, and each brain cannula was connected to a catheter flame-sealed at one end and filled with a Marcain solution (5 mg/ml, bupivacaine, Astra, Sweden). The rats were placed on a rodent treadmill with arterial, venous, and brain catheters extended to the outside. A priming dose of $2.1 \mu\text{Ci D-[3-}^3\text{H]glucose}$ (NET 331, New England Nuclear) followed by a constant

rate infusion of 93 nCi/min was delivered intravenously by a precision pump (Unita 1b, B. Braun, FRG). Blood sampling was started 60 min after the onset of the tracer infusion ($t = -20$ min). The rats were randomly assigned to resting or running groups infused with $3 \mu\text{g}$ Marcain/min for 5 min in each VMH (Marcain rats) or to resting or running control groups receiving no infusion (control rats). A control group receiving no infusion was chosen for two reasons: 1) the pH of the Marcain vehicle and Marcain solution differs (12.0 vs. 6.5, respectively) and, most importantly, 2) a possible solvent effect in rats infused with the vehicle alone most probably would not manifest itself in Marcain rats because of anesthesia of the infused neural tissue. Marcain was chosen as the anesthetic agent because 1) the onset of action is rapid, 5–10 min; 2) the drug is hydrophilic, a fact which reduces diffusion after application; and 3) the duration of anesthesia is longer for Marcain (4–10 h) than for any other commonly used local anesthetic. Marcain infusion was started at $t = -13$ min (see figures) and was delivered by two $5\text{-}\mu\text{l}$ Hamilton syringes positioned in a precision pump. To ensure that Marcain rats received the infusate, the distance traveled by an air bubble in each brain catheter was measured. At $t = 0$ min, Marcain and control rats either continued to rest on the treadmill or started running at a speed of 26 m/min for 20 min.

Tissue sampling and histology. Immediately after the 20 min of exercise or rest the rats were anesthetized with pentobarbital sodium (5 mg/100 mg body wt intra-arterially), and liver and muscle biopsies for determination of glycogen content were taken within 3 min, freeze-clamped with aluminum tongs precooled in liquid N_2 , and stored at -80°C until analysis. Muscle biopsies were taken from the superficial (fast-twitch white) and deep (fast-twitch red) part of the medial gastrocnemius muscle and from the soleus muscle (consisting mainly of slow-twitch red fibers). The rats were decapitated, and the heads were fixated in a 4% Formalin solution for 1 wk after which the brains were removed from the crania. After a few days in a 30% sucrose solution, the brains were quickly frozen and cut vertically in $40\text{-}\mu\text{m}$ slices on a cryostat microtome. The slices were stained with a 1% solution of cresyl fast violet and examined under a light microscope for determination of cannula placement. All 25 rats used in the study had the tip of the brain cannulas bilaterally situated just above the VMH. Three rats were excluded from the study due to wrong cannula placement.

Blood sampling and analytical procedures. Blood samples were taken frequently throughout the experiment as depicted in Figs. 1–3. Sampled blood was replaced by citrated blood from Immobilon-anesthetized donor rats. Treatment of blood samples and analytical procedures for glucagon, glucose, glycogen, and $[^3\text{H}]$ glucose have been described recently (30). The supernatant of $100 \mu\text{l}$ plasma and $200 \mu\text{l}$ 0.33 N perchloric acid was used for measuring glycerol and lactate concentrations by enzymatic fluorometric methods (4, 17). Plasma insulin was determined by radioimmunoassay using guinea pig serum M8309 as antiserum (NOVO, Denmark) (21). Plasma corticosterone, epinephrine, and norepinephrine concentrations were determined by high-pressure liquid chro-

matography in combination with ultraviolet light detection for corticosterone and electrochemical detection for the catecholamines (21). Glucose production and utilization were calculated according to Steele's equations (26, 30). The volume of distribution in which rapid changes in glucose concentration and specific activity of [^3H]glucose take place was set to 188 ml/kg body wt (19).

Statistically significant ($P < 0.05$, two-tailed testing) changes with time in glucose turnover and plasma concentrations of glucose, catecholamines, insulin, lactate, and glycerol were assessed by Friedman's nonparametric analysis of variance. Wilcoxon's nonparametric rank-sum test for paired data was applied to locate the differences when significant changes with time existed. Remaining data were evaluated by Wilcoxon's rank-sum test for paired and Mann-Whitney's rank-sum test for unpaired data.

RESULTS

Glucose turnover (Fig. 1). In resting control rats, turnover and plasma concentration of glucose were constant throughout the whole experiment. In resting Marcain rats, hepatic glucose production increased over basal values after Marcain infusion, the increase being significant in the time interval -5 to 5 min ($P < 0.05$). Glucose utilization remained unchanged throughout the experiment, and a large increment in plasma glucose concentration, significant from $t = -5$ min, was accordingly observed in resting Marcain rats.

In exercising control rats, hepatic glucose production increased rapidly during exercise, exceeding the increase in glucose utilization and thus giving rise to an increase in plasma glucose concentration during running. Production, utilization, and plasma concentrations of glucose were elevated over resting levels from $t = 2.5$ min and throughout the rest of the exercise period ($P < 0.05$). During the first 2.5 min of exercise, the increase in hepatic glucose production was insignificant in Marcain rats ($P > 0.10$) and tended ($P < 0.10$) to be smaller than in control rats. Hepatic production remained smaller in Marcain compared with control rats throughout the exercise period ($P < 0.05$ in the time interval 2.5–10 min). Glucose utilization was smaller in Marcain compared with control rats during the first 2.5 min of exercise. During continued exercise, however, glucose utilization did not differ between Marcain and control rats. Correspondingly, the exercise-induced increase in plasma glucose concentration was significantly attenuated in Marcain compared with control rats after 10 min of exercise ($P < 0.05$).

Hormonal responses (Fig. 2 and Table 1). Plasma levels of norepinephrine, epinephrine, insulin, and corticosterone were stable throughout the experiment in resting control rats. In resting Marcain rats, norepinephrine, epinephrine, and corticosterone levels increased after Marcain administration ($P < 0.05$), whereas insulin remained unchanged. During running, insulin concentrations decreased ($P < 0.05$), and corticosterone and catecholamine concentrations increased ($P < 0.05$) in both groups. Plasma insulin and corticosterone concentrations did not differ between exercising Marcain and

control rats. The initial increase in both catecholamines at $t = 2.5$ min, however, was insignificant in Marcain rats and smaller in Marcain compared with control rats ($P < 0.05$). During continued exercise, both catecholamines tended to be lower in Marcain compared with control rats. Plasma levels of glucagon were higher at rest in Marcain compared with control rats at the end of the experiment. In the 20th min of exercise, plasma levels of glucagon were similar in Marcain and control rats and higher than corresponding resting values.

Lactate and glycerol (Fig. 3). Plasma levels of lactate and glycerol were stable throughout the experiment in resting control rats. In resting Marcain rats, both lactate and glycerol increased after Marcain administration (at $t = 2.5$ min and $t = -5$ min, respectively, $P < 0.05$). In exercising rats, lactate and glycerol concentrations increased over resting levels and remained elevated throughout the exercise period ($P < 0.05$), the concentrations always being similar in Marcain and control rats.

Glycogen metabolism (Table 2). In resting rats, liver glycogen tended ($P < 0.10$) to be, and muscle glycogen content was, lower in Marcain compared with control rats. In rats of both exercising groups, postexercise liver and muscle glycogen concentrations were lower than glycogen concentrations in corresponding resting groups. Postexercise glycogen levels were similar in the two groups.

DISCUSSION

In the present study, anesthetic ablation of the VMH in conscious rats was used to examine the role played by this region in the regulation of glucose mobilization during exercise. The principal new finding is that the exercise-induced increases in plasma concentrations of norepinephrine, epinephrine, and glucose as well as hepatic glucose production are attenuated by anesthesia of the VMH. These results indicate that the VMH is involved in the regulation of glucose production during exercise at least in part by enhancing epinephrine release from the adrenal medulla, epinephrine being a potent stimulator of hepatic glucose production during exercise (25, 32).

Plasma concentrations of catecholamines and hepatic glucose production were lower in Marcain than in control rats throughout exercise. The difference did not, however, achieve statistical significance late during exercise, but in this period glucose levels were lower in Marcain compared with control rats, which may in a feedback manner have elicited compensatory increases in plasma concentrations of norepinephrine, epinephrine, glucagon, corticosterone, and decreases in plasma insulin, with subsequent increases in hepatic glucose production (2, 3, 8). Thus the importance of the VMH in regulation of catecholamine release and mobilization of glucose during exercise may have been underestimated. Furthermore, exercise-induced changes in the release of hormones from the endocrine pancreas and from corticotropes in the pituitary may also be elicited by the VMH. This may be so, since the similar plasma levels of glucagon, insulin, and corticosterone during exercise in Marcain and con-

FIG. 1. Glucose production and utilization and plasma glucose concentration in control ($n = 7$) and Marcain-infused ($n = 6$) rats studied at rest only and in control ($n = 6$) and Marcain-infused ($n = 6$) rats studied during a 20-min resting period followed by 20 min of running at a speed of 26 m/min. Marcain was infused in ventromedial hypothalamus. Values are means $\pm$ SE. * Difference between control and Marcain-infused rats in resting or exercising groups.

control rats, while glucose levels were lower in Marcain rats, indicate a relative inhibition of glucagon and corticosterone release and a relative enhancement of insulin release in Marcain rats (8). The VMH may also directly increase sympathetic outflow to the liver during exercise and thus activate hepatic glycogenolysis directly (24). The importance of liver nerves in the regulation of hepatic glucose production during exercise in rats, however, is still unsettled (25, 32).

The findings in the present study are supported by anatomic studies and functional studies at rest that have demonstrated numerous connections between the VMH and the peripheral sympathetic nervous system (6, 12, 13, 24, 27). Furthermore, the involvement of the VMH in regulating the mobilization of glucose during exercise

is supported by the recent finding in swimming rats of an abolished exercise-induced increase in plasma glucose concentration when the VMH was infused with the α -adrenergic blocker phentolamine (22).

Exercise-induced increases in sympathetic outflow from the VMH may be elicited by stimulation from higher brain areas (central command) that, in parallel, activate locomotion and substrate mobilization (7, 8, 30, 31). More direct evidence has recently been presented in favor of such a central command regulation of substrate mobilization during exercise (11, 16). Alternatively or additionally, increased sympathetic outflow from the VMH during exercise may have been elicited by neural feedback from working muscles. Neural feedback from muscles and central neural feedforward mechanisms

FIG. 2. Plasma concentrations of norepinephrine, epinephrine, and insulin in control ($n = 7$) and Marcaïn-infused ($n = 6$) rats studied at rest only and in control ($n = 6$) and Marcaïn-infused ($n = 6$) rats studied during a 20-min resting period followed by 20 min of running at a speed of 26 m/min. Marcaïn was infused in ventromedial hypothalamus. Values are means $\pm$ SE. * Difference between control and Marcaïn-infused rats in resting or exercising groups.

TABLE 1. Plasma corticosterone and glucagon concentrations at rest and in 20th min of running in control rats and in rats infused with Marcaïn in ventromedial hypothalamus

	Control		Marcaïn	
	Resting group	Exercising group	Resting group	Exercising group
n	7	6	6	6
Corticosterone ($t = -20$ min)	0.95 ± 0.14	0.89 ± 0.17	0.87 ± 0.33	0.96 ± 0.21
Corticosterone ($t = 20$ min)	0.87 ± 0.14	$1.79 \pm 0.22^*$	$1.35 \pm 0.15^{*\dagger}$	$1.55 \pm 0.15^*$
Glucagon ($t = 20$ min)	20 ± 3	$58 \pm 10^\ddagger$	$37 \pm 2^\ddagger$	$72 \pm 7^\ddagger$

Values are means $\pm$ SE; n , no. of rats. Blood for corticosterone analysis was sampled at rest ($t = -20$ min) in both resting and exercising groups. In exercising groups, exercise was started at $t = 0$ min. Blood for glucagon analysis was only taken at end of experiment due to large plasma volume (0.7 ml) required for determination of this hormone. Plasma corticosterone concentration was measured in $\mu\text{mol/l}$ and glucagon concentration was measured in pmol/l . * Change from $t = -20$ min to $t = 20$ min ($P < 0.05$); $\dagger$ difference between control and Marcaïn-infused rats ($P < 0.05$); $\ddagger$ difference between glucagon levels at rest and during the 20th min of exercise ($P < 0.05$).

have for a long time been recognized as important regulatory mechanisms involved in the control of cardiovascular adjustments to exercise (15). Such neural mechanisms by which working muscles are rapidly supplied with energy-rich substrates are more efficient than sole

regulation of substrate mobilization in exercise by slower acting blood-borne metabolic feedback mechanisms that hitherto has been believed to be the mechanism by which metabolism primarily is regulated during exercise (5, 10).

It may be argued that the effect of recent surgery on

FIG. 3. Plasma concentrations of lactate and glycerol in control ($n = 7$) and Marcaïn-infused ($n = 6$) rats studied at rest only and in control ($n = 6$) and Marcaïn-infused ($n = 6$) rats studied during a 20-min resting period followed by 20 min of running at a speed of 26 m/min. Marcaïn was infused in ventromedial hypothalamus. Values are means $\pm$ SE. * Difference between control and Marcaïn-infused rats in resting or exercising groups.

TABLE 2. Liver and muscle glycogen content in rats infused with Marcaïn in ventromedial hypothalamus and in control rats immediately after resting or running at a speed of 26 m/min for 20 min

	Control		Marcaïn	
	Rest	Exercise	Rest	Exercise
<i>n</i>	7	6	6	6
Liver	250 $\pm$ 47	124 $\pm$ 42*	180 $\pm$ 30	86 $\pm$ 25*
Medial gastrocnemius				
White	42 $\pm$ 3	36 $\pm$ 3	37 $\pm$ 2†	35 $\pm$ 2
Red	40 $\pm$ 2	27 $\pm$ 4*	35 $\pm$ 1†	23 $\pm$ 1*
Soleus	21 $\pm$ 2	6 $\pm$ 1*	15 $\pm$ 1†	8 $\pm$ 2*

Values are means $\pm$ SE; n , no. of rats. Liver and muscle glycogen content were measured in mmol glucose units/kg wet wt. * Difference from resting values ($P < 0.05$); † difference between Marcaïn and control rats ($P < 0.05$).

hormonal and metabolic responses to exercise in the present study may have obscured effects due to exercise per se and thus obscured differences in catecholamine and glucose responses between control and Marcaïn rats. It is well known that rats after insertion of carotid and jugular catheters lose ~ 10 g in body weight the first 24 h after the operation (25, 30). Some influence of surgery on exercise responses can therefore not be completely rejected. However, interaction of surgical aftereffects on exercise responses probably did not play a major role for the following reasons. 1) The body weight had returned to preoperative levels on the day before the experiment, indicating that the rats had recovered well from brain cannulation. 2) The weight loss of 14 ± 1 g in ~ 15 h from the insertion of arterial and venous catheters to the

experiment corresponds to the weight loss in rats that only fast (29). Thus the rats used in the study were not dehydrated. 3) Resting plasma levels of catecholamines and corticosterone and the hormonal and metabolic responses to exercise in rats of the present study were similar compared with levels in rats that were not instrumented or allowed a 1-wk recovery period from arterial and venous operations (25, 29, 30, 33). 4) Running performance in the experiment did not differ from running performance in the days just before insertion of catheters.

In contrast to the smaller exercise-induced increases in hepatic glucose production and plasma concentrations of norepinephrine, epinephrine, and glucose observed in Marcaïn compared with control rats, Marcaïn anesthesia of the VMH at rest elicited a metabolic and endocrine response similar to that observed in exercise. Thus Marcaïn infusion elicited increases in plasma concentrations of norepinephrine, epinephrine, corticosterone, glucose, lactate, and glycerol and an inhibition of plasma insulin levels relative to plasma glucose concentrations. Furthermore, hepatic glucose production and hepatic and muscular glycogenolysis were increased by Marcaïn infusion. These findings could not be explained by increased physical activity in resting Marcaïn rats. It therefore appears that Marcaïn neutralizes activity in areas of the VMH that have an inhibitory influence at rest on the sympathetic nervous system and corticotropes in the anterior pituitary. Interestingly, this means that during exercise the involvement of the VMH in regulation of substrate mobilization may not only be dependent on

increased activity in areas of the VMH that enhance sympathetic activity and pituitary secretion but may also include decreased activity in areas of the VMH that inhibit sympathetic activity and pituitary secretion. In this context, it is of interest to draw a parallel to the well-known phenomenon that the increase in pulse rate during exercise at higher work loads is a result of simultaneous decrease in vagal tone and increase in sympathetic tone to the heart (20).

The VMH is generally recognized as being functionally related to sympathetic facilitation (6, 9, 13, 23, 24, 27) as also suggested by the findings during exercise in this study. Evidence in favor of the view that the VMH also may contain areas that are functionally related to sympathetic inhibition at rest, as suggested by the present study, is furthermore provided by studies where the VMH was lesioned electrolytically. Increases in the sympathetic discharge to the adrenal medulla along with increases in secretion of catecholamines (34), increases in hepatic glycogenolysis (35), and increases in plasma concentrations of corticosterone (35) and glucose (28, 35) have been observed after VMH lesioning in rats. The possibility exists that postlesion edema of close neighboring areas, induced by electrolytic lesions, and, in our study, diffusion of Marcain into close neighboring areas of the VMH may have inactivated a sympathetic inhibitory area just outside the VMH. One of the three rats excluded from the study due to incorrect placement of brain cannulas was a resting Marcain rat with a 1 mm off target placement of cannulas. The plasma glucose was analyzed in this rat to investigate the importance of Marcain diffusing to neighboring areas. Interestingly, plasma glucose concentration did not increase initially after Marcain infusion and only increased 1.0 mmol/l late in the experiment as opposed to a 4.5 ± 0.8 mmol/l increase in rats with correct cannula placements. Thus, within the time frame of the experiment, Marcain does not seem to diffuse to any great extent into neighboring areas of the VMH.

Because exercise-induced elimination of activity in sympathetic inhibitory areas in the VMH may not be a physiological response to exercise, complete anesthesia of both sympathetic inhibitory and facilitating areas in the VMH, induced by Marcain administration, may lead to an underestimation of the importance of sympathetic facilitating areas in the VMH for substrate mobilization during exercise. Accordingly, differences in exercise-induced increases in glucose production and plasma concentrations of epinephrine, norepinephrine, glycerol, and glucose between exercising Marcain and control rats may have been underestimated and differences in plasma corticosterone, insulin, and glucagon levels may have been masked.

Except for a small increase in plasma glycerol after Marcain infusion in resting Marcain rats, fat metabolism, as indicated by plasma glycerol concentrations, was similar in resting Marcain and control rats as well as in the two running groups. It therefore appears that the VMH is not significantly involved in the regulation of fat mobilization during exercise. Muscle metabolism during exercise was similar in Marcain and control rats as

indicated by muscle glycogenolysis, glucose utilization, and plasma concentrations of lactate and glycerol. Similar glucose utilization in the two exercising groups in the face of lower plasma glucose levels in the Marcain rats means that muscular glucose clearance must have been higher in Marcain compared with control rats, a fact that may partly be explained by lower epinephrine levels in Marcain rats (14).

In conclusion, the present study provides evidence for a regulation of glucose mobilization during exercise by the VMH, involving stimulation of sympathoadrenal activity by the VMH. The results are consistent with the view that during exercise, the VMH increases sympathetic activity by increasing activity in areas of the VMH that facilitate sympathetic activity and decreasing activity in areas of the VMH that inhibit sympathetic activity.

Expert technical assistance was performed by Regitze Kraunsøe, Jan Bruggink, Lisbeth Kall, and Vibeke Staffeldt. Excellent secretarial work was done by Janet Wright and Pamela Kay Maass. The advice and support from Dr. Jere H. Mitchell, Univ. of Texas Southwestern Medical Center, Dallas, Texas, is gratefully acknowledged.

This study was supported by the Danish Medical Research Council (12-5995, 12-6420, 12-6352), the Carlsberg Foundation, the NOVO Foundation, the Danish Sports Research Council, and the Foundation for Fundamental Biological Research of 'The Netherlands' Organization for Scientific Research.

Address for reprint requests: J. Vissing, Dept. of Medical Physiology B, The Panum Institute, Univ. of Copenhagen, Blegdamsvej 3C, DK-2200 Copenhagen N, Denmark.

Received 20 September 1988; accepted in final form 29 December 1988.

REFERENCES

1. BAGBY, G. J., H. J. GREEN, S. KATSUTA, AND P. D. GOLLNICK. Glycogen depletion in exercising rats infused with glucose, lactate, or pyruvate. *J. Appl. Physiol.* 45: 425-429, 1978.
2. CHERRINGTON, A. D., R. W. STEVENSON, K. E. STEINER, M. A. DAVIS, S. R. MYERS, B. A. ADKINS, N. N. ABUMRAD, AND P. E. WILLIAMS. Insulin, glucagon and glucose as regulators of hepatic glucose uptake and production in vivo. *Diabetes Metab. Rev.* 3: 307-332, 1987.
3. DUNER, H. The influence of the blood glucose level on the secretion of adrenaline and noradrenaline from the suprarenal. *Acta Physiol. Scand.* 28, Suppl. 102: 1-77, 1953.
4. EGGSTEIN, M., AND F. H. KREUTZ. Eine neue Bestimmung der Neutralfette im Blutserum und Gewebe. I. Prinzip, Durchführung und Besprechung der Methode. *Klin. Wochenschr.* 44: 262-267, 1966.
5. FELIG, P., AND J. WAHREN. Role of insulin and glucagon in the regulation of hepatic glucose production during exercise. *Diabetes* 28, Suppl. 1: 71-75, 1979.
6. FROHMAN, L. A., AND L. L. BERNARDIS. Effect of hypothalamic stimulation on plasma glucose, insulin, and glucagon levels. *Am. J. Physiol.* 221: 1596-1603, 1971.
7. GALBO, H., B. SONNE, J. VISSING, M. KJAER, K. MIKINES, AND E. A. RICHTER. Lack of accuracy of fuel mobilization in exercise. In: *Biochemical Aspects of Physical Exercise*, edited by G. Benzi, L. Packer, and N. Siliprandi. Amsterdam: Elsevier, 1986, p. 221-234.
8. GALBO, H. *Hormonal and Metabolic Adaptation to Exercise*. Stuttgart, FRG: Thieme-Stratton, 1983, p. 1-116.
9. INOUE, S., AND G. A. BRAY. Role of the autonomic nervous system in the development of ventromedial hypothalamic obesity. *Brain Res. Bull.* 5, Suppl. 4: 119-125, 1980.
10. JENKINS, A. B., S. M. FURLER, D. J. CHISHOLM, AND E. W. KRAEGEN. Regulation of hepatic glucose output during exercise by circulating glucose and insulin in humans. *Am. J. Physiol.* 250 (Regulatory Integrative Comp. Physiol. 19): R411-R417, 1986.
11. KJAER, M., N. H. SECHER, F. W. BACH, AND H. GALBO. Role of motor center activity for hormonal changes and substrate mobilization in humans. *Am. J. Physiol.* 253 (Regulatory Integrative

- Comp. Physiol.* 22): R687–R695, 1987.
12. KUMON, A., A. TAKAHASHI, T. HARA, AND T. SHIMAZU. Mechanism of lipolysis induced by electrical stimulation of the hypothalamus in the rabbit. *J. Lipid Res.* 17: 551–558, 1976.
 13. LUITEN, P. G. M., G. J. TER HORST, AND A. B. STEFFENS. The hypothalamus, intrinsic connections and outflow pathways to the endocrine system in relation to the control of feeding and metabolism. *Prog. Neurobiol.* 28: 1–54, 1987.
 14. MIKINES, K. J., B. SONNE, E. A. RICHTER, N. J. CHRISTENSEN, AND H. GALBO. Glucose turnover during insulin-induced hypoglycemia in liver-denervated rats. *Am. J. Physiol.* 248 (*Endocrinol. Metab.* 11): E327–E332, 1985.
 15. MITCHELL, J. H. Cardiovascular control during exercise: central and reflex neural mechanisms. *Am. J. Cardiol.* 55: 34D–41D, 1985.
 16. MITCHELL, J. H., J. VISSING, K. J. RYBICKI, H. GALBO, AND G. A. IWAMOTO. Parallel activation of locomotion, metabolism and circulation from the hypothalamus (Abstract). *FASEB J.* 2: A1325, 1988.
 17. OLSEN, C. An enzymatic fluorimetric micromethod for the determination of acetoacetate, beta-hydroxybutyrate, pyruvate and lactate. *Clin. Chim. Acta* 33: 293–300, 1971.
 18. PAXINOS, G., AND C. WATSON. *The Rat Brain in Stereotaxic Coordinates*. New York: Academic, 1982.
 19. PROIETTO, J., F. ROHNER-JEANRENAUD, E. IONESCU, J. TERRETTAZ, J. F. SAUTER, AND B. JEANRENAUD. Non-steady-state measurement of glucose turnover in rats by using a one-compartment model. *Am. J. Physiol.* 252 (*Endocrinol. Metab.* 15): E77–E84, 1987.
 20. ROWELL, L. *Human Circulation. Regulation During Physical Stress*. New York: Oxford Univ. Press, 1986, p. 1–416.
 21. SCHEURINK, A. J. W., A. B. STEFFENS, H. BOURITIUS, G. DRETELER, R. BRUNTINK, R. REMIE, AND J. ZAAGSMA. Adrenal and sympathetic catecholamines in exercising rats. *Am. J. Physiol.* 256 (*Regulatory Integrative Comp. Physiol.* 25): R155–R160, 1989.
 22. SCHEURINK, A. J. W., A. B. STEFFENS, AND L. BENTHEM. Central and peripheral adrenoreceptors affect glucose, FFA and insulin in exercising rats. *Am. J. Physiol.* 255 (*Regulatory Integrative Comp. Physiol.* 24): R547–R556, 1988.
 23. SHIMAZU, T., AND K. ISHIKAWA. Modulation by the hypothalamus of glucagon and insulin secretion in rabbits: studies with electrical and chemical stimulations. *Endocrinology* 108: 605–611, 1981.
 24. SHIMAZU, T. Neuronal control of intermediate metabolism. In: *Neuroendocrinology*, edited by S. Lightman and B. Everitt. Oxford, UK: Blackwell, 1986, p. 304–330.
 25. SONNE, B., K. J. MIKINES, E. A. RICHTER, N. J. CHRISTENSEN, AND H. GALBO. Role of liver nerves and adrenal medulla in glucose turnover of running rats. *J. Appl. Physiol.* 59: 1640–1646, 1985.
 26. STEELE, R. Influences of glucose loading and of injected insulin on hepatic glucose output. *Ann. NY Acad. Sci.* 82: 420–430, 1959.
 27. STEFFENS, A. B., G. DAMSMA, J. VAN DER GUGTEN, AND P. G. M. LUITEN. Circulating free fatty acids, insulin, and glucose during chemical stimulation of hypothalamus in rats. *Am. J. Physiol.* 247 (*Endocrinol. Metab.* 10): E765–E771, 1984.
 28. STEFFENS, A. B., G. J. MOGENSEN, AND J. A. F. STEVENSON. Blood glucose, insulin, and free fatty acids after stimulation and lesions of the hypothalamus. *Am. J. Physiol.* 222: 1446–1452, 1972.
 29. VISSING, J., J. L. WALLACE, AND H. GALBO. Effect of liver glycogen content on glucose production in running rats. *J. Appl. Physiol.* 66: 318–322, 1989.
 30. VISSING, J., B. SONNE, AND H. GALBO. Role of metabolic feedback regulation in glucose production of running rats. *Am. J. Physiol.* 255 (*Regulatory Integrative Comp. Physiol.* 24): R400–R406, 1988.
 31. VISSING, J., T. OHKUWA, T. PLOUG, AND H. GALBO. Effect of prior immobilization on muscular glucose clearance in resting and running rats. *Am. J. Physiol.* 255 (*Endocrinol. Metab.* 18): E456–E462, 1988.
 32. WINDER, W. W. Regulation of hepatic glucose production during exercise. *Exer. Sports Sci. Rev.* 13: 1–31, 1985.
 33. YANG, H. T., R. L. HAMMER, T. L. SELLERS, J. AROGYASAMI, D. T. CARRELL, AND W. W. WINDER. Adrenomedullation affects endurance but not hepatic fructose 2,6-bisphosphate. *Am. J. Physiol.* 254 (*Regulatory Integrative Comp. Physiol.* 23): R572–R577, 1988.
 34. YOSHIMATSU, H., Y. OOMURA, T. KATAFUCHI, A. NIJIMA, AND A. SATO. Lesions of the ventromedial hypothalamic nucleus enhance sympatho-adrenal function. *Brain Res.* 339: 390–392, 1985.
 35. ZAIA, T. B. V., C. M. P. OLLER DO NASCIMENTO, C. TIMO-IARIA, AND M. S. DOLNIKOFF. Time course of insulin, corticosterone and metabolic changes caused by lesion of the ventromedial hypothalamus in the rat. *Physiol. Behav.* 39: 707–714, 1987.