University of Groningen

Hydrolysis and Synthesis of ATP by Membrane-Bound ATPase from a Motile Streptococcus

Drift, C. van der; Janssen, D.B.; Wezenbeek, P.M.G.F. van

Published in:
Archives of Microbiology

DOI:
[10.1007/BF00407924](https://doi.org/10.1007/BF00407924)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1978

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Drift, C. V. D., Janssen, D. B., & Wezenbeek, P. M. G. F. V. (1978). Hydrolysis and Synthesis of ATP by Membrane-Bound ATPase from a Motile Streptococcus. *Archives of Microbiology*, 119(1).
<https://doi.org/10.1007/BF00407924>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Hydrolysis and Synthesis of ATP by Membrane-Bound ATPase from a Motile *Streptococcus*

C. van der Drift, D. B. Janssen, and P.M.G.F. van Wezenbeek

Department of Microbiology, Faculty of Science, University of Nijmegen, Nijmegen, The Netherlands

Abstract. ATPase was detected in the membranes of a motile *Streptococcus*. Maximal enzymic activity was observed at pH 8 and ATP/Mg²⁺ ratio of 2. Mn²⁺ and Ca²⁺ could replace Mg²⁺ to some extent. Besides ATP, GTP and ITP were substrates. The enzyme was inhibited by N,N'-dicyclohexylcarbodiimide but not by sodium azide, uncouplers or bathophenanthroline.

An electrochemical gradient of protons, which was artificially imposed across the membranes of *Streptococcus* cells by manipulation of either the K⁺ diffusion potential or the transmembrane pH gradient, led to ATP synthesis. ATP synthesis was abolished by proton conductors, an inhibitor of the ATPase or an increase in the extracellular K⁺ concentration. A comparison between the phosphate potential and the electrochemical proton gradient showed that the data found are in agreement with a stoichiometry of 2 protons translocated per molecule ATP synthesized.

Key words: Motile *Streptococcus* – Membrane-bound ATPase – DCCD – Valinomycin – Uncouplers – Protonmotive force – ATP synthesis.

Energization of various membrane-bound processes, such as active transport of neutral and ionic solutes and ATP synthesis via the membrane-bound proton-translocating ATPase (adenosine phosphohydrolase, EC 3.6.1.3), is brought about by chemiosmotic cou-

pling (Mitchell, 1966; Harold, 1972; Hamilton, 1975; Haddock and Jones, 1977). In aerobic and facultatively anaerobic organisms the driving force for these processes – the so-called protonmotive force or electrochemical proton gradient ($\Delta\mu_{H^+}$) – is generated by the redox reactions of the electron transport chain. Anaerobic fermentative and facultatively anaerobic organisms in the absence of oxygen generate a $\Delta\mu_{H^+}$ via an electrogenic proton-translocating ATPase. The $\Delta\mu_{H^+}$ is composed of an electrical and a chemical parameter according to the relationship $\Delta\mu_{H^+} = \Delta\psi - Z\Delta pH$, where $\Delta\psi$ represents the membrane potential and ΔpH the transmembrane pH difference. Z is equal to 2.3 RT/F and amounts to 59 mV at 25° C.

Recently it was shown that an electrochemical gradient of protons is involved in motility and chemotaxis. Proton circulation not ATP delivers the energy necessary to drive the flagellar motor (Larsen et al., 1974; Manson et al., 1977) and the transmittance of a signal from a chemoreceptor to the flagellar motor is thought to occur via fluctuations in the $\Delta\mu_{H^+}$ or in the flux of a particular ion (de Jong et al., 1976; Szmecman and Adler, 1976; Manson et al., 1977; Matsuura et al., 1977; de Jong and van der Drift, 1978). Previously the motile and chemotactic behavior of a motile *Streptococcus* was studied (van der Drift et al., 1975; Manson et al., 1977). Since this *Streptococcus* is a primarily fermentative organism, $\Delta\mu_{H^+}$ will be generated mainly via the membrane-bound ATPase. In this study we report on the properties of the membrane-bound ATPase of this motile *Streptococcus*. Both hydrolysis of ATP and synthesis of ATP under influence of an artificially imposed $\Delta\mu_{H^+}$ was studied. ATP synthesis in bacterial systems by means of an artificially imposed $\Delta\mu_{H^+}$ previously was shown for whole cells of *Escherichia coli* (Grinius et al., 1975), *Streptococcus lactis* (Maloney and Wilson, 1975) and *Halobacterium halobium* (Danon and Caplan, 1976), membrane vesicles of *E. coli* (Tsuchiya and Rosen,

Abbreviations. $\Delta\mu_{H^+}$ = electrochemical gradient of protons; DMO = 5,5-dimethyl-2,4-oxazolidinedione; CCCP = carbonylcyanide *m*-chlorophenylhydrazine; FCCP = carbonylcyanide *p*-trifluoromethoxyphenylhydrazine; DCCD = N,N'-dicyclohexylcarbodiimide; DNP = 2,4-dinitrophenol

1976) and a thermophilic bacterium (Sone et al., 1977), and chromatophores of *Rhodospirillum rubrum* (Gromet-Elhanan and Leiser, 1975).

Materials and Methods

Microorganism. The motile *Streptococcus* Strain V4051 (van der Drift et al., 1975) was used in all experiments.

Media, Growth Conditions and Preparation of Cell Suspensions. The strain was grown at 37° C in 3.7% brain heart infusion broth (Oxoid) supplemented with 0.5% KCl. Cells were harvested by centrifugation (8,000 × g for 10 min at 4° C) in the early stationary growth phase at a cell density of 0.5 mg of dry weight per ml. The cells were washed twice with 100 mM sodium phosphate buffer (pH 7.0) and resuspended in this buffer to 0.4–0.6 mg of dry weight per ml.

Preparation of Membranes. Cells were harvested by centrifugation and washed twice with 100 mM Tris-HCl buffer (pH 8.0) containing 2.5 mM MgCl₂. The cells were resuspended in this buffer (5 ml buffer per g wet weight) and lysed by passage through a French pressure cell (American Instrument Co.) at 140,000 kPa. The suspension was centrifuged at 20,000 × g for 10 min and the supernatant solution was centrifuged at 100,000 × g for 1 h at 4° C. The membranes were washed and resuspended in Tris-Mg²⁺ buffer (pH 8.0) (about 5 mg of protein per ml), frozen in liquid nitrogen and stored at –20° C.

Assays. The ATPase activity of the isolated membranes was assayed by measurement of inorganic phosphate (Bonting et al., 1961) liberated from ATP at 37° C. The incubation mixture routinely contained 100 mM Tris-HCl (pH 8.0), 5 mM ATP, 2.5 mM MgCl₂ and a suitable amount of membrane protein (200–600 µg per ml). One unit of activity is defined as the amount of enzyme which liberates 1 µmol inorganic phosphate per min at pH 8.0 and 37° C. Specific activity is expressed as units per mg of protein.

Intracellular ATP was extracted as described by Maloney and Wilson (1975). The extracted ATP was measured according to Cole et al. (1967). ADP was measured after its conversion to ATP in the presence of pyruvate kinase and excess phosphoenolpyruvate (Chapman et al., 1971). Intracellular inorganic phosphate was extracted by treatment of the cells for 15 min at 0° C with 1 N HClO₄ (10 mg of dry weight per ml HClO₄). After neutralization with KOH and removal of cells and precipitate by centrifugation inorganic phosphate was measured (Bonting et al., 1961).

Intracellular potassium was determined by flame photometry (Maloney and Wilson, 1975).

The intracellular pH was calculated from the distribution of the weak acid [¹⁴C]-5,5-dimethyl-2,4-oxazolidinedione (DMO) between the intracellular space of the cells and the medium using [³H]-inulin as marker for extracellular water (Rottenberg, 1975).

Cell water was determined with the use of ³H₂O and [¹⁴C]-inulin assuming that water freely permeates the cell membrane, whereas inulin does not.

Protein was measured according to Lowry et al. (1951) using bovine serum albumin as standard.

Proton Conduction. The rate of H⁺ entry at 24° C was measured according to Harold and Baarda (1968).

ATP Synthesis. Washed cells were resuspended (0.4–0.6 mg dry weight per ml) in 100 mM sodium phosphate buffer of the desired pH. An artificial Δμ_{H⁺} was applied at 24° C either by addition of the potassium ionophore valinomycin, a pH shift or a combination of both. At different time intervals samples were taken and assayed for ATP content. The size of the Δμ_{H⁺} was calculated from the contributions made by both the membrane potential and the pH gradient. The membrane potential was determined from the measured ratio of internal to external potassium using the Nernst equation. The activity coefficient for K⁺ was assumed to be 1 both for

the internal space and the extracellular medium. The pH gradient was determined from measurements of internal and external pH and expressed in terms of millivolts (59 ΔpH).

The phosphate potential — the free energy of synthesis of ATP — was calculated from the intracellular concentrations of ATP, ADP and inorganic phosphate and using a value of 28.5 kJ/mol for the standard free energy of hydrolysis of ATP at pH 7.0, 10 mM Mg²⁺, 0.2 ionic strength and 25° C (Rosing and Slater, 1972). Since the intracellular inorganic phosphate concentration (37 mM) was much higher than the nucleotide concentration, changes in this concentration were neglected.

Chemicals. ATP, ADP, ITP, GTP and carbonylcyanide *p*-trifluoromethoxyphenylhydrazone (FCCP) were obtained from Boehringer, Mannheim; valinomycin and carbonylcyanide *m*-chlorophenylhydrazone (CCCP) from Calbiochem, San Diego, CA; N,N'-dicyclohexylcarbodiimide (DCCD) from British Drug Houses, Poole; bathophenanthroline from Merck, Darmstadt; firefly lantern extract (FLE-50) from Sigma, St. Louis, MI; [³H]-inulin from The Radio Chemical Centre, Amersham; and [¹⁴C]-5,5-dimethyl-2,4-oxazolidinedione (DMO) and ³H₂O from New England Nuclear Corp., Boston, MA. Solutions of valinomycin, CCCP and DCCD were made in methanol.

Results

Proton Conduction

Cells took up protons slowly when the pH of a lightly buffered cell suspension was lowered from 6.5 to 5.3. Addition of valinomycin resulted in an increase of H⁺ uptake and subsequent addition of either CCCP or DNP permitted a very rapid entry of H⁺ into the cells (Fig. 1). Pretreatment of the cells for 15 min at 24° C with DCCD (10^{–4}M) resulted after acidification of the cell suspension in a H⁺ uptake which was about half that of untreated cells. After addition of valinomycin to DCCD-treated cells a light increase in H⁺ uptake was observed, while addition of uncoupler resulted in a very rapid H⁺ uptake (Fig. 1).

Hydrolysis of ATP

Cells of *Streptococcus* strain V4051 were fairly resistant to lysozyme even after acetylation of the cell wall as described by Eisenberg and Lillmars (1975). Therefore membranes were prepared by using a French pressure cell. The specific ATPase activity of such membranes amounted to 0.3. Membranes with a similar specific activity were obtained if the cells were broken by ultrasonic disintegration. The streptococcal membrane-bound ATPase showed optimal activity at pH 8.0. The enzyme required Mg²⁺ for activity and the reaction velocity was maximal at an ATP/Mg²⁺ ratio of 2. Mg²⁺ could be partially replaced by Mn²⁺ and Ca²⁺: in the case 80% and 35%, respectively, of the activity with Mg²⁺ was measured. The enzyme exhibited Michaelis-Menten kinetics with a K_m value for ATP of 1.8 mM. GTP, ITP and ADP were hydrolyzed by the enzyme at a rate which was 100%, 70% and less

Fig. 1. Proton uptake by *Streptococcus* strain V4051. Cells were suspended (0.21 mg dry weight/ml) at 24°C in KCl (50 mM)-MgCl₂ (2 mM)-glycylglycine (1 mM). To 10-ml portions (initial pH 6.5) 0.05 ml 10 mM HCl was added followed by further additions as indicated by the arrows. The dashed line represents cells treated for 15 min at 24°C with DCCD (10⁻⁴ M) before the addition of HCl. Concentrations used were: valinomycin (val), 10⁻⁶ M; CCCP, 10⁻⁵ M; DNP, 5 × 10⁻⁴ M

Fig. 2a and b. ATP synthesis driven by a pH gradient. Cells (0.53 mg dry weight/ml) were suspended in 100 mM sodium phosphate (pH 8.0). A pH gradient was applied by addition of various amounts of HCl. ATP levels were determined as described in "Materials and Methods". **a** Time-dependent ATP synthesis after a pH shift from 8 to 3.5. ●—● none addition; ■—■ cells preincubated for 15 min with 10⁻⁴ M DCCD; △—△ cells preincubated for 1 min with 10⁻⁵ M CCCP. **b** ATP synthesis as function of the pH gradient. The maximal level of ATP was determined for all shifts applied in external pH. All experiments were carried out at 24°C

Fig. 3a and b. ATP synthesis driven by a combination of a membrane potential and a pH gradient. Cells (0.5 mg dry weight/ml) were suspended in 100 mM sodium phosphate buffer (pH 7.0). An $\Delta\mu_{H^+}$ was imposed by addition of HCl, valinomycin or a combination of both. ATP levels were determined as described in "Materials and Methods". **a** Time-dependent ATP synthesis. △—△ valinomycin (10⁻⁵ M); ●—● the external pH was shifted from 7 to 5.2; ■—■ the external pH was shifted to 5.2 and valinomycin (10⁻⁵ M) was added. **b** Effect of external K⁺ on ATP synthesis. ■—■ both the external pH was shifted to 5.2 and valinomycin (10⁻⁵ M) was added, external K⁺ 0.3 mM; △—△ external K⁺ 1.3 mM; ●—● external K⁺ 10.3 mM

than 2%, respectively, of that observed with ATP as substrate. The enzyme was inhibited by DCCD: preincubation for 15 min at 24°C with DCCD (2 × 10⁻⁴ M) lowered the ATPase activity about 70%. No inhibition was observed with sodium azide (100 mM), CCCP (0.05 mM), FCCP (0.01 mM), DNP (1 mM) and bathophenanthroline (1 mM). Preincubation of the membrane-bound ATPase for 10 min at 37°C with trypsin (0.5 mg/ml) lowered the activity to 75%.

No effective and selective solubilization of the membrane-bound enzyme was obtained when the membranes were either washed with buffers of low ionic strength in the absence of Mg²⁺ (Abrams, 1965) or treated with NH₄HCO₃ and EDTA at pH 9 (Carreira et al., 1973).

ATP Synthesis Driven by a pH Gradient

The ATP level in nonstarved and starved cells of *Streptococcus* strain V4051 is about the same (about 100 pmol/mg dry weight). The intracellular pH of the cells was 6.8 as measured by [¹⁴C]-DMO distribution. This value is in agreement with that determined by Manson et al. (1977) after lysis of the cells in *n*-butanol. The establishment of an inwardly directed $\Delta\mu_{H^+}$ by imposing a pH gradient (alkaline inside) across the cell membrane resulted in ATP synthesis. At pH 3.5 the

ATP level increased maximally about 40-fold. No ATP synthesis after a pH jump was observed if CCCP, a proton conductor, or DCCD, an ATPase inhibitor, was present (Fig. 2a). The magnitude of the applied pH gradient determined the maximal increase in the ATP level. A transition from pH 8 to pH 3.5 was optimal for ATP synthesis, while a shift from pH 8 to pH 5 did not result in ATP formation (Fig. 2b).

ATP synthesis was dependent on the growth phase of the cells. It appeared that maximal ATP synthesis occurred in cells from the early stationary phase. Therefore such cells were used in all experiments.

ATP Synthesis Driven by a Membrane Potential and a pH Gradient

Cells harvested in the early stationary phase have a K⁺ content of 600 mM. The K⁺ content of washed cells used in our experiments was 220 mM. Addition of valinomycin (10⁻⁵ M) to cells suspended in potassium-free sodium phosphate buffer (pH 7.0) resulted in the synthesis of a very small amount of ATP (see also Manson et al., 1977). If both an artificially imposed membrane potential and pH gradient were applied a considerable amount of ATP was synthesized; the pH gradient by itself was insufficient to induce ATP synthesis (Fig. 3). ATP synthesis was abolished by

Table 1. Correlation between ATP synthesis and the electrochemical proton gradient. Cells (0.5 mg dry weight/ml) were suspended in 100 mM sodium phosphate buffer (pH 6.8) with different concentrations of K^+ . An $\Delta\mu_{H^+}$ was applied by addition of valinomycin ($10^{-5}M$) and/or HCl. ATP synthesis was measured as described in "Materials and Methods". The size of the $\Delta\mu_{H^+}$ was calculated from the contributions of the pH gradient and the membrane potential

Maximal ATP synthesis (nmol/mg dry weight)	$\Delta\psi$ (mV)	$-Z\Delta pH$ (mV)	$\Delta\mu_{H^+}$ (mV)
0.1	104	0	104
0.5	137	8	141
0.7	104	67	171
0.9	104	72	176
0.9	137	40	177
1.0	137	41	178
1.0	104	76	180
1.4	137	60	197
1.5	137	66	203
1.3	104	168	272
1.2	137	204	341

either CCCP or DCCD (data not shown). Addition of KCl to the external medium reduced ATP synthesis as would be expected since the driving force for ATP synthesis was mainly composed of a K^+ diffusion potential.

Relation between ATP Synthesis and Electrochemical Proton Gradient

Table 1 shows the correlation between ATP synthesis and the applied $\Delta\mu_{H^+}$. At all values of $\Delta\mu_{H^+}$ net ATP synthesis was observed. Maximal synthesis seemed to be reached at about 200 mV, while below 140 mV only a slight increase in ATP levels was observed.

Comparison of Phosphate Potential and Electrochemical Proton Gradient

Table 2 shows that the phosphate potential measured under the experimental conditions used would require an $\Delta\mu_{H^+}$ of about 200 mV if a stoichiometry of 2 protons translocated per ATP synthesized is assumed. The values of $\Delta\mu_{H^+}$, which were measured simultaneously, were in reasonable agreement with the phosphate potential based on the above-mentioned assumption.

Discussion

The properties of the membrane-bound ATPase of *Streptococcus* strain V4051 resemble those of other membrane-bound bacterial ATPases (Abrams and Smith, 1974; Panet and Sanadi, 1976). The enzyme

Table 2. Correlation between phosphate potential and electrochemical proton gradient. Experimental conditions are given in Table 1. The phosphate potential and the $\Delta\mu_{H^+}$ were measured as described in "Material and Methods"

Phosphate potential		$\Delta\mu_{H^+}$ (mV)	
ATP ADP	$\Delta G'$ (kJ/mol)	$\Delta\mu_{H^+}$ required for $\Delta G'$ assuming $2H^+/ATP$	$\Delta\mu_{H^+}$ applied
0.3	33.7	174	161
1.6	37.8	196	176
2.5	38.9	202	197
2.7	39.1	202	187
3.1	39.5	205	187
4.7	40.5	209	176
6.3	41.2	213	198

shows maximal activity at pH 8.0 and a ATP/Mg^{2+} ratio of 2. Mg^{2+} can be partially replaced by Mn^{2+} and Ca^{2+} ions. Besides ATP, GTP and ITP are substrates. The enzyme is inhibited by DCCD, but not by sodium azide, CCCP, FCCP and DNP. Bathophenanthroline, a lipophilic chelator, which inhibits the *Escherichia coli* ATPase (Sun et al., 1975) has no effect on the streptococcal enzyme. Trypsin, which stimulates the ATPase activity of various bacteria (Carreira et al., 1973; Panet and Sanadi, 1976) does not activate the streptococcal ATPase, but inactivates it just as was observed with the enzyme from *Clostridium pasteurianum* (Riebeling and Jungermann, 1976). The streptococcal ATPase remains membrane-bound when washing procedures are employed using dilute buffers in the absence of Mg^{2+} ions (Abrams, 1965; Carreira et al., 1973; Abrams and Smith, 1974).

An artificially imposed electrochemical gradient of protons either brought about by a pH gradient or a pH gradient in combination with valinomycin results in a transient synthesis of ATP. This synthesis does not take place in the presence of a proton conductor or an inhibitor of the ATPase and furthermore is dependent on the external K^+ concentration. Cells pretreated with DCCD show a much lower proton uptake than untreated cells indicating the role of the ATPase in proton translocation. These observations can be fully explained by the chemiosmotic hypothesis (Mitchell, 1966; Harold, 1972) and are in agreement with observations made with other bacteria (Grinius et al., 1975; Maloney and Wilson, 1975; Danon and Caplan, 1976).

It was proposed that 2 protons are translocated per ATP molecule hydrolyzed or synthesized (Moyle and Mitchell, 1973). A comparison of the $\Delta\mu_{H^+}$ values generated artificially across the cell membrane of *Streptococcus* strain V4051 with the values of the

phosphate potential assuming a H^+ /ATP ratio of 2 shows that the values are in reasonable agreement. A translocation of 3 protons per ATP synthesized was observed with mitochondria (Nicholls, 1974), chloroplasts (Avron, 1976) and chromatophores of *Rhodospirillum rubrum* (Leiser and Gromet-Elhanan, 1977). However, in chromatophores of *Rhodopseudomonas capsulata* the measured $\Delta\mu_{H^+}$ values were identical to or in excess of those required by the measured phosphate potential assuming a H^+ /ATP ratio of 2 (Casadio et al., 1974). Recently it was proposed that in mitochondria 3 protons are ejected per pair of electrons traversing each energy-conserving site of the respiratory chain and that 2 protons return to the cytoplasm through the ATPase to form ATP from ADP and phosphate or alternatively that 4 protons are ejected per site followed by return of 3 protons through the ATPase (Brand and Lehninger, 1977).

Our data indicate that in *Streptococcus* strain V4051 the H^+ /ATP ratio is 2. However, this ratio may vary with different organisms and further work will be needed to elucidate this.

Acknowledgement. We wish to thank Dr. F. M. Harold for critical reading of the manuscript.

References

- Abrams, A.: The release of bound adenosine triphosphatase from isolated bacterial membranes and the properties of solubilized enzyme. *J. Biol. Chem.* **240**, 3675–3681 (1965)
- Abrams, A., Smith, J. B.: Bacterial ATPase. In: *The enzymes*, Vol. X, 3rd ed. (P. D. Boyer, ed.), pp. 395–429. New York-London: Academic Press 1974
- Avron, M.: Energy transduction in isolated chloroplast membranes. In: *The structural basis of membrane function* (Y. Hatefi, L. Djavadi-Ohanian, eds.), pp. 227–238. New York-London: Academic Press 1976
- Bonting, S. L., Simon, K. A., Hawkins, N. M.: Studies on sodium-potassium-activated adenosine triphosphatase. I. Quantitative distribution in several tissues of the cat. *Arch. Biochem. Biophys.* **95**, 416–423 (1961)
- Brand, M. D., Lehninger, A. L.: H^+ /ATP ratio during ATP hydrolysis by mitochondria: modification of the chemiosmotic theory. *Proc. Nat. Acad. Sci. U.S.A.* **74**, 1955–1959 (1977)
- Carreira, J., Leal, J. A., Rojas, M., Munoz, E.: Membrane ATPase of *Escherichia coli* K12. Selective solubilization of the enzyme and its stimulation by trypsin in the soluble and membrane-bound states. *Biochim. Biophys. Acta* **307**, 541–556 (1973)
- Casadio, R., Baccarini-Melandri, A., Zannoni, D., Melandri, B. A.: Electrochemical proton gradient and phosphate potential in bacterial chromatophores. *FEBS Lett.* **49**, 203–207 (1974)
- Chapman, A. G., Fall, L., Atkinson, D. E.: Adenylate energy charge in *Escherichia coli* during growth and starvation. *J. Bacteriol.* **108**, 1072–1086 (1971)
- Cole, H. A., Wimpenny, J. W. T., Hughes, D. E.: The ATP pool in *Escherichia coli*. I. Measurement of the pool using a modified luciferase assay. *Biochim. Biophys. Acta* **143**, 445–453 (1967)
- Danon, A., Caplan, S. R.: Stimulation of ATP synthesis in *Halobacterium halobium* R_1 by light-induced or artificially created proton electrochemical potential gradients across the cell membrane. *Biochim. Biophys. Acta* **423**, 133–140 (1976)
- de Jong, M. H., van der Drift, C.: Control of the chemotactic behavior of *Bacillus subtilis* cells. *Arch. Microbiol.* **116**, 1–8 (1978)
- de Jong, M. H., van der Drift, C., Vogels, G. D.: Proton-motive force and the motile behavior of *Bacillus subtilis*. *Arch. Microbiol.* **111**, 7–11 (1976)
- Eisenberg, R. J., Lillmars, K.: A method for gentle lysis of *Streptococcus sanguis* and *Streptococcus mutans*. *Biochem. Biophys. Res. Commun.* **65**, 378–384 (1975)
- Grinius, L., Slušnyte, R., Griniuvienė, B.: ATP synthesis driven by protonmotive force imposed across *Escherichia coli* cell membranes. *FEBS Lett.* **57**, 290–293 (1975)
- Gromet-Elhanan, Z., Leiser, M.: Postillumination adenosine triphosphate synthesis in *Rhodospirillum rubrum* chromatophores. II. Stimulation by a K^+ diffusion potential. *J. Biol. Chem.* **250**, 90–93 (1975)
- Haddock, B. A., Jones, C. W.: Bacterial respiration. *Bacteriol. Rev.* **41**, 47–99 (1977)
- Hamilton, W. A.: Energy coupling in microbial transport. *Adv. Microb. Physiol.* **12**, 1–53 (1975)
- Harold, F. M.: Conservation and transformation of energy by bacterial membranes. *Bacteriol. Rev.* **36**, 172–230 (1972)
- Harold, F. M., Baarda, J. R.: Inhibition of membrane transport in *Streptococcus faecalis* by uncouplers of oxidative phosphorylation and its relationship to proton conduction. *J. Bacteriol.* **96**, 2025–2034 (1968)
- Larsen, S. H., Adler, J., Gargus, J. J., Hogg, R. W.: Chemo-mechanical coupling without ATP. The source of energy for motility and chemotaxis in bacteria. *Proc. Nat. Acad. Sci. U.S.A.* **71**, 1239–1243 (1974)
- Leiser, M., Gromet-Elhanan, Z.: Comparison of the electrochemical proton gradient and phosphate potential maintained by *Rhodospirillum rubrum* chromatophores in the steady state. *Arch. Biochem. Biophys.* **178**, 79–88 (1977)
- Lowry, O. H., Rosebrough, N. J., Farr, A. L., Randall, R. J.: Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**, 265–275 (1951)
- Maloney, P. C., Wilson, T. H.: ATP synthesis driven by a protonmotive force in *Streptococcus lactis*. *J. Membrane Biol.* **25**, 285–310 (1975)
- Manson, M. D., Tedesco, P., Berg, H. C., Harold, F. M., van der Drift, C.: A protonmotive force drives bacterial flagella. *Proc. Nat. Acad. Sci. U.S.A.* **74**, 3060–3064 (1977)
- Matsuura, S., Shioi, J.-I., Imae, Y.: Motility in *Bacillus subtilis* driven by an artificial proton motive force. *FEBS Lett.* **82**, 187–190 (1977)
- Mitchell, P.: Chemiosmotic coupling in oxidative and photosynthetic phosphorylation. *Biol. Rev.* **41**, 445–502 (1966)
- Moyle, J., Mitchell, P.: Proton translocation quotient for the adenosine triphosphatase of rat liver mitochondria. *FEBS Lett.* **30**, 317–320 (1973)
- Nicholls, D. G.: The influence of respiration and ATP hydrolysis on the proton electrochemical gradient across the inner membrane of rat liver mitochondria as determined by ion distribution. *Eur. J. Biochem.* **50**, 305–315 (1974)
- Panet, R., Sanadi, D. R.: Soluble and membrane ATPase of mitochondria, chloroplasts and bacteria: molecular structure, enzymatic properties, and functions. *Curr. Top. Membr. Transp.* **8**, 99–160 (1976)
- Riebeling, V., Jungermann, K.: Properties and function of clostridial membrane ATPase. *Biochim. Biophys. Acta* **430**, 434–444 (1976)
- Rosing, J., Slater, E. C.: The value of ΔG° for the hydrolysis of ATP. *Biochim. Biophys. Acta* **267**, 275–290 (1972)

- Rottenberg, H.: The measurement of transmembrane electrochemical proton gradients. *J. Bioenerg.* **7**, 61–74 (1975)
- Sone, N., Yoshida, M., Hirata, H., Kagawa, Y.: Adenosine triphosphate synthesis by electrochemical proton gradient in vesicles reconstituted from purified adenosine triphosphatase and phospholipids of thermophilic bacterium. *J. Biol. Chem.* **252**, 2956–2960 (1977)
- Sun, I. L., Phelps, D. C., Crane, F. L.: Lipophilic chelator inhibition of *Escherichia coli* membrane-bound ATPase activity and prevention of inhibition by uncouplers. *FEBS Lett.* **54**, 253–258 (1975)
- Szmelcman, S., Adler, J.: Change in membrane potential during bacterial chemotaxis. *Proc. Nat. Acad. Sci. U.S.A.* **73**, 4387–4391 (1976)
- Tsuchiya, T., Rosen, B. P.: ATP synthesis by an artificial proton gradient in right-side-out membrane vesicles of *Escherichia coli*. *Biochem. Biophys. Res. Commun.* **68**, 497–502 (1976)
- van der Drift, C., Duiverman, J., Bexkens, H., Krijnen, A.: Chemotaxis of a motile *Streptococcus* toward sugars and amino acids. *J. Bacteriol.* **124**, 1142–1147 (1975)

Received April 26, 1978