University of Groningen

Torsionally Distorted Olefins. Resolution of cis- and trans-4,4'-Bi-1,1',2,2',3,3'-hexahydrophenanthrylidene

Wynberg, Hans; Feringa, Bernard

Published in:
Journal of the American Chemical Society

DOI:
[10.1021/ja00444a046](https://doi.org/10.1021/ja00444a046)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1977

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Wynberg, H., & Feringa, B. (1977). Torsionally Distorted Olefins. Resolution of cis- and trans-4,4'-Bi-1,1',2,2',3,3'-hexahydrophenanthrylidene. *Journal of the American Chemical Society*, 99(2).
<https://doi.org/10.1021/ja00444a046>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Communications to the Editor

Torsionally Distorted Olefins. Resolution of *cis*- and *trans*-4,4'-Bi-1,1',2,2',3,3'-hexahydrophenanthrylidene

Sir:

Only few examples are known of optically active stereoisomers of alkenes bearing no chiral centers.¹ In principle both the *cis* as well as the *trans* form of olefins should be capable of existence as stable optically active stereoisomers, provided bulky substituents cause a permanent out-of-plane distortion of the double bond. This expectation has now been realized by the synthesis and resolution of *dl-cis*- (II) and *dl-trans*-4,4'-bi-1,1',2,2',3,3'-hexahydrophenanthrylidene (III) using a novel method of resolution (as applied to olefins).

When 4.1 g (21 mmol) of 4-keto-1,2,3,4-tetrahydrophenanthrene (I)² in 25 ml of dry THF was added, under nitrogen, to the black suspension resulting from mixing 6.5 g of TiCl₃ and 0.75 g of LiAlH₄ in 20 ml of THF, an immediate reaction occurred.³ The resulting mixture was stirred and heated for 4 h under reflux and after a normal workup 4.0 g of an oil could be isolated. The apolar fraction of the complex reaction mixture consisted of 0.80 g of colorless crystalline olefins, separable into *cis* (II, mp 192–193 °C) and *trans* isomers (III, mp 234–235 °C) via preparative TLC (deactivated alumina) (Scheme I).⁴

The structures of both olefins are based upon the following observations. Correct elemental analysis of both olefins and spectral data in agreement with structures II and III were obtained. The olefins II and III were reconverted into starting ketone I by oxidation using RuO₄. Hydrogenation of either olefin gave the same dihydro product (5% Pd/C, 3 atm H₂, ethanol solution). Both olefins showed the expected C₂ symmetry in their ¹³C NMR spectra. Assignment of the *cis* configuration to the lower melting isomer (II) is based upon the 100-MHz ¹H NMR spectra and the UV spectra. In the *cis* isomer an upfield shift was observed for protons in the 5,5',6,6',7,7',8,8' positions. (ABCD system, δ 6.36, 6.48, 6.70, and 6.86). These protons are influenced by the shielding zones of the aromatic rings, indicating a helix-like shape of this isomer. In the 100-MHz ¹H NMR spectrum of the *trans* isomer a multiplet was observed for the protons in the 5,5',6,6',7,7',8,8',9,9',10,10' positions (δ 7.1–7.8). A multiplet at δ 8.2 is probably due to the protons in the 5,5' positions, deshielded as a result of steric interaction.⁵

A hypsochromic shift is observed for the low energy band in the UV spectrum of the *cis* isomer (II) compared to that of the *trans* isomer (III) (Figure 1 and 2). This effect is well known for the *cis*- and *trans*-di-1-naphthylstilbene system.⁶ The structural relationship of II and III was clearly demonstrated by the slow isomerization of either pure *cis*- or pure *trans* isomer to a *cis*-*trans* mixture.

Scheme I

Figure 1. UV(—), ORD(---), and CD(· · ·) spectra of *cis*-olefin II.

Both *dl-cis* and *dl-trans* olefins could be resolved using HPLC techniques by employing an alumina column impregnated with (+)-TAPA.⁷ Partial resolution of the *cis* isomer was achieved (*d*-(+)-II: $[\alpha]^{22}_{578} +88^\circ c 1.60 \times 10^{-3}$, *n*-hexane). In addition the *trans* isomer could be resolved completely (*l*-(-)-III: $[\alpha]^{22}_{578} -508^\circ c 3.14 \times 10^{-3}$, *n*-hexane). The ORD and CD spectra of II (Figure 1) and of III (Figure 2) showed large rotations and ellipticities.

Recently published CD data and calculations on rotational strength indicate torsion of the olefin chromophore as the dominant contribution in determining the sign of the Cotton effects.⁹ However, in this case the entire conjugated system rather than an isolated twisted olefin chromophore must be involved in the interpretation of the spectra. Important features to be taken into considerations are: The direction of the principal C₂ axes. Note that in the *cis* olefin this axis is perpendicular to a plane through the π -orbitals of the double bond, while in the *trans* olefin the C₂ axis lies in the plane of the π -orbitals of the double bond (we realize that it is awkward to speak of a "plane" through the twisted olefin π -orbitals). Thus the relations between the transition moments and the C₂ axes in the two olefins differ. This must be evident in the spectra. Whether true exciton splitting is present cannot at this time be answered unequivocally. To a first approximation the fact that we are dealing with a conjugated system would argue against it.

The *trans* isomer could in principle exist in two diastereomeric forms; one *dl-trans* (III) and one non-dissymmetric form (IV). The resolution of the *trans* olefin is proof of the structure as III and not IV. The partial stereoselectivity of the reductive dimerization of I, resulting in the formation of *dl-trans* olefin (III) only, might give an indication about the structure of the intermediate Ti complex.¹⁰ Stereoselective formation of the *cis* isomer (II) could also be achieved: reaction of I under the

Figure 2. UV (—), ORD (---), and CD (···) spectra of trans-olefin III.

above conditions, but in the presence of (–)-trans-2,3-methylenedioxybutane as solvent gave cis-isomer II as the only olefinic product. This product was nevertheless optically inactive.

Some additional aspects of the structures of the cis and trans olefins need to be mentioned. The structure of II resembles that of a heptahelicene (more accurately a tetrahydroheptahelicene) with a double bond replacing an aromatic ring. The olefin III has a double helix structure and resembles a double pentahelicene with a double bond replacing two aromatic rings.^{11a} Both chiral as well as achiral double helix structures have been reported in the helicene case.^{11b} The role of π -bond torsion has been the subject of many investigations,^{1,9} although almost all of the attention has been focused on *trans*-cyclooctene. NMR studies connected with π -bond torsion have been reported for systems such as bisfluorenylidene and bianthrone, but no stable enantiomers have been obtained thus far.¹² To our knowledge, olefins II and III represent the first instance in which a *dl*-cis and *dl*-trans form of one inherently dissymmetric olefin have been isolated and resolved into enantiomers.

The isomerization and racemizations of II and III may furnish further information about the torsional distortion in both olefins. This complex series of interconversions will be the subject of further investigations.

Acknowledgment. The authors wish to thank Dr. R. Helder and Dr. H. Numan for their assistance during the HPLC resolutions, Dr. H. Hiemstra for recording and interpreting the NMR spectra, and the Netherlands Organization for Pure

Research (Z.W.O.) for a graduate fellowship. We thank Professor A. Moscovitz for a brief but highly valuable discussion.

References and Notes

- (1) A. C. Cope, C. R. Ganellin, H. W. Johnson, Jr., T. V. van Auken, and H. J. S. Winkler, *J. Am. Chem. Soc.*, **85**, 3276 (1963); A. C. Cope, and A. S. Metha, *ibid.*, **86**, 5626 (1964); G. Krow, *Top. Stereochem.*, **5**, 31 (1970); M. Tichy and J. Sicher, *Tetrahedron Lett.*, 4609 (1969); M. Rösner and G. Köbrich, *Angew. Chem.*, **87**, 715 (1975).
- (2) G. Schroeter and H. Müller, *Ber.*, **62**, 645 (1929); R. D. Haworth, *J. Chem. Soc.*, 1125 (1932).
- (3) J. E. McMurry and M. P. Fleming, *J. Am. Chem. Soc.*, **96**, 4708 (1974); J. E. McMurry, *Acc. Chem. Res.*, **7**, 281 (1974); H. Wynberg, K. Lammertsma, and L. A. Hulshof, *Tetrahedron Lett.*, 3749 (1975); E. J. Corey, R. L. Danheiser, and S. Chandrasekaran, *J. Org. Chem.*, **41**, 260 (1976).
- (4) In the case of II and III, only one of the possible enantiomers is shown in Scheme I.
- (5) In the 100-MHz ¹H NMR spectra of both *cis*-II and *trans*-III isomers, multiplets were observed for the protons in the 1,1',2,2',3,3' positions. (II: δ 1.7–3.1; III: δ 1.6, 2.2, and 2.7). For the *cis* isomer an AB system was observed for the protons in the 9,9',10,10' positions.
- (6) Ch. Goeddicke and H. Stegemeyer, *Ber. Bunsenges. Phys. Chem.*, **73**, 782 (1969); H. Suzuki, "Electronic Absorption Spectra and Geometry of Organic Molecules", Academic Press, New York, N.Y., 1967, Chapter 14 and 18.
- (7) The first report of the use of (+)-TAPAB^a in chromatographic separation of optical isomers is due to Klemm et al.^{8b} Gil-Av et al.^{8c} have recently reported its use in HPLC for the resolution of helicenes. We have applied the technique independently to the resolution of heterohelicenes.^{8d}
- (8) (a) M. S. Newman and D. Lednicher, *J. Am. Chem. Soc.*, **87**, 4765 (1965); (b) L. H. Klemm and D. Reed, *J. Chromatogr.*, **3**, 264 (1960); (c) F. Mikes, G. Boshart, and E. Gil-Av, *J. Chem. Soc., Chem. Commun.*, 99 (1976); (d) H. Numan, R. Helder, and H. Wynberg, *Recl. Trav. Chim. Pays-Bas*, **95**, 211 (1976).
- (9) N. H. Andersen, C. R. Costin, and J. R. Shaw, *J. Am. Chem. Soc.*, **96**, 3692 (1974); C. C. Levin and R. Hoffmann, *ibid.*, **94**, 3446 (1972); A. D. Scott and A. D. Wrixon, *J. Chem. Soc. D*, 1182 (1969); *Tetrahedron*, **26**, 3695 (1970); A. Moscovitz, *Proc. R. Soc. London*, **297**, 16 (1967), and references cited therein.
- (10) Dreiding models of the intermediate Ti complexes as suggested by McMurry and co-workers³ indicate that, in going from the precursors of III and IV to the olefins, the formation of IV is disfavored due to steric hindrance.
- (11) (a) Whether we view II as an "open helicene", a "chiral and rigid stilbene", or a novel helical conjugated π -system is a matter of semantics. (b) R. Martin, *Angew. Chem.*, **86**, 727 (1974); W. H. Laarhoven and Th. J. H. M. Cuppen, *Tetrahedron Lett.*, 163 (1971); R. H. Martin, Ch. Eyndels, and N. Defay, *ibid.*, 2731 (1972).
- (12) I. Agranat and Y. Tapuhi, *J. Am. Chem. Soc.*, **98**, 615 (1976); H. O. Kalinowsky and H. Kessler, *Top. Stereochem.*, **7**, 295 (1973); I. R. Gault, W. D. Ollis, and I. O. Sutherland, *Chem. Commun.*, 269 (1970).

Ben Feringa, Hans Wynberg*

Department of Organic Chemistry, The University
Groningen, The Netherlands

Received July 13, 1976

Synthesis and Spectral Data of a Stable, Optically Active 1,2-Dioxetane

Sir:

Some years ago the synthesis and structure proof of the stable 1,2-dioxetane, derived from adamantylideneadamantane,¹ was reported. Subsequently this molecule has been studied in considerable detail.² Using the same olefin moiety, we have now prepared and characterized an *optically active* 1,2-dioxetane.³

In addition to the criteria we enumerated for the successful synthesis of the adamantylideneadamantane-1,2-dioxetane¹ (electron-rich olefin; avoidance of "ene" reaction and no steric hindrance), stereochemical considerations played a crucial role in our present synthetic strategy. (i) The possible difficulties in resolving a 1,2-dioxetane made us decide to prepare an optically active (and if possible optically pure) olefin, as precursor of the 1,2-dioxetane. (ii) Condition i requires either the synthesis of an olefin with a functional group allowing resolution, or the synthesis of an optically active olefin precursor. (iii) The position and nature of the substituent, which imparts chirality