University of Groningen

EPR and optical studies of the lowest $n\pi^*$ triplet state of p-benzoquinone in an isotopic mixed crystal

Veenvliet, Hendrik; Wiersma, Douwe A.

Published in:
The Journal of Chemical Physics

DOI:
[10.1063/1.1681097](https://doi.org/10.1063/1.1681097)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1974

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Veenvliet, H., & Wiersma, D. A. (1974). EPR and optical studies of the lowest $n\pi^*$ triplet state of p-benzoquinone in an isotopic mixed crystal. *The Journal of Chemical Physics*, 60(2), 704-705.
<https://doi.org/10.1063/1.1681097>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

*A. P. Sloan Fellow, 1971–1973.

- ¹A. D. Buckingham and P. J. Stephens, *Ann. Rev. Phys. Chem.* **17**, 399 (1966); P. J. Stephens, *J. Chem. Phys.* **52**, 3489 (1970); P. M. Schatz and A. J. McCaffery, *Quart. Rev.* **23**, 552 (1969).
- ²C. Djerassi, E. Bunnenberg, and D. L. Elder, *Pure Appl. Chem.* **25**, 57 (1971); J. P. Larkindale and D. J. Simkin, *J. Chem. Phys.* **55**, 5668 (1971); W. G. Dauben, J. I. Seeman, P. H. Wendschuh, G. Barth, E. Bunnenberg, and C. Djerassi, *J. Org. Chem.* **37**, 1209 (1972); L. Seamans, A. Moscowitz, G. Barth, E. Bunnenberg, and C. Djerassi, *J. Amer. Chem. Soc.* **94**, 6464 (1972); D. J. Shieh, S. H. Lin, and H. Eyring, *J. Phys. Chem.* **76**, 1844 (1972); A. J. McHugh, M. Gouterman, and C. Weiss, Jr., *Theor. Chim. Acta* **24**, 346 (1972); H. H. Dearman, *J. Chem. Phys.* **58**, 2135 (1973).
- ³D. J. Caldwell and H. Eyring, *The Theory of Optical Activity* (Wiley-Interscience, New York, 1971), p. 184; *J. Chem. Phys.* **58**, 1149 (1973); L. Seamans and A. Moscowitz, *ibid.* **56**, 1099 (1972).
- ⁴While this work was in progress, a successful calculation using the finite perturbation method and gauge-invariant orbitals appeared: L. Seamans and J. Linderberg, *Mol. Phys.* **24**, 1393 (1972).
- ⁵For π systems, see J. Linderberg, *Chem. Phys. Lett.* **1**, 39 (1967). A generalization for σ systems is possible (J. Linderberg, personal communication).
- ⁶R. G. Parr, *Quantum Theory of Molecular Electronic Structure* (Benjamin, New York, 1963).
- ⁷J. A. Pople and D. L. Beveridge, *Approximate Molecular Orbital Theory* (McGraw-Hill, New York, 1970).
- ⁸Except for alternant hydrocarbons in which perfect pairing causes the calculated B values to vanish: L. H. Seamans, Ph.D. dissertation, University of Minnesota, 1971, p. 24.

EPR and optical studies of the lowest $n\pi^*$ triplet state of *p*-benzoquinone in an isotopic mixed crystal

Hendrik Veenliet and Douwe A. Wiersma

Laboratory for Physical Chemistry of the University of Groningen, Groningen, The Netherlands

(Received 22 October 1973)

In this Communication we wish to report our preliminary results of magnetic resonance and phosphorescence studies of *p*-benzoquinone- h_4 (PBQ- h_4) as a trap in PBQ- d_4 . Previous Stark effect studies^{1,2} on pure PBQ- h_4 crystals gave strong support for the assignment of the lowest triplet state in the pure crystal as the $B_{1g}n\pi^*$ triplet state with the unobserved origin in the pure crystal at 18602.8 cm^{-1} . It was further suggested that this lowest state was so severely distorted that for certain b_{1u} type nuclear motions the molecule had acquired a double minimum potential well along these nuclear motions, with the unobserved origin being the *g*-inversion level of this well. It thus seems very important to further establish the nature of the lowest excited state in PBQ since this molecule seems very suitable for a detailed study of the subtle effects that may occur in a molecule where the electronic and nuclear motions cannot be considered independent. In such a molecule small nuclear perturbations (as isotopic substitution) may drastically disturb the spin density distribution and possibly spin-orbit coupling selectivity. We now wish to present some preliminary results of our experiments on the *p*-benzoquinone isotopic mixed crystal system that focus on these suggestions.

Single crystals of PBQ- d_4 doped with 1 mol% PBQ- h_4 were grown from a Bridgeman oven and oriented conoscopically. A selection of the phosphorescence bands observed (all polarized along the long molecular axis) in a 1% mixed crystal at 1.8°K is given in Table I.

Crucial to the analysis of this spectrum is the fact that the origin (*g*-inversion level) of the PBQ- h_4 trap was placed at $18613.0 \pm 1\text{ cm}^{-1}$. This position was deduced from our Stark² and mixed crystal absorption studies.³

With phase sensitive detection of the phosphorescence at different chopping frequencies it was possible to separate b_{1u} , b_{2u} , and b_{3u} modes in the emission arising from independently decaying T_x , T_y , and T_z spin substates, respectively. This is as expected for emission from a $^3B_{1g}n\pi^*$ state that is spin-vibronic coupled to a strongly allowed $^1B_{1u}(\pi\pi^*)$ state. At 4.2°K emission exclusively from the T_x spin substate is observed and all bands of assigned b_{2u} and b_{3u} symmetry are completely absent. The same situation can be created at 1.8°K when a magnetic field is applied.

Further conformation of the analysis presented came from ODMR experiments. Zero field transitions at

TABLE I. Microwave effect on and assignment of selected phosphorescence bands of *p*-benzoquinone- h_4 (1 mol%) in *p*-benzoquinone- d_4 at 1.8°K [$L = z$ (B_{1u}), $M = y$ (B_{2u})].

Position (VAC cm^{-1})	Assignment ^a	Microwave effect (% $\pm$ 2%)	
		1927 MHz	2165 MHz
18 613	Origin; not observed		
18 195	418 (ν_{21}, b_{2u})	- 18	None
18 082	531 (ν_{25}, b_{3u})	None	- 4
17 863	750 (ν_{16}, b_{1u})	+ 41	+ 15
17 832	781 (ν_{25}, b_{3u}) + (ν_{30}, b_{2g})	+ 41	+ 15
17 747	866 (ν_{21}, b_{2u}) + (ν_6, a_g)	- 2	None
17 705	908 (ν_{24}, b_{3u})	None	- 9
17 666	947 (ν_{15}, b_{1u})	+ 26	+ 12
17 422	1191 (ν_{21}, b_{2u}) + (ν_5, a_g)	- 7	None
17 294	1319 (ν_{19}, b_{2u})	- 7	None
17 242	1371 (ν_{14}, b_{1u})	+ 25	+ 12
16 953	1660 (ν_{13}, b_{1u})	+ 41	+ 12

^aGround state fundamentals have been taken from A. H. Francis, Ph.D. thesis, University of Michigan, Ann Arbor, MI, 1969.

1927 $\pm$ 5 MHz and 2164 $\pm$ 5 MHz were detected with this technique and the percentages change in phosphorescence observed at these frequencies for the individual bands are also given in Table I. It must be concluded that T_x is the most radiative ($\tau_x = 100$ μ sec) and underpopulated level at 1.8 $^\circ$ K. The third zero field transition could not be detected with this method but was inferred to arise at 237 $\pm$ 10 MHz from ODMR experiments in a magnetic field. These measurements also gave the relative signs of D and E and the g -tensor principal values:

$$|(D+E)/h| = 1927 \pm 5 \text{ MHz} \quad g_{xx} = 2.0043 \pm 0.0010$$

$$|(D-E)/h| = 2164 \pm 5 \text{ MHz} \quad g_{yy} = 2.0010 \pm 0.0006$$

$$|2E/h| = 237 \pm 10 \text{ MHz} \quad g_{zz} = 2.0080 \pm 0.0006$$

The large deviation of g_{zz} from the free electron value shows the expected effect of direct spin-orbit coupling of the T_x spin sublevel with the ground state. Quite surprising is also the fact that our measured ZFS parameters of PBQ- h_4 in PBQ- d_4 are completely different from those measured by Attia *et al.*⁴ in a p -dibromobenzene host crystal:

$$|(D+E)/h| = 5224 \text{ MHz}; \quad |(D-E)/h| = 5379 \text{ MHz},$$

and

$$|2E/h| = 155 \text{ MHz}.$$

This fact already indicates that the p -benzoquinone B_{1g} state is extremely sensitive to perturbations. This is also evident from the drastic change that occurs when the ZFS parameters of PBQ- dh_3 in the same host crystal (PBQ- d_4) are measured. We report the following numbers: $|(D+E)/h| = 2316 \pm 5 \text{ MHz}$, $g_{xx} = 2.0040 \pm 0.0010$; $|(D-E)/h| = 2526 \pm 5 \text{ MHz}$, $g_{yy} = 2.0012 \pm 0.0006$; $|2E/h| = 210 \pm 10 \text{ MHz}$, $g_{zz} = 2.0080 \pm 0.0006$. Such an enormous change in D can only be accounted for on the basis of an electronic state that is extremely sensitive to the exact form of the vibrational force field. In the case of a dou-

ble minimum potential well, asymmetric ring substitution favors one of the distorted configurations that may result in a drastic change in the spin density distribution. We therefore take the observed hypersensitivity of the ZFS parameters of PBQ to asymmetric deuteration as a strong argument in favor of the proposed² double minimum well of the $^3B_{1g}n\pi^*$ state. Further EPR, ENDOR, and Stark-experiments on PBQ and its deuterio isotopes are now underway and will be reported in a future paper.

Finally we would like to report observation of strong level anticrossing and cross relaxation effects in the phosphorescence of PBQ at 1.8 $^\circ$ K completely similar to the effects observed by Veeman and van der Waals⁵ on the lowest $n\pi^*$ triplet state of benzophenone.

We are most indebted to Philip Vergragt of the University of Leiden for his invaluable help in obtaining the ODMR spectra of p -benzoquinone.

We are also grateful to A. H. Francis for sending us copies of papers dealing with p -benzoquinone prior to their publication.

¹R. M. Hochstrasser, L. W. Johnson, and H. P. Trommsdorff, Chem. Phys. Lett. 21, 251 (1973).

²H. Veenliet and D. A. Wiersma, Chem. Phys. Lett. 22, 87 (1973).

³H. Veenliet and D. A. Wiersma, Proc. 6th Mol. Cryst. Symp. Elmau, Germany, 1973, p. 90.

⁴A. I. Attia, B. H. Loo, and A. H. Francis, Chem. Phys. Lett. 22, 537 (1973).

⁵W. S. Veeman and J. H. van der Waals, Chem. Phys. Lett. 7, 65 (1970).

Theoretical calculations of the second virial coefficient of water

J. G. Lane* and H. W. Leidecker

Department of Physics, The American University, Washington, D.C. 20016
(Received 2 October 1973)

In the process of investigating the water pair potential we have calculated the second virial coefficient for a number of potentials which have in common a Lennard-Jones spherically symmetric part and various representations of the water molecule electrostatic charge interaction. The numerical calculation yields results with a precision of 1%, which is better than the experimental measurements. The results of these calculations also allow us to estimate the second virial coefficient for the Hartree-Fock potential surface calculated by Del Bene

and Pople.¹

The electrostatic-type potentials, $U_{M1} = U_{LJ} + U_{\mu\mu}$ (Stockmayer²), $U_{M2} = U_{LJ} + U_{\mu\mu} + U_{\mu Q}$, $U_{M3} = U_{LJ} + U_{\mu\mu} + U_{\mu Q} + U_{QQ} + U_{\mu\Omega}$, and $U_{BNS} = U_{LJ} + U_{\text{point charge}}$ (Ben-Naim and Stillinger³), form a sequence with increasingly more complete descriptions of the electrostatic interactions. The multipole moments used for $M1$, $M2$, and $M3$ (calculated relative to the center of mass of $H_2^{16}O$) are $\mu_z = 1.85 \times 10^{-18}$ esu $\cdot$ cm, $Q_{zz} = -.13 \times 10^{-26}$ esu $\cdot$ cm², Q_{xx}