University of Groningen

Cytogenetic studies on *Solanum tuberosum* L. and some of its relatives

Koopmans, Aaltje

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

1951

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Koopmans, A. (1951). *Cytogenetic studies on *Solanum tuberosum* L. and some of its relatives*. [Thesis fully internal (DIV), University of Groningen]. Martinus Nijhoff.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supposing like ÖSTERGREN does (1949) that the spindle figure is a tactoid, and does not exist of fibres but of particles in dynamic equilibrium, continuously exchanging position with one another as in a liquid, but still preserving their characteristic parallel orientation it is possible to imagine that the working on the nuclear plasm is in such a way that may be by change of viscosity, no orientation of the particles occurs.

An important field of research still lies open: it seems more and more necessary not only to look at the final result, but to study the colchicine action much more in detail as a cytological, and not a purely karyological, process.

SUMMARY

- 1) Native materials, obtained through the courtesy of Professor Dr A. C. BOERGER, Uruguay, and considered to be *Solanum commersonii*, have a somatic number of chromosomes of $2n = 24$. Therefore it seems probable that the plants having a somatic number of $2n = 36$ as studied by other writers, do not belong to this species.
- 2) On the strength of the results obtained from crosses between different species it was tried to establish a relationship between the genomes.
- 3) The cytological assumption that the basic number of chromosomes in the genus *Solanum* is 6, was strengthened by genetical data.
- 4) It was demonstrated that the action of colchicine is primarily cytological, and secondarily karyological.
- 5) Three new species are mentioned, viz. *Solanum antipophureja* ($2n = 72$), *S. antipochacoense* ($2n = 72$) and *S. demissorosum* ($2n = 120$).

SAMENVATTING

- 1) In het wild groeiend materiaal, ontvangen van Prof. Dr. A. C. BOERGER, Uruguay, dat door hem als *Solanum commersonii* beschouwd wordt, heeft een somatisch chromosomenaantal van $2n = 24$. In verband hiermede lijkt het waarschijnlijk, dat de