University of Groningen

Homogeneous cis-dihydroxylation and epoxidation of olefins with high H₂O₂ efficiency by mixed manganese/activated carbonyl catalyst system

Brinksma, J.; Schmieder, L.; van Vliet, G.; Boaron, R.; Hage, R.; De Vos, D.E.; Alsters, P.L.; Feringa, B.L.

Published in:
Tetrahedron Letters

DOI:
[10.1016/S0040-4039\(02\)00292-7](https://doi.org/10.1016/S0040-4039(02)00292-7)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2002

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Brinksma, J., Schmieder, L., van Vliet, G., Boaron, R., Hage, R., De Vos, D. E., Alsters, P. L., & Feringa, B. L. (2002). Homogeneous cis-dihydroxylation and epoxidation of olefins with high H₂O₂ efficiency by mixed manganese/activated carbonyl catalyst system. *Tetrahedron Letters*, 43(14), 2619 - 2622. [PII S0040-4039(02)00292-7]. [https://doi.org/10.1016/S0040-4039\(02\)00292-7](https://doi.org/10.1016/S0040-4039(02)00292-7)

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Homogeneous *cis*-dihydroxylation and epoxidation of olefins with high H₂O₂ efficiency by mixed manganese/activated carbonyl catalyst system

Jelle Brinksma,^a Lizette Schmieder,^b Gerbert van Vliet,^b Rob Boaron,^b Ronald Hage,^c
Dirk E. De Vos,^d Paul L. Alsters^{b,*} and Ben L. Feringa^{a,*}

^aLaboratory of Organic Chemistry, Stratingh Institute, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands

^bDSM Fine Chemicals, Advanced Synthesis and Catalysis, PO Box 18, 6160 MD Geleen, The Netherlands

^cUnilever Research Laboratory Vlaardingen, PO Box 114, 3130 AC Vlaardingen, The Netherlands

^dCentre for Surface Chemistry and Catalysis, Katholieke Universiteit Leuven, Kasteelpark Arenberg 23, 3001 Leuven, Belgium

Received 18 January 2002; accepted 8 February 2002

Abstract—The use of [Mn₂O₃(tmtacn)₂](PF₆)₂ (tmtacn = 1,4,7-trimethyl-1,4,7-triazacyclononane) in combination with glyoxylic acid methylester methyl hemiacetal (GMHA) results in a highly active and hydrogen peroxide efficient catalyst for the epoxidation of olefins as well as the first homogeneous catalytic *cis*-dihydroxylation system with H₂O₂ and with turnover numbers up to 420. © 2002 Elsevier Science Ltd. All rights reserved.

For economic and environmental reasons catalytic olefin oxidations based on O₂ or H₂O₂ are preferred over traditional stoichiometric oxidations, e.g. epoxidation with peracids and *cis*-dihydroxylation with permanganate.¹ Several methods are available for catalytic epoxidation with aqueous H₂O₂ (notably Re-, W-, and Mn-based catalysts),² but high turnovers for *cis*-dihydroxylation are only achieved using Os.³ However, the high cost and toxicity of Os hamper large scale application and provide a strong incentive to develop benign Fe- or Mn-based *cis*-dihydroxylation catalysts. Que et al. recently reported the first *cis*-dihydroxylation with H₂O₂ catalysed by a non-heme iron complex.⁴ Although this system shows good *cis*-diol selectivities, turnover numbers (t.o.n.) are rather low (up until 22 t.o.n.).

Recently Mn complexes based on tmtacn were found to be highly active in catalytic oxidation,⁵ but apart from high turnover, it is essential to develop catalytic systems that employ H₂O₂ very efficiently, as many Mn or Fe

catalysts are known to be particularly effective in decomposition of H₂O₂.⁶ This can be suppressed by working in acetone;⁷ however, this solvent is not acceptable for large-scale applications because of the risk of formation of explosive cyclic peroxides. H₂O₂ decomposition by Mn–tacn complexes can be suppressed by addition of oxalate^{8a} or ascorbic acid^{8b} as co-catalysts, or by anchoring the tacn ligand to a solid support.⁹ Significant amounts of *cis*-diols besides epoxides are formed during alkene oxidation with this heterogeneous Mn catalyst.

Herein, we report efficient alkene oxidations with H₂O₂ catalysed by Mn–tmtacn **1**¹⁰ in the presence of GMHA **2** (Fig. 1). Surprisingly, the latter not only strongly suppresses H₂O₂ decomposition by the Mn catalyst, but also imparts the first homogeneous catalytic *cis*-dihydroxylation activity to a Mn catalyst. Catalytic oxidations were performed by adding aqueous, 50% hydrogen peroxide (1.3 equiv. with respect to the sub-

Figure 1. L = 1,4,7-trimethyl-1,4,7-triazacyclononane.

Keywords: hydrogen peroxide; *cis*-diols; epoxides; manganese; oxidation.

* Corresponding authors. Tel.: +31 50 363 4278; fax: +31 50 363 4296; e-mail: b.l.feringa@chem.rug.nl

strate) in 6 h to a mixture of alkene, Mn–tmtacn catalyst (0.1 mol%), and GMHA (25 mol%) in MeCN at 0°C. Under these reaction conditions, the presence of GMHA as the co-catalyst enables high conversions with only a 30% excess of oxidant with respect to substrate (Table 1) and the H₂O₂ efficiency is drastically improved compared to previous Mn systems using excess of H₂O₂.⁵ In most cases the conversions were also significantly higher than those obtained with oxalate as co-catalyst and 1.3 equiv. of H₂O₂ (for similar experiments with a larger excess of H₂O₂, see Ref. 8a).

In particular, styrene afforded a high yield of styrene oxide in the presence of GMHA, whereas the use of oxalate resulted in only modest conversion of styrene. When a mixture of GMHA (25 mol%) and oxalate (0.3 mol%) was used as the co-catalyst system, the epoxide yields even surpassed those obtained with the Mn/GMHA catalytic system, and high epoxide yields at complete conversion were obtained for the non-sterically hindered alkenes (Table 1, entries 1–3, 6, 8). Unexpectedly, substantial amounts of *cis*-diols were formed besides epoxides when *only* GMHA was present as the co-catalyst. The epoxide/*cis*-diol ratio depends

strongly on the alkene structure. The highest amount of *cis*-diol was found for cyclooctene (entry 3), which afforded the *cis*-diol as the main product (42%, 420 t.o.n.) besides the epoxide (36%, 360 t.o.n.). Minor amounts of 2-hydroxycyclooctanone were also found due to overoxidation of the diol. The ring size of cycloalkenes has a profound influence on the epoxide/*cis*-diol ratio (entries 1–4). For these cyclic olefins, almost no *trans*-diol could be detected (ratio *cis*-diol/*trans*-diol >99.5/0.5). *cis*-Diol formation is also observed for aliphatic acyclic alkenes. Yields of diol are significantly lower for *trans*-2-hexene than from *cis*-2-hexene, but the epoxide/*cis*-diol ratio was similar for both substrates. The aryl-substituted alkenes (entries 7, 8) yield epoxide nearly exclusively under these conditions. Limited *cis/trans* isomerisation is observed in the epoxide formation of *cis*-2-hexene. The *cis/trans* isomerisation points to epoxidation via a Mn oxo species, with formation of epoxides from C-centred radical intermediates with a lifetime sufficient for some C–C bond rotation prior to reaction to the epoxide.¹² In line with this mechanism, olefins that form a relatively long-lived radical intermediate, such as *cis*-stilbene, show substantial loss of configuration in epoxide. No diols were formed on replacing the substrate by epox-

Table 1. Oxidation of selected olefins by H₂O₂ with catalyst 1¹¹

Entry	Substrate	Conversion (%) ^c			Product ^d	Turnover number (t.o.n.) ^c		
		Additive: GMHA ^a (2)	Additive: 2 + oxalate ^b	Additive: oxalate ^c		Additive: GMHA ^a (2)	Additive: 2 + oxalate ^b	Additive: oxalate ^c
1	Cyclopentene	97	100	92	Epoxide	610	860	770
					<i>cis</i> -Diol	260	0	0
					Cyclopentenone	80	10	100
2	Cyclohexene	88	100	73	Epoxide	590	820	610
					<i>cis</i> -Diol	90	0	0
					2-Cyclohexenone	80	80	70
3	Cyclooctene	90	100	52	Epoxide	360	840	410
					<i>cis</i> -Diol	420	60	0
					2-HO-cyclooctanone	220	80	0
4	Norbornylene	95	88	59	<i>exo</i> -Epoxide	540	530	360
					<i>exo-cis</i> -Diol	180	60	0
5	<i>trans</i> -2-Hexene	77	93	37	<i>trans</i> -Epoxide	210	550	240
					<i>cis</i> -Epoxide	50	30	0
					<i>RR/SS</i> -Diol	150	90	0
					<i>RS/SR</i> -Diol	0	0	0
					<i>cis</i> -Epoxide	450	820	640
6	<i>cis</i> -2-Hexene	93	100	80 ^f	<i>trans</i> -Epoxide	40	30	20
					<i>SR/RS</i> -Diol	280	10	10
					<i>RR/SS</i> -Diol	10	0	0
					<i>cis</i> -Epoxide	260	260	110
					<i>trans</i> -Epoxide	200	160	150
7	<i>cis</i> -Stilbene	82	77	32	<i>meso</i> -Hydrobenzoin	40	30	10
					Hydrobenzoin	40	20	10
					Epoxide	860	860	380
8	Styrene	97	100	44	Ph(CH)(OH)CH ₂ OH	60	50	0
					PhC(O)CH ₂ OH	10	30	0

^a Co-catalyst GMHA (25 mol%).

^b Co-catalyst mixture of GMHA (25 mol%) and oxalate (0.3 mol%).

^c Co-catalyst oxalate (0.3 mol%), for similar experiments with a larger excess of H₂O₂, see Ref. 8a.

^d All products were identical to independent samples and identified by GC and ¹H NMR.

^e Determined by GC.

^f 32 mmol scale

Figure 2. Time profile of the oxidation of cyclooctene by H_2O_2 with catalyst **1** and GMHA; ■ epoxide; ● *cis*-diol.

ide, thus excluding epoxide hydrolysis.¹³ This conclusion is further supported by the time course profile of the **1**/GMHA-catalysed oxidation of cyclooctene, which shows that both epoxide and diol increase progressively with time (Fig. 2).

Attempts to increase the *cis*-diol selectivity by addition of acid or base resulted in a dramatic decrease of conversion. Besides GMHA, a few other carbonyl compounds were tested as co-catalyst (25 mol%) for the oxidation of cyclooctene with H_2O_2 (1.3 equiv.). In the presence of glyoxylic acid hydrate, the epoxide was formed in 840 t.o.n. and *cis*-diol formation was almost completely suppressed. Modest *cis*-diol yields (<250 t.o.n.) and predominant formation of the epoxide were seen with diethyl ketomalonate or 2-ketoglutaric acid as the co-catalysts. However, *cis*-diol formation (370 t.o.n.) predominated over epoxide (310 t.o.n.) formation in the presence of chloral hydrate. Thus, a variety of carbonyl compounds with an adjacent electron withdrawing group are able to reduce H_2O_2 decomposition by **1** and to impose *cis*-dihydroxylation activity on manganese catalyst **1**, albeit with variable *cis*-diol/epoxide ratios.

Since *cis*-diol formation through Mn-catalysed epoxide hydrolysis can be excluded (vide supra), we propose that the *cis*-diol is formed by reaction of the alkene with an Mn oxo-hydroxo species. As in the case of oxalate, hydrated activated carbonyl compounds¹⁴ might break down the catalase active binuclear Mn complex **1**¹⁵ into a mononuclear Mn species via complexation to the Mn centre. *cis*-Diol formation from an Mn oxo-hydroxo species with a coordinated hydrated carbonyl ligand could be induced through a hydrogen bonded six-membered ring transition state (concerted pathway, Scheme 1). Reoxidation of the Mn centre with H_2O_2 , release of the diol from Mn, and hydration of the carbonyl compound closes the catalytic cycle. In conclusion, use of activated carbonyl compounds in

Scheme 1. Proposed *cis*-dihydroxylation mechanism (L = tmtacn, X = CO_2Me , CCl_3).

combination with Mn–tmtacn results in a highly active (up to 860 t.o.n.) and H_2O_2 efficient epoxidation system. Besides epoxidation this new catalytic system also provides, to the best of our knowledge, the most active Os-free homogeneous catalyst for *cis*-dihydroxylation (up to 420 t.o.n.). Compared with the anchored Mn–tmtacn catalyst,⁹ the present homogeneous **1**/activated carbonyl compound system is much more accessible, since both **1** and many activated carbonyl compounds have large scale applications. Efforts to enhance diol selectivity and to elucidate the *cis*-dihydroxylation mechanism are in progress.

Acknowledgements

We thank Dr. D. Schipper (DSM Food Specialities) for NMR experiments on the GMHA/50% aq. H_2O_2 system and DSM Fine Chemicals Austria for supply of GMHA.

References

1. (a) Hill, C. L.; Prosser-McCartha, C. M. *Coord. Chem. Rev.* **1995**, *143*, 407–455; (b) Sato, K.; Aoki, M.; Noyori, R. *Science* **1998**, *281*, 1646–1646; (c) Ten Brink, G. J.; Arends, I. W. C. E.; Sheldon, R. A. *Science* **2000**, *287*, 1636–1639.
2. (a) Sato, K.; Aoki, M.; Ogawa, M.; Hashimoto, T.; Noyori, R. *J. Org. Chem.* **1996**, *61*, 8310–8311; (b) Hermann, W. A.; Fischer, R. W.; Marz, D. W. *Angew. Chem., Int. Ed. Engl.* **1991**, *30*, 1638–1641; (c) Rudolph, J.; Reddy, K. L.; Chiang, J. P.; Sharpless, K. B. *J. Am. Chem. Soc.* **1997**, *119*, 6189–6190; (d) Larrow, J. F.; Jacobsen, E. N.; Gao, Y.; Hong, Y.; Nie, X.; Zepp, C. M. *J. Org. Chem.* **1994**, *59*, 1939–1942; (e) Hosoya, N.; Hatayama, A.; Yanai, K.; Fujii, H.; Irie, R.; Katsuki, T. *Synlett* **1993**, 641–645; (f) Hage, R.; Iburg, J. E.; Kerschner, J.; Koek, J. H.; Lempers, E. L. M.; Martens, R. J.; Racherla, U. S.; Russell, S. W.; Swarthoff, T.; Van Vliet, M. R. P.; Warnaar, J. B.; Van Der Wolf, L.; Krijnen, B. *Nature* **1994**, *369*, 637–639; (g) Lane, B. S.; Burgess, K. *J. Am. Chem. Soc.* **2001**, *123*, 2933–2934.
3. (a) Kolb, H. C.; VanNieuwenhze, M. S.; Sharpless, K. B. *Chem. Rev.* **1994**, *94*, 2483–2547; (b) Döbler, C.; Mehlretter, G. M.; Sundermeier, U.; Beller, M. *J. Am. Chem. Soc.* **2000**, *122*, 10289–10290; (c) Jonsson, S. Y.; Färnegårdh, K.; Bäckvall, J.-E. *J. Am. Chem. Soc.* **2001**, *123*, 1365–1371.
4. Chen, K.; Que, L., Jr. *Angew. Chem., Int. Ed.* **1999**, *38*, 2227–2229.
5. (a) De Vos, D. E.; Bein, T. *Chem. Commun.* **1996**, 917–918; (b) Zondervan, C.; Hage, R.; Feringa, B. L. *Chem. Commun.* **1997**, 419–420.
6. Recently Jacobsen reported a very efficient Fe-based epoxidation catalyst: White, M. C.; Doyle, A. G.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2001**, *123*, 7194–7195.
7. Knops-Gerrits, P.-P.; De Vos, D. E.; Jacobs, P. A. *J. Mol. Catal. A.* **1997**, *117*, 57–70.
8. (a) De Vos, D. E.; Sels, B. F.; Reynaers, M.; Subba Rao, Y. V.; Jacobs, P. A. *Tetrahedron Lett.* **1998**, *39*, 3221–3224; (b) Berkessel, A.; Sklorz, C. A. *Tetrahedron Lett.* **1999**, *40*, 7965–7968.
9. De Vos, D. E.; De Wildeman, S.; Sels, B. F.; Grobet, P. J.; Jacobs, P. A. *Angew. Chem., Int. Ed.* **1999**, *38*, 980–983.
10. Wiegardt, K.; Bossek, U.; Nuber, B.; Weiss, J.; Bonvoisin, J.; Corbella, M.; Vitols, S. E.; Girerd, J. J. *J. Am. Chem. Soc.* **1988**, *110*, 7398–7411.
11. Typical oxidation procedure with Mn–tmtacn/GMHA system: H₂O₂ (50% aq., 52 mmol) was added in 6 h at 0°C to a solution of Mn₂O₃(tmtacn)₂(PF₆)₂ (**1**, 0.04 mmol), GMHA (10 mmol), substrate (40 mmol) and internal standard (1,2-dichlorobenzene, 20 mmol) in acetonitrile (40 mL). Samples for GC (HP 6890, column HP1 15×0.3 mm×2.65 μm, polymethylsiloxane) analysis were taken 1 h after complete addition of oxidant.
12. Jacobsen, E. N.; Deng, L.; Furukawa, Y.; Martinez, L. E. *Tetrahedron* **1994**, *50*, 4323–4334. With the combined epoxidation/*cis*-dihydroxylation by a heterogenised Mn–tmtacn complex also limited *cis/trans* isomerisation was observed (see Ref. 9).
13. Quee-Smith, V. C.; DelPizza, L.; Jureller, S. H.; Kerschner, J. L.; Hage, R. *Inorg. Chem.* **1996**, *35*, 6461–6465.
14. ¹GMHA is an equilibrium mixture, which also contains some hydrated methyl glyoxylate. NMR experiments showed that formation of peroxyhydrate from GMHA and aqueous H₂O₂ is very slow.
15. Hage, R. *Recl. Trav. Chim. Pays-Bas* **1996**, *115*, 385–395.