University of Groningen

A chiral Lewis-acid-catalyzed Diels-Alder reaction. Water- enhanced enantioselectivity

Otto, S.; Boccaletti, G.; Engberts, J.B.F.N.

Published in:
Journal of the American Chemical Society

DOI:
[10.1021/ja972772+](https://doi.org/10.1021/ja972772+)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1998

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Otto, S., Boccaletti, G., & Engberts, J. B. F. N. (1998). A chiral Lewis-acid-catalyzed Diels-Alder reaction. Water- enhanced enantioselectivity. *Journal of the American Chemical Society*, 120(17), 4238 - 4239.
<https://doi.org/10.1021/ja972772+>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

A Chiral Lewis-Acid-Catalyzed Diels–Alder Reaction. Water-Enhanced Enantioselectivity

Sijbren Otto, Giovanni Boccaletti, and Jan B. F. N. Engberts*

Department of Organic and Molecular
Inorganic Chemistry, University of Groningen
Nijenborgh 4, 9747 AG Groningen, The Netherlands

Received August 8, 1997

Revised Manuscript Received March 16, 1998

Water is becoming increasingly popular as a medium for organic reactions.¹ Apart from the obvious economic and environmental benefits,² the aqueous medium can have favorable effects on many organic transformations.³ In the field of Lewis-acid catalysis, the use of water is still in its infancy.⁴ Accordingly, reports on Lewis-acid catalysis of Diels–Alder reactions in water⁵ or mixtures of organic solvents with small amounts of water⁶ are scarce. Examples of *enantioselective* Lewis-acid-catalyzed Diels–Alder reactions in water have been completely lacking to date.

In organic solvents, enantioselective Lewis-acid catalysis of Diels–Alder reactions has been studied extensively.⁷ The choice of the solvent for these reactions is rather limited. Donating solvents are generally avoided since they exert detrimental effects on rate and enantioselectivity.⁸ Nevertheless, on the basis of our previous work on Lewis-acid catalysis of Diels–Alder reactions in water,^{5a,b} we envisaged that even in aqueous media enantioselective catalysis should be possible.

In recent studies,^{5a,b} we have already shown that water can have a beneficial effect on the rate of the Lewis-acid-catalyzed Diels–Alder reaction of 3-phenyl-1-(2-pyridyl)-2-propen-1-one (**1**) with cyclopentadiene (**2**) (Scheme 1). We now report that for this reaction enantioselectivity can be induced by coordinating chiral commercially available α -amino acids to a catalytically active metal ion. Derivatives of α -amino acids have already been reported to induce large enantioselectivities in Diels–Alder reactions in organic solvents.^{8a,9}

In the presence of 10 % percent of copper(II) complexes of glycine, L-valine, L-leucine, L-phenylalanine, L-tyrosine, L-tryptophan, and *N* α -methyl-L-tryptophan (L-abrine), the Diels–Alder adduct **3** is obtained after 48 h in yields generally exceeding

Scheme 1

90%.¹⁰ Most interestingly, enantioselectivity is observed for the chiral ligands (Table 1). With L-abrine, an enantiomeric excess (ee) of 74% can be achieved. The catalyst solution can be reused without decrease in enantioselectivity. To the best of our knowledge, this is the first example of an enantiomeric excess in a chiral Lewis-acid-catalyzed reaction in water.

Significant enantioselectivities are observed exclusively for the α -amino acids containing aromatic side groups. This suggests that arene–arene interactions are important in discriminating between the two pathways leading to the enantiomeric Diels–Alder adducts. The importance of these interactions was further assessed by examining the effect of the α -amino acid ligands on the following: (1) the equilibrium constants (K_1 in Scheme 1) for binding of the dienophile to the copper–ligand complex and (2) the second-order rate constants (k_2) for reaction of **2** with the thus-formed complex. The results are shown in Table 1. When compared to the aquo complex, the presence of the aliphatic α -amino acids glycine, L-valine, and L-leucine at the copper center causes a decrease of the equilibrium constants for binding of **1** to this copper ion. Presumably, this effects is a consequence of the reduction of the number of coordination sites available on the copper center in combination with a reduction of the Lewis acidity of the metal ion as a result of the presence of the Lewis-basic α -amino acid ligand.

Interestingly, for the aromatic α -amino acids an additional and counteractive effect is operative, which *raises* the equilibrium

(10) In a typical procedure a solution of 0.175 mmol of L-abrine and 0.175 mmol of NaOH in 1 mL of water was added to a solution of 0.100 mmol of Cu(NO₃)₂ in 100 mL of water. The pH was adjusted to 6–6.5. The catalyst solution was cooled to 0 °C, and a solution of 1.0 mmol of **1** in a minimal amount of ethanol was added, together with 2.0 mmol of **2**. After 48 h of stirring at 0 °C, the reaction mixture was extracted with ether affording 0.94 mmol (94%) of **3**. All attempts to determine the absolute configuration of **3** and several derivatives thereof by X-ray diffraction have been as yet unsuccessful, due to problems in obtaining suitable crystals.

(1) (a) Li, C. *Chem. Rev.* **1993**, 93, 2023. (b) Lubineau, A. *Chem. Ind.* **1996**, 123. (c) Cornils, B.; Herrmann, W. A.; Eckl, R. W. *J. Mol. Catal., A Chem.* **1997**, 116, 27. (d) *Organic Synthesis in Water*; Grieco, P. A., Ed.; Blackie Academic and Professional: London, 1998.

(2) Cornils, B.; Wiebus, E. *Recl. Trav. Chim. Pays-Bas* **1996**, 115, 211.

(3) Particularly the Diels–Alder reaction benefits strongly from the use of water as the solvent, as was discovered by Breslow (Rideout, D. C.; Breslow, R. J. *J. Am. Chem. Soc.* **1980**, 102, 7816). For recent references, see: (a) Otto, S.; Blokzijl, W.; Engberts, J. B. F. N. *J. Org. Chem.* **1994**, 59, 5372. (b) Engberts, J. B. F. N. *Pure Appl. Chem.* **1995**, 67, 823.

(4) Engberts, J. B. F. N.; Feringa, B. L.; Keller, E.; Otto, S. *Recl. Trav. Chim. Pays-Bas* **1996**, 115, 457.

(5) (a) Otto, S.; Engberts, J. B. F. N. *Tetrahedron Lett.* **1995**, 36, 2645. (b) Otto, S.; Bertoncin, F.; Engberts, J. B. F. N. *J. Am. Chem. Soc.* **1996**, 118, 7702. (c) Loh, T.; Pei, J.; Lin, M. J. *J. Chem. Soc., Chem. Commun.* **1996**, 2315. (d) Yu, L.; Chen, D.; Wang, P. G. *Tetrahedron Lett.* **1996**, 37, 2169. (e) Yu, L.; Li, J.; Ramirez, J.; Chen, D.; Wang, P. G. *J. Org. Chem.* **1997**, 62, 903. (f) Zhu, Z.; Espenson, J. H. *J. Am. Chem. Soc.* **1997**, 119, 3507.

(6) (a) Kobayashi, S.; Hachiya, I.; Araki, M.; Ishitani, H. *Tetrahedron Lett.* **1993**, 34, 3755. (b) Kobayashi, S. *Synlett* **1994**, 689.

(7) For a review, see: Kagan, H. B.; Riant, O. *Chem. Rev.* **1992**, 92, 1007. (8) (a) Sartor, D.; Saffrich, J.; Helmchen, G.; Richards, C. J.; Lambert, H. *Tetrahedron: Asymmetry* **1991**, 2, 639. (b) Nevalainen, V. *Tetrahedron: Asymmetry* **1991**, 2, 827.

(9) (a) Takasu, M.; Yamamoto, H. *Synlett* **1990**, 194. (b) Sartor, D.; Saffrich, J.; Helmchen, G. *Synlett* **1990**, 197. (c) Corey, E. J.; Loh, T. *J. Am. Chem. Soc.* **1991**, 113, 8966. (d) Corey, E. J.; Loh, T.; Roper, T. D.; Azimioara, M. D.; Noe, M. *J. Am. Chem. Soc.* **1992**, 114, 8290.

Table 1. Effects of Ligands on the Equilibrium Constants (K_1)^a for Binding of **1** to the Copper(II)–Ligand Complex, the Second-order Rate Constants (k_2)^a for Reaction of **2** with This Complex, and the Enantioselectivity^b of the Latter Process at 25 °C^c

ligand	$10^{-3} K_1$ (M ⁻¹) ^d	k_2 (M ⁻¹ s ⁻¹) ^d	ee (%) ^{e,f,g}
water	1.2	2.56	0
glycine	0.74	1.89	0
L-valine	0.57	1.90	3
L-leucine	0.51	2.01	3
L-phenylalanine	0.87	2.01	14
L-tyrosine	1.4	1.68	26
L-tryptophan	3.0	1.44	25
L-abrine	5.0	1.47	74 ^h

^a Determined as described previously.^{5b} ^b Determined by HPLC analysis using a Daicel Chiracel OD column, eluting with a 60:1 hexane/2-propanol mixture. ^c α -Amino acid solutions were adjusted to pH 4.6–4.9 (glycine pH 5.7). ^d Constant ionic strength (2.00 M) was ensured by adding KNO₃. ^e Conditions: [Cu(NO₃)₂] = [α -amino acid] = 2.4 mM, [**1**] = 0.96 mM, [**2**] = 2.4 mM. ^f Only the results for the major endo (>90%) isomer of the Diels–Alder adduct are shown. ^g Reproducible to within 2%. ^h At 0 °C, pH 6.0 and 1.75 equiv of L-abrine relative to Cu(NO₃)₂.

Figure 1. Transition-state assembly suggested¹⁵ for the aqueous Lewis-acid-catalyzed Diels–Alder reaction of **1** with **2**.

constant even above the value observed for the aquo complex. This implies that an attractive interaction has to be involved, most likely between the aromatic ring of the α -amino acid and the pyridine ring of **1** as depicted in Figure 1. There is literature precedent for this type of interaction in the fields of aqueous coordination chemistry¹¹ as well as enantioselective catalysis.^{9c,d,12}

Surprisingly, the second-order rate constants of the reaction

(11) (a) Brookes, G.; Pettit, L. D. *J. Chem. Soc., Dalton Trans.* **1977**, 1918. (b) Fischer, B. E.; Sigel, H. *J. Am. Chem. Soc.* **1980**, *102*, 2998. (c) Kim, S.; Martin, B. *J. Am. Chem. Soc.* **1984**, *106*, 1707. (d) Aoki, K.; Yamazaki, H. *J. Chem. Soc., Dalton Trans.* **1987**, 2017. (e) Borghesani, G.; Pulidori, F.; Remelli, M.; Purello, R.; Rizzarelli, E. *J. Chem. Soc., Dalton Trans.* **1990**, 2095. (f) Dallavalle, F.; Folesani, G.; Leporati, E.; Galaverna, G. *Helv. Chim. Acta* **1996**, *79*, 1818. (g) Sugimori, T.; Masuda, H.; Ohata, N.; Koiwai, K.; Odani, A.; Yamauchi, O. *Inorg. Chem.* **1997**, *36*, 576 and references therein.

(12) (a) Corey, E. J.; Becker, K. B.; Varma, R. K. *J. Am. Chem. Soc.* **1972**, *94*, 8616. For reviews, see: (b) Sawamura, M.; Ito, Y. *Chem. Rev.* **1992**, *92*, 857. (c) Jones, G. B.; Chapman, B. J. *Synthesis* **1995**, 475.

Table 2. Solvent Effects on the Enantioselectivity of the Diels–Alder Reaction of **1** with **2**, Catalyzed by Cu(L-abrine)⁺ at 0 °C

medium	ee (%) ^{a,b}
acetonitrile ^c	17
THF ^c	24
ethanol ^c	39
chloroform ^c	44
water ^d	74

^a Only the results for the major (>90%) endo isomer of the Diels–Alder adduct are shown. ^b Reproducible to within 2%. ^c Conditions: [Cu(OTf)₂] = 1.00 mM; [L-abrine] = [Et₃N] = 1.75 mM; [**1**] = 10 mM; [**2**] = 80 mM. ^d Conditions: see footnote 10.

are not sensitive to variation of the ligand. Coordination of a strongly Lewis-basic α -amino acid ligand to Cu²⁺ reduces the Lewis acidity of this ion, so that activation of **1** is somewhat less efficient in the presence of α -amino acids, but the effects are small.

Finally, we compared the results obtained in water with the enantioselectivity in organic solvents. Table 2 clearly demonstrates that enantioselectivity benefits considerably from the use of water as the solvent. This is in line with literature observations that arene–arene interactions are less efficient in organic solvents than in water.¹³

Since arene–arene interactions are held responsible for the enantioselectivity in many reactions involving chiral catalysts, we infer that the enhancement of enantioselectivity by water might well be a general phenomena. Moreover, we want to point out that water can be expected to facilitate mechanistic studies of the interactions underlying enantioselectivity. Two effects that usually complicate the study of enantioselective catalysis in organic solvent, ion pairing¹⁴ and clustering of the catalyst species, will normally not occur in water. The remaining rate and equilibrium constants of the different catalytic steps can be determined with relative ease, as is exemplified in this paper.

Acknowledgment. We gratefully acknowledge the Syncom company for the use of the chiral HPLC column and the Research School Netherlands Institute for Catalysis Research for financial support.

JA972772+

(13) (a) Granot, J. *J. Am. Chem. Soc.* **1978**, *100*, 6745. (b) Mitchell, P. R. *J. Chem. Soc., Dalton Trans.* **1980**, 1079. Exceptions to this behavior have also been observed: (c) Sigel, H.; Schneller, K. H.; Haring, U. K.; Malini-Balakrishnan, R. *Inorg. Chim. Acta* **1983**, *79*, 277. (d) Sigel, H.; Tribolet, R.; Scheller, K. H. *Inorg. Chim. Acta* **1985**, *100*, 151.

(14) Evans, D. A.; Murry, J. A.; von Matt, P.; Norcross, R. D.; Miller, S. *J. Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 798.

(15) We are currently investigating whether the coordination sphere of the copper ion is actually trans and whether the dienophile reacts in the s-trans configuration.