University of Groningen

Erratum

Salis, L.; Lof, M.; van Asch, M.; Visser, M. E.

Published in:
Oikos

DOI:
[10.1111/oik.04819](https://doi.org/10.1111/oik.04819)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Salis, L., Lof, M., van Asch, M., & Visser, M. E. (2017). Erratum: Modeling winter moth *Operophtera brumata* egg phenology: Nonlinear effects of temperature and developmental stage on developmental rate. – Oikos 125: 1772–1781. *Oikos*, 126(10), 1522. <https://doi.org/10.1111/oik.04819>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Erratum

Salis, L., Lof, M., van Asch, M. and Visser, M. E. 2016. Modeling winter moth *Operophtera brumata* egg phenology: nonlinear effects of temperature and developmental stage on developmental rate. – Oikos 125: 1772–1781.

1) On page 1774 Eq. 4 is

$$rate(T, D)_{DSD} = \begin{cases} rate(T)_{SS} & , \text{for } D > D_{th} \\ rate(T)_{SS} + (D - D_{th}) * S_T * (rate(T)_{ss} - \rho) & , \text{for } D \geq D_{th} \end{cases} \quad (4)$$

But it should be:

$$rate(T, D)_{DSD} = \begin{cases} rate(T)_{SS} & , \text{for } D < D_{th} \\ rate(T)_{SS} + (D - D_{th}) * S_T * (rate(T)_{ss} - \rho) & , \text{for } D \geq D_{th} \end{cases} \quad (4)$$

2) On page 1777, in the last paragraph of the results, H_H should be substituted with H_L (since H_H was not included in Eq. 3, nor in the list of the parameters).

3) and the rate at T_{ref} (ρ). This parameter caused a large difference in predicted egg-hatching date, compared to the egg-hatching date predicted by the model with the original estimated parameter values (Supplementary material Appendix 1 Fig. A1d). The second most influential parameter was T_{ref} (Supplementary material Appendix 1 Fig. A1g). The parameter that least influenced predicted egg-hatching date was H_H (Supplementary material Appendix 1 Fig. A1f).

Discussion

Timing of winter moth egg-hatching, thus the developmental rate of the eggs, is affected by temperature and develop-

3) In Fig. 2a the line designations are in a reversed order. It should be as below.