University of Groningen

Quantifying the non-breeding provenance of staging Ruffs, *Philomachus pugnax*, using stable isotope analysis of different tissues

Schmaltz, Lucie E.; Loonstra, A. H. Jelle; Wymenga, Eddy; Hobson, Keith A.; Piersma, Theunis

Published in:
Journal of Ornithology

DOI:
[10.1007/s10336-017-1488-x](https://doi.org/10.1007/s10336-017-1488-x)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Schmaltz, L. E., Loonstra, A. H. J., Wymenga, E., Hobson, K. A., & Piersma, T. (2018). Quantifying the non-breeding provenance of staging Ruffs, *Philomachus pugnax*, using stable isotope analysis of different tissues. *Journal of Ornithology*, 159(1), 191-203. <https://doi.org/10.1007/s10336-017-1488-x>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Quantifying the non-breeding provenance of staging Ruffs, *Philomachus pugnax*, using stable isotope analysis of different tissues

Lucie E. Schmaltz¹ · A. H. Jelle Loonstra¹ · Eddy Wymenga² · Keith A. Hobson^{3,4} · Theunis Piersma^{1,5}

Received: 7 April 2017 / Revised: 28 June 2017 / Accepted: 17 August 2017 / Published online: 30 August 2017
© Dt. Ornithologen-Gesellschaft e.V. 2017

Abstract International conservation efforts for migratory populations are most effectively based on quantification of the geographical linkages between wintering, staging, and breeding areas, patterns that may not remain constant in times of global change. We used stable isotope ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$, and $\delta^2\text{H}$) measurements of different tissues representing distinct periods of dietary integration to quantify the non-breeding provenance of a threatened staging population of Ruffs *Philomachus pugnax*. In 199 staging Ruffs captured in 2012 during northward migration in the Netherlands, we compared the multi-isotope patterns of feathers grown at wintering grounds, with the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ profiles of blood cells and plasma representative of staging areas. Few birds had the ^{13}C -depleted and ^{15}N -enriched feathers suggestive of wintering quarters in European agricultural areas. Most Ruffs had higher feather $\delta^{13}\text{C}$ values, suggesting that they wintered in sub-Saharan

Africa. Feather $\delta^2\text{H}$ values were not informative due to the overlap of values from European and African moulting sites. Blood cell $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values indicated that sub-Saharan Ruffs fuelled on low trophic-level foods in habitats dominated by C3 terrestrial or freshwater aquatic primary production, e.g. the rice fields in Africa or the Mediterranean. Stable isotope ratios in plasma suggested that Ruffs made stopovers in southern European agricultural areas. Stable isotopes thus enabled assessments of wintering origin in large numbers of birds. We further propose that conservation measures to protect Ruffs must include the adequate management of sub-Saharan wetlands, based on a better understanding of the role of human-made rice fields for fuelling birds.

Keywords *Philomachus pugnax* · Wintering · Spring migration · Migratory connectivity · Stable isotopes · Shorebirds

Communicated by N. Chernetsov.

✉ Lucie E. Schmaltz
lucie.schmaltz@gmail.com

¹ Conservation Ecology Group, Groningen Institute for Evolutionary Life Sciences (GELIFES), University of Groningen, PO Box 11103, 9700 CC Groningen, The Netherlands

² Altenburg & Wymenga Ecological Consultants, Súderwei 2, 9269 TZ Feanwâlden, The Netherlands

³ Environment Canada, 11 Innovation Blvd., Saskatoon, SK S7N 3H5, Canada

⁴ Department of Biology, University of Western Ontario, London, ON N6A 5B7, Canada

⁵ Department of Coastal Systems, NIOZ Royal Netherlands Institute for Sea Research, Utrecht University, PO Box 59, 1790 AB Den Burg, Texel, The Netherlands

Zusammenfassung

Quantifizierung der Herkunft rastender Kampfläufer *Philomachus pugnax* mit Hilfe der stabilen Isotopen-Analyse von unterschiedlichen Gewebetypen

Internationale Schutzbemühungen für ziehende Populationen sind am effektivsten, wenn sie auf der Quantifizierung der geografischen Verbindungen zwischen Überwinterungs-, Rast- und Brutgebiet basieren, da in Zeiten des globalen Wandels deren Muster nicht konstant bleiben. Mit Hilfe stabiler Isotopen-Messungen ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$ und $\delta^2\text{H}$) aus verschiedenen Gewebetypen, die unterschiedliche Perioden der Nahrungsaufnahme repräsentieren, um die Herkunftsgebiete einer bedrohten Rastpopulation von

Kampfläufern *Philomachus pugnax* zu quantifizieren. 2012 wurden dazu während des Heimzuges 199 rastende Kampfläufer in den Niederlanden gefangen. Wir verglichen die Multi-Isotopen-Muster der Federn, die in den Überwinterungsgebieten gewachsen waren, mit den $\delta^{13}\text{C}$ und $\delta^{15}\text{N}$ Profilen der Blutzellen und Plasma, die typisch sind für die Rastgebiete. Einige Vögel hatten ^{13}C -verminderte und ^{15}N -angereicherte Federn, was auf Überwinterungsgebiete in europäischen Agrargebieten hinweist. Die meisten Kampfläufer hatten höhere $\delta^{13}\text{C}$ Werte in den Federn, was auf eine Überwinterung in Afrika südlich der Sahara hindeutet. Federn mit $\delta^2\text{H}$ Werten lieferten keine Informationen aufgrund der Überlappung der Werte aus Europa und afrikanischen Mausergebieten. $\delta^{13}\text{C}$ und $\delta^{15}\text{N}$ Werte der Blutzellen deuten darauf hin, dass sich südlich der Sahara überwinternde Kampfläufer mit Nahrung niedriger Trophiestufen „auftanken“ in Habitaten, dominiert von C_3 aus terrestrischer oder Süßwasser-Primärproduktion, z. B. Reisfelder in Afrika oder im mediterranen Raum. Die Verhältnisse stabiler Isotope im Plasma legen nahe, dass Kampfläufer in Agrargebieten Südeuropas rasten. Stabile Isotope erlauben daher einen Rückschluss auf die winterlichen Herkunftsgebiete bei einer großen Anzahl von Vögeln. Darüber hinaus sollten Maßnahmen zum Schutz von Kampfläufern ein adäquates Feuchtgebietsmanagement in der Subsahara beinhalten, basierend auf einem besseren Verständnis der Rolle von Reisfeldern für die Fettdeposition für den Zug.

Introduction

Migratory birds occur all over the globe, travelling from one biome to another to take advantage of favourable conditions seasonally (Alerstam 1990; Newton 2008). Along their flyways these widely ranging species experience increasing human pressures on their environment, which often contribute to population declines (Baker et al. 2004; Wilcove and Wikelski 2008; Runge et al. 2015; Piersma et al. 2016). Effective conservation measures should be based on information of migratory connectedness of populations, taking into consideration the full life cycle requirements of species and critical bottlenecks or sites (Webster et al. 2002; Piersma 2007; Taylor and Norris 2010; Iwamura et al. 2013). Unfortunately, such information is often incomplete or lacking for migratory birds (Iwamura et al. 2013; Gilroy et al. 2016; Márquez-Ferrando et al. 2014). This hampers the development of effective demographic monitoring and international conservation strategies (Iwamura et al. 2014). Species using diffuse networks of inland freshwater habitats, or those with poorly described migratory pathways, provide

the greatest challenges for the conservation of wetland-associated species (Skagen et al. 2003).

Numerous tools are currently available to quantify geographic linkages of migrating populations. The use of individual ring recoveries, recaptures, or resightings (Thorup et al. 2014) is presently augmented by the use of modern tracking devices including satellite telemetry (e.g. Gill et al. 2009) and geolocation (e.g. Tøttrup et al. 2012; Ouwehand et al. 2015). For cases where birds move between sites with food webs showing distinct isotopic compositions, a complementary approach has been the use of stable isotope measurements in animal tissues (Hobson 1999, 2005; Hobson and Wassenaar 2008). Interpretation of stable isotope analyses rely on the existence of predictable spatial patterns in isotopic ratios of substrates that vary due to a variety of biogeochemical processes. These processes ultimately shape isotopic “landscapes” or “isoscapescapes”, with such variations being passed on to higher trophic levels (e.g. Catry et al. 2016; Christianen et al. 2017). These isoscapes may serve as a basis to examine retrospectively the geographical origin of sampled animal tissues, hence the origin and movements of organisms themselves (Hobson 1999, 2005; Hobson and Wassenaar 2008; Catry et al. 2016). Isotopic values of tissues incorporated from the environment at the time of growth are maintained until the tissue is renewed or replaced. Based on a careful choice of tissues, stable isotope analyses provide a complementary and cost-effective method to monitor migratory linkages over time (Hobson and Wassenaar 2008; Dietz et al. 2010; Yerkes et al. 2008).

Here we used stable isotope measurements to quantify the current geographical linkages of an endangered staging population of Ruffs, *Philomachus pugnax*, migrating along the East Atlantic Flyway (Verkuil et al. 2012; Schmaltz et al. 2015). Ruffs are lek-breeding shorebirds, well known for their strong sexual dimorphism and the extravagant nuptial plumage of males (van Rhijn 1991). The species is strongly associated with inland wetlands habitats, whose migration typically takes place on a broad front between northern Eurasia where they breed, and Africa and India where they winter (Cramp and Simmons 1983). Ruffs along the East Atlantic Flyway stage in the Netherlands during spring migration then to move on to breeding areas at the sub-arctic and arctic latitudes of Eurasia (Jukema et al. 2001), before returning to the floodplains of the Senegal Delta, Inner Niger Delta in Mali and Chad Basin in winter (Zwarts et al. 2009). A few thousand males, however, do not cross the Sahara and winter instead in northwest Europe (Prater 1973; Castelijn 1994; Gill et al. 1995; Devos et al. 2012; Hornman et al. 2013), Iberia (Hortas and Masero 2012), and Morocco (Qninba et al. 2006—Fig. 1). Jukema et al. (2001) suggested that European winterers are the earliest arriving birds at the Dutch staging areas in spring, followed by sub-Saharan wintering males and late females.

Fig. 1 **a** Distribution range of Ruffs with the location of the collection sites of tissue samples. The wintering areas of Europe are displayed in *blue*, wintering range in sub-Saharan Africa is in *green* and the breeding range of the species in *yellow*. *Dashed arrows* indicate the East Atlantic Flyway corridor. Staging birds were captured on their main spring staging area in southwest Friesland, the

Netherlands. One wintering female was sampled in Mopti, Mali. **b** Annual cycle of Ruffs wintering in sub-Saharan Africa and in Europe, with respective time window of isotopic integration for plasma, blood cells, and feathers, relative to the period of spring capture on the staging site in Friesland (indicated by the *red stripe*). *Colour code* is the same as for **a** (colour figure online)

The staging population in spring in the Netherlands has shown a severe decline since the late twentieth century. Of the 20,000 birds counted during peaks of spring migration on communal roosts in the 1990s, fewer than 5000 remained 15 years later (Verkuil et al. 2012; Schmaltz et al. 2015). The decline may at least partially be explained by a redistribution towards Eastern Europe (Rakhimberdiev et al. 2010; Verkuil et al. 2012). In parallel, however, low female survival, perhaps linked to greater exposure to heavy hunting pressures in Mali than males, could have severe negative consequences for this population (Schmaltz et al. 2015). In sub-Saharan Africa, the control of the hydrology of the Senegal River, and partly also the Niger River, enabled the development of irrigated agriculture, but greatly disrupted the natural dynamics of the floodplains on which Ruffs and many other wintering birds relied on traditionally (Zwarts et al. 2009). Within continental Europe and the Mediterranean, the anthropogenic pressures on natural wetlands continues along with associated intensive land-use for agriculture. Birds are more and more exposed to man-made wetlands, such as ricefields north and south of the Sahara that might constitute at least temporarily, favourable foraging habitat for numbers of waterbirds (Wymenga and Zwarts 2010; Sánchez-Guzmán et al. 2007).

In the context of landscape changes occurring along the entire flyway, we assessed the current wintering quarters and

northward migratory patterns of Ruffs using the Netherlands as their main staging area. We inferred wintering quarters of Ruffs from multi-isotopic values ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$, $\delta^2\text{H}$) in their 9th primary feather, which is replaced during early winter on their final wintering destination (Pearson 1981; Koopman 1986; OAG Münster 1998). As a metabolically inert tissue, feathers “lock in” the spatial isotopic signals at time of growth (Hobson and Clark, 1992; Bearhop et al. 2002). Unlike feathers, blood is continuously renewed, and in medium-sized shorebirds like Ruffs a ^{13}C turnover rate will be about 2 weeks for plasma to a month for blood cells (Evans Ogden et al. 2004; Klaassen et al. 2010). This way, we inferred current and recent refuelling areas for arriving Ruffs in the Netherlands by comparing $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ measurements in plasma and blood cells separately. Using these results, we examined the proposed tendency for European wintering birds to arrive at the Dutch staging area before the birds from sub-Saharan wintering areas.

Materials and methods

Field methods

Ruffs were captured during spring migration in 2012 at our study site, the core of their staging area in the province of

Friesland in the north of the Netherlands. The area consists of agricultural grasslands intensively managed for dairy farming along the shore of Lake IJsselmeer between the villages of Makkum (53°3.37'N, 05°24.19'E) in the north, Laaksum (52°51.15'N 05°24.77'E) in the south, and It Heidenskip (52°56.93'N, 05°30.11'E) in the east (Schmaltz et al. 2016). During their stay, Ruffs feed on soil invertebrates and above-ground insects in the grasslands, while they rest along shorelines and scattered inland wetlands during midday and at night (Verkuil and de Goeij 2003; Schmaltz et al. 2016; Onrust et al. 2017). Between 10 March and 15 May 2012, 199 adult Ruffs (187 males and 12 females) were caught using traditional clap nets (“wilsternets”, Jukema et al. 2001). The low numbers of females in our sample is explained by their rarity on this staging area, where there is a striking male-bias (Jukema et al. 2001; Schmaltz et al. 2015). Birds were measured, weighed (± 1 g), and banded with a metal band and an individual combination of four color bands and one leg flag. Sexing was based on wing length (Prater et al. 1977; Jukema and Piersma 2006). Age was determined on the basis of leg colour (orange for adults, greenish for first-year birds), the presence or absence of juvenile inner median coverts (Prater et al. 1977), and also by examining breeding plumage development (Meissner and Scebba 2005; Karlionova et al. 2008). For stable isotope analyses, we collected feather material for all birds captured and blood for 173 of them (163 males and 10 females), which were then used for stable isotope analysis.

To infer the winter origin of adult staging Ruffs, we measured $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, and $\delta^2\text{H}$ values in the vane of the 9th primary. For each bird, we clipped 2 cm of the inner vane at the feather's base using small scissors. Samples were then stored in paper envelopes until analysis. We did not consider young birds, as only a small proportion of them replace the outermost primaries in their first winter (Melter and Sauvage 1997). In parallel, we also retrieved samples of 9th primary (i.e. similar as above) of known origin: eight from Ruffs captured in November and December 2004 and 2005 in the Netherlands, and one from an adult female sold for consumption in the market of Mopti in Mali (14°29'45 N, 04°11'55 W) in February 2012.

After ringing, we took a blood sample (200 μL) by puncturing the wing vein and drawing blood into heparinized capillaries. Within 3 h after collection, blood samples were centrifuged for 12 min at 6900g in order to separate plasma and blood cells. Until freeze drying, plasma and blood cell samples were stored in glass vials in a freezer (-20°C).

Stable isotope analysis

Proximal feather vane material was washed in a 2:1 chloroform:methanol solution and dried.

For $\delta^2\text{H}_\text{f}$ analysis, dried washed feather material was weighed (c. 0.35 mg) into silver capsules and combusted under helium flow in a Hekatek furnace at 1350°C interfaced with a Eurovector 3000 (Milan, Italy—www.eurovector.it) elemental analyser. The resultant H_2 gas was measured for $\delta^2\text{H}$ in an Isoprime (Crewe, UK) continuous flow stable isotope mass spectrometer and corrected for H exchange using the comparative equilibration technique of Wassenaar and Hobson (2003), using three keratin calibrated standards: Caribou Hoof Standard (CBS -197‰), Commercial Keratin Standard (SPK -121.6‰), and Kudu Horn Standard (KHS -54‰). All measurements are reported in δ -notation as the non-exchangeable feather H component in parts per thousand (‰) relative to the Vienna Standard Mean Ocean Water (VSMOW)—Standard Light Antarctic Precipitation (SLAP) scale. Based on replicate within-run measurements of standards ($n = 5$ per run) measurement error was estimated to be $\pm 2\text{‰}$.

For $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analyses, between 0.5 and 1.0 mg of feather or dried blood material was weighed into tin capsules and combusted online using a Eurovector 3000 (Milan, Italy—www.eurovector.it) elemental analyser. The resulting CO_2 and N_2 was introduced into a Nu Horizon (Nu Instruments, Wrexham, UK—www.nu-ins.com) triple-collector isotope-ratio mass-spectrometer via an open split and compared to CO_2 or N_2 reference gas. Stable nitrogen ($^{15}\text{N}/^{14}\text{N}$) and carbon ($^{13}\text{C}/^{12}\text{C}$) isotope ratios were expressed in δ notation, as parts per thousand (‰) deviation from the primary standards, atmospheric AIR and Vienna Pee Dee Belemnite (VPDB). Using previously calibrated internal laboratory standards (powdered keratin BWB II: $\delta^{13}\text{C} = -20.0\text{‰}$, $\delta^{15}\text{N} = -14.1\text{‰}$ and gelatine: $\delta^{13}\text{C} = -13.6\text{‰}$, $\delta^{15}\text{N} = -4.7\text{‰}$) within run ($n = 5$) precision for $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ measurements was $\pm 0.15\text{‰}$.

Assigning wintering origin

We lack sufficient ground-truth samples to enable assignment by means of a discriminant function analysis. Therefore, we chose to follow the approach of Yerkes et al. (2008) to establish feather isotopic thresholds (cut-off values) to delineate wintering biomes of staging Ruffs. Isotopic thresholds were set on the basis of: (1) the relative difference in the abundance of plants using C3 and C4 photosynthesis pathways in Europe versus sub-Saharan Africa which would respectively show ^{13}C -depleted and enriched tissue values (mean C3 plant $\delta^{13}\text{C} -28\text{‰}$, C4 plant $\delta^{13}\text{C} -13\text{‰}$, Peterson and Fry 1987), (2) on the land-use practices with high nitrogen input (manure) increasing $\delta^{15}\text{N}$ values (Hebert and Wassenaar 2001, 2005; Yerkes et al. 2008) and (3) the negative gradient in $\delta^2\text{H}$ feather values existing from sub-Saharan West Africa towards northwestern Europe (Bowen et al. 2005). Samples of

known moulting locations were then used to assess the relevance of our inferences.

First, we expected that European winterers would show lower $\delta^{13}\text{C}$ feather values. Most of these Ruffs winter in open agricultural areas and replace their 9th primary in early September (Koopman 1986) while they forage on soil invertebrates and spilled grain found in meadows or arable lands (Castelijns 1994) with C3 crops. In contrast, feathers grown in sub-Saharan Africa should present higher $\delta^{13}\text{C}$ values. Sub-Saharan Ruffs moult primary 9 in early October for males to a month later for the delayed females (Pearson 1981; Koopman 1986; OAG Münster 1998). During this time, Ruffs rely on food based on a C4-dominated environment, invertebrates, but also grass seeds (e.g. *Echinochloa* sp., *Panicum* sp.) found along the edges of the floodplains, which are then at their highest levels after the rains of June and July (Tréca 1990, 1994; van der Kamp et al. 2002a). Eventually, we expect feather $\delta^{13}\text{C}$ values to reflect a broad C3 and C4 isotopic range from -25.8 to -10.8‰ taking into account a diet feather discrimination factor of 2.16‰ (Caut et al. 2009); see also Werner et al. 2016). We considered the median threshold $\delta^{13}\text{C}$ value of -18.3‰ to distinguish birds that fed on a C3 vs. C4 dominated diet at time of moult. The presence of cultivated maize (*Zea mays*, a C4 plant) in Europe and the presence of Rice (*Oryza* sp., a C3 plant) in sub-Saharan Africa should not confound our expectations. By the time Ruffs are replacing primary 9, neither the rice in West Africa nor the maize in Europe is harvested and available for Ruffs.

We assumed that $\delta^{15}\text{N}$ feather values higher than 10‰ were indicative of birds feeding in agricultural areas subjected to high nitrogen input. In Europe, intensive agricultural practices are common with regular application of manure on lands. Animal waste, as an additional source of nitrogen, may have typically elevated $\delta^{15}\text{N}$ values (from 10 to 25‰ compared to synthetic fertilizer (-4 to $+7\text{‰}$; Kendall 1998), which would be reflected in feather $\delta^{15}\text{N}$ values (e.g. Hebert and Wassenaar 2001, 2005; Yerkes et al. 2008; Coulton et al. 2010). In comparison, land use in sub-Saharan Africa is extensive, but locally Ruffs may use pastures with high livestock densities. This would make it possible to encounter sub-Saharan individuals showing elevated feather $\delta^{15}\text{N}$ values too.

Eventually, in the case of feathers showing low $\delta^{13}\text{C}$ and low $\delta^{15}\text{N}$ values, we assumed that birds used natural or extensive terrestrial or freshwater C3 habitats (i.e. not subjected to input of manure) while feeding on low trophic levels, as would indicate low $\delta^{15}\text{N}$ values (Post 2002). Because Ruffs are associated primarily with freshwater (Cramp and Simmons 1983), this corresponds to natural freshwater wetland areas in Europe or perhaps ricefields. However, considering that rice cultivation is widespread in Africa and the high heterogeneity in farming practices

there, we cannot rule out the use of non-harvested rice fields during the period of moult. To be conservative, we did not predict a specific geographical origin for these birds.

We thus ended up with four possible isotopic environments reflecting a bird's early wintering location and habitat (Fig. 2): (A) European agricultural areas with high N input with C3 foods (i.e. $\delta^{13}\text{C} < -18.3\text{‰}$, $\delta^{15}\text{N} > 10\text{‰}$), (B) Natural wetlands or rice fields in unknown wintering quarters ($\delta^{13}\text{C} < -18.3\text{‰}$, $\delta^{15}\text{N} < 10\text{‰}$), (C) sub-Saharan floodplains with C4 foods and areas grazed by livestock ($\delta^{13}\text{C} > -18.3\text{‰}$, $\delta^{15}\text{N} > 10\text{‰}$), (D) sub-Saharan floodplains ($\delta^{13}\text{C} > -18.3\text{‰}$, $\delta^{15}\text{N} < 10\text{‰}$). We fully recognize that these cutoffs are somewhat arbitrary and that birds can be misplaced using these criteria. However, this approach was the most parsimonious given the data.

We chose to not set an isotopic threshold for $\delta^2\text{H}$ and kept a more explorative approach. This is because Ruffs use freshwater wetlands and agricultural areas that are subjected to many processes and anthropogenic activities (e.g. intense evaporation, use of water of mixed origin, and ground waters) leading to large variation in the incorporation of deuterium into the diet and then in animal tissues (Bowen et al. 2005; Oppel et al. 2011; Hobson et al. 2012; Gutiérrez-Esposito et al. 2015). Nevertheless, at time of growth of primary 9, we expected to find a tendency for higher $\delta^2\text{H}$ values in feathers grown in sub-Saharan than in feathers grown in Europe (Bowen et al. 2005). In sub-Saharan Africa, water feeding the floodplains of Senegal and Mali may importantly originate from May to July rains in the Guinean Highlands, which $\delta^2\text{H}$ values range from -22 to 10‰ (SD ranged from 9.6 to 12.8‰) according to $\delta^2\text{H}$ monthly mean values from the Global Network of Isotopes in Precipitation (GNIP) database administrated by the International Atomic Energy Association and World Meteorological Organization (IAEA/WMO 2001). In comparison, the $\delta^2\text{H}$ values of September precipitation over Western Europe may range from -54 to -22‰ (SD ranged from 8 to 9.6‰). Finally, we examined the contribution of deuterium to the observed variation in feather $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ measurements with principal component analysis (PCA). This approach allowed us to incorporate our assignment criteria based on $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ feather values (i.e. isotopic environment A, B, C, D as supplementary qualitative variables), while taking into account the part of the overall observed variation explained by $\delta^2\text{H}$ measures in feathers.

Migration stages during northward migration

We isotopically characterized the staging grounds of Ruffs used 2 weeks to a month before their arrival on our study site in the Netherlands from $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ measures in

Fig. 2 **a** Expected wintering isotopic environments for spring migrant Ruffs based on $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ thresholds values in feathers, **b** primary feathers $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ reflecting isotopes ratios of moulting areas in either sub-Saharan Africa or Europe. *Filled* and *open dots*, respectively, represent samples from males and females captured on the staging area in Friesland and for which wintering origin in unknown. *Filled triangles* (in blue) represent samples from males at known wintering site in the Netherlands, the *filled triangle* (in green) represent a sample from a female collected on the local food market in Mopti, Mali. **c** Principal Component (PC) biplot of $\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and $\delta^2\text{H}$ measurements in feather samples from staging Ruffs collected in Friesland during spring migration 2012. Ellipses grouped 67% of the individuals assigned to each isotopic environment A, B, C, and D (see **a**)—here as qualitative supplementary variables. PC1 and PC2 refer to the principal component axes scores on the *first* and *second axes*, respectively. The amount of variation accounted for by the *axes* are shown in *brackets* (colour figure online)

carbon, we considered an isotopic threshold value of $\delta^{13}\text{C} = -20.4$ ‰, which corresponds to the median value of C3 versus C4 dominated diet ($\delta^{13}\text{C} = -20.5$ ‰) adjusted with a carbon diet-whole blood discrimination factor of 0.09‰ ($\Delta^{13}\text{C}_{\text{diet-whole blood}} = -0.199 \times \delta^{13}\text{C}_{\text{diet}} - 3.986$ ‰, Caut et al. 2009). For nitrogen, we considered a threshold value of $\delta^{15}\text{N} = 8.4$ ‰, which is the nitrogen threshold value for feather ($\delta^{15}\text{N} = 10$ ‰) to which we subtracted the difference of 1.6‰ between nitrogen diet-feather and nitrogen diet-whole blood discrimination factors ($\Delta^{15}\text{N}_{\text{diet-whole blood}} = 2.25$ ‰, $\Delta^{15}\text{N}_{\text{diet-feather}} = 3.84$ ‰, Caut et al. 2009).

We kept the same rationale as in feathers but took into consideration that possible isotopic environments (Group A, B, C, D) reflected the birds' early spring locations and habitats. For isotopic environment B ($\delta^{13}\text{C} < -20.4$ ‰, $\delta^{15}\text{N} < 8.4$ ‰) this implies that the use of rice fields becomes more likely. In Africa, Ruffs can feed on spilled kernels in dry harvested ricefields (Tréca 1990), whereas in Europe Ruffs may benefit from the increasing tendency for post-harvest inundation of rice fields (Hortas and Masero 2012; Pernollet et al. 2015), which follows the first spring rains, and later the flooding of rice paddies for the new sowing (Wymenga 1999; Bacetti et al. 1998). After a change in isotope values of diet, tissues follow an exponential pattern of change (Karasov and Martinez del Rio 2007). We assumed a half-life of 6 and 15 days for plasma and blood cells, for carbon and nitrogen stable isotopes (see above). Ruffs, on average, stage 19–23 days in the Netherlands (Verkuil et al. 2010), which in principle allowed for enough time (i.e. three half-lives) for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ plasma values of migrants to reach equilibrium levels on the local Dutch diet and for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ blood cell values to nearly approach it.

Accordingly, we examined the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of blood cells and plasma of Ruffs caught before 1 April.

plasma and blood cells. We assumed a half-life of 6 and 15 days for plasma and blood cells, respectively, as determined isotopically in a similarly sized shorebird, the red knot *Calidris canutus* for carbon stable isotopes (Klaassen et al. 2010). We considered that incorporation rate to be equivalent for the nitrogen stable isotope (van Gils and Ahmedou Salem 2015; but see Dietz et al. 2013). Then we used the isotopic thresholds set above for feathers, which we adjusted for blood cells and plasma tissues. For

Capture date should be close to true arrival date (Verkuil et al. 2010), and their blood isotope values likely reflect more of the isotopic environment of any previous staging area. From the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ plasma values of these early birds, we verified the proposition that some of the Ruffs make a nonstop flight across the Sahara and Europe to the Netherlands (Jukema et al. 2001). Later, we considered Ruffs caught during the entire season to explore temporal patterns of changes in plasma and blood cell $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values. We calculated log-linear time trends of changes in plasma and blood cell $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in Ruffs assigned to different wintering areas. Females were considered aside of males because of their delayed migration.

For birds wintering in European agricultural areas (A in Fig. 2a), we considered that plasma and blood cells isotopic values of these European Ruffs caught after 20 April were equivalent to the end points of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in plasma and blood cells at equilibrium with the local diet.

All statistical tests involved were performed in R version 3.3.1 (R Development Core team 2016).

Results

Wintering origin

Values of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in feathers of known origin were in agreement with the independently set isotopic thresholds to discern wintering areas (Fig. 2). The $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in feathers grown in the Netherlands (mean $\pm$ SD, $\delta^{13}\text{C}$, $-23.5 \pm 1.68\text{‰}$; $\delta^{15}\text{N}$, $13.0 \pm 1.35\text{‰}$), confirmed our expectation for an agricultural wintering habitat with high nitrogen inputs dominated by C3 foods (A: $\delta^{13}\text{C} < -18.3\text{‰}$, $\delta^{15}\text{N} > 10\text{‰}$; see Fig. 2a, b). The isotope values of the one feather from a female wintering in Mali ($\delta^{13}\text{C} = -10.3\text{‰}$; $\delta^{15}\text{N} = 11.7\text{‰}$), were consistent with the suggestion that feathers grown in West Africa may present $\delta^{13}\text{C}$ values close to those expected for a diet based on C4 plants, presumably an important food resource for Ruffs there.

Among the 199 Ruffs sampled during their stage in the Netherlands, 75% ($n = 149$) showed feather isotopic values consistent with those assumed to have a sub-Saharan wintering origin (Fig. 2a, c, d: $\delta^{13}\text{C} > -18.3\text{‰}$), of which 129 (given $\delta^{15}\text{N}$ feather values $> 10\text{‰}$ —Fig. 2b) may have used areas grazed by cattle and/or feeding on higher trophic level foods. All females in the sample from Friesland were assigned to have wintered in sub-Saharan Africa (Fig. 2b). In contrast, only 29 male Ruffs (15%) seemed to have wintered in agricultural areas in Europe (A: $\delta^{13}\text{C} < -18.3\text{‰}$, $\delta^{15}\text{N} > 10\text{‰}$; Fig. 2b), while 21 other males showed isotopic values with both depleted values in ^{13}C and in ^{15}N , probably indicating moulting areas in

European freshwater wetlands or perhaps also rice fields, possibly in Africa (Chi square test: $\chi^2 = 169.3$, $df = 3$, $p < 0.001$). The $\delta^2\text{H}$ values in feathers assigned to moulting areas in Europe (mean $\pm$ SD, $-33.9 \pm 24.4\text{‰}$, $n = 29$) and in sub-Saharan ($-40.6 \pm 15.0\text{‰}$, $n = 149$) both showed large variations and did not differ from each other (Mann–Whitney test, $W = 2619$, $P = 0.07$).

From the PCA analysis, the first two principal components (eigenvalues > 1) accounted for 38 and 34% of the observed variations in feather isotopic values. The $\delta^{13}\text{C}$ and $\delta^2\text{H}$ feather values contributed the most (52 and 30%) to the first principal component (PC1), and were negatively correlated. Second principal component (PC2) was best described by $\delta^{15}\text{N}$ (61%) and then by $\delta^2\text{H}$ feather values (39%), in both cases positively.

Spring staging grounds before arrival in Friesland

Among the 29 Ruffs assigned to European wintering quarters (group A), five birds were caught after the 20th of April, from which we took the average $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in blood cells and plasma to predict the expected endpoint values at equilibrium with the diet on the Dutch staging area as $\delta^{13}\text{C}$: $-25.7 \pm 1.2\text{‰}$, $\delta^{15}\text{N}$: $12.3 \pm 0.6\text{‰}$ for blood cells and $\delta^{13}\text{C}$: $-26.7 \pm 0.9\text{‰}$, $\delta^{15}\text{N}$: $12.2 \pm 0.7\text{‰}$ for plasma (mean $\pm$ SD, $n = 5$).

Out of all birds sampled, 55 Ruffs were captured before 1 April. Five were allocated to a European origin (9%), 42 (76%) to sub-Saharan Africa (38 in isotopic environment C, including one female and four in isotopic environment D), and eight birds (15%) could not be assigned to a particular location (isotopic environment B). The proportion of European wintering Ruffs in early catches did not differ from their proportion in overall catches [binomial test, $p = 0.26$, confidence interval (0.3–20%)].

Among the 41 early males that wintered in Africa, all had $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ plasma values indicating the use of agricultural areas in Europe as early as 2 weeks before arrival on the study site (Chi squared test, $\chi^2 = 123$, $df = 3$, $p < 0.001$) and so did the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ blood cells values for 30 of them, as early as a month before arrival in Friesland (73%, Chi squared test, $\chi^2 = 56.66$, $df = 3$, $p < 0.001$, Fig. 3). However, for 10 birds (24%), the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ blood cell values were more contrasted to plasma values, falling in isotopic environment B and indicating the use of natural freshwater wetland areas in Europe or perhaps also rice fields (Fig. 3). A last male presented blood cell $\delta^{13}\text{C}$ values indicative of C4 based diet (Group C), suggesting his presence in Africa in the month before his capture (Fig. 3). The single early arriving female showed plasma and blood cells isotope profiles indicative of agricultural habitat in Europe (Fig. 3a).

Over the entire passage period, all but one Ruff assigned to sub-Saharan Africa and of unknown wintering quarters (groups B, C, and D) presented $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in plasma already converged towards the local end point values for plasma (Fig. 4). This male caught on April 20 (plasma: $\delta^{13}\text{C}$ -18.5‰ and $\delta^{15}\text{N}$ 14.6‰ , i.e. group C) provide the unique suggestion of a bird present in Africa within 2 weeks before his capture in the Netherlands.

The $\delta^{15}\text{N}$ blood cell values of males assigned to Africa or unknown wintering quarters (groups B, C, and D), increased over time (Fig. 4, B: $F_{1-7} = 13.75$, $p = 0.02$, C & D: $F_{1-118} = 41.2$, $p \leq 0.001$) converging towards $\delta^{15}\text{N}$ end point values expected in the Netherlands (Fig. 4) while it was not the case for blood cells $\delta^{13}\text{C}$ values, all below the carbon threshold of -20.4‰ and even decreasing for sub-Saharan males (Fig. 4C, D: $F_{1-118} = 19.99$,

$p < 0.001$). There was no such patterns for Ruffs that supposedly wintered in Europe (A: $F_{1-22} = 0.26$, $p = 0.61$). Finally, for all male Ruffs, we also observed a slight decrease in plasma $\delta^{13}\text{C}$ values over the season, (Fig. 4A: $F_{1-22} = 6.91$, $p = 0.02$, B: $F_{1-7} = 4.66$, $p = 0.05$, C, D: $F_{1-118} = 23.95$, $p < 0.001$).

As for females, two individuals captured on May 1 presented $\delta^{13}\text{C}$ blood cells values indicative of C4 based diet (Fig. 4— $\delta^{13}\text{C}_{\text{blood cells}}$ of -18.3 and -19.6‰) and their presence in Africa in the last month. The eight other females had $\delta^{13}\text{C}$ blood cells values already converged within expected values of the Netherlands.

Discussion

Quantitative information on the migratory connectivity of endangered populations can establish the contemporary importance of critical sites or habitats, and prioritize management actions and conservation decisions. Our study shows that in 2012, the remnant staging population of Ruffs in the Netherlands consisted mainly of individuals wintering in sub-Saharan Africa, but nevertheless involved a considerable proportion of the small population of male Ruffs wintering in Europe. For Ruffs wintering south of the Sahara, rice fields either in Africa or in the Mediterranean are likely to represent important fuelling areas. We found no evidence that Ruffs routinely make non-stop flights from West Africa to the staging site in the Netherlands. Instead, our results suggested that prior to their arrival in Friesland, Ruffs usually stopover in more southerly intensive agricultural areas.

The multi-isotopic values of winter-grown feathers from Ruffs staging in spring in the Netherlands showed differences between European and sub-Saharan wintering quarters although there was high variability (Fig. 2b, c). Intensive agricultural areas of Europe were satisfactorily delineated by the high $\delta^{15}\text{N}$ threshold values characteristic of regular manure input on these lands. Above the $\delta^{13}\text{C}$ threshold of -18.3‰ , indicating a sub-Saharan origin, feathers showed high isotopic variability suggestive of the opportunistic feeding behaviour of Ruffs and/or the isotopic complexity of sub-Saharan freshwater wetlands (see Oppel et al. 2011). Our data indicate that during the growth of primary 9, sub-Saharan Ruffs occupied habitat dominated by C4 grasses, possibly at the interface between inundated zone and more xeric habitat given a relatively large range of feather $\delta^{13}\text{C}$ values (e.g. Marra et al. 1998). The mostly enriched $\delta^{15}\text{N}$ feather values may also reflect the influence of livestock. This could suggest wintering quarters in and near the Inner Niger Delta where livestock is an important resource and where Ruffs are abundant and could forage in floodplains habitats subject to grazing (Wymenga et al. 2002; Zwarts et al. 2009). Note that Ruffs

Fig. 3 a Blood cells and plasma $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ (‰) of staging Ruffs caught before April 1st 2012 in southwest Friesland that were previously assigned to sub-Saharan wintering origin and mean values (error bars represent standard errors). These are all males, except for one female designated with open symbols. Dashed lines are indicative of isotopic thresholds for $\delta^{13}\text{C}$ (-20.4‰) and $\delta^{15}\text{N}$ (8.4‰) for blood cells and plasma tissues b Proportion of sub-Saharan Ruffs with blood cells and plasma isotopic signatures reflecting isotopic environments A, B, C, or D with 95% CI

Fig. 4 Temporal changes in the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values (‰) of blood cells (dark red circles) and plasma (in yellow triangles) of male ruffs in the course of the 2012 spring migration season in southwest Friesland. Only significant temporal trends are emphasized with a trend line and a 95% confidence contour. The $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values (‰) of blood cells and plasma of females assigned to wintering quarters in sub-Saharan Africa, are represented by open dark red dots and open yellow triangles, respectively. Dashed lines are indicative of isotopic thresholds for $\delta^{13}\text{C}$ (−20.4‰) and $\delta^{15}\text{N}$ (8.4‰) for blood cells and plasma tissues (colour figure online)

have practically vanished from the Senegal River delta (Triplet et al. 2014). In contrast, the $\delta^2\text{H}$ values of feathers presumably grown in Europe and those grown in sub-Saharan Africa did not differ.

Globally, our study provides new evidence that the assignment to wintering areas is challenging for ground-foraging species closely associated with freshwater wetlands, and notably tropical wetlands subjected to intense

evaporation (Oppel et al. 2011; Hobson et al. 2012). Our study needs to be further augmented by more ground-truth samples from Europe and sub-Saharan wintering quarters, and ideally combine multi-isotopic data with movement data from satellite tags. If isotopic values in tissues with crucial information on individual habitat use during the winter time could be linked, we could evaluate the feasibility of multi-isotopic large-scale geographical assignments for species strongly associated to wetlands (López-Calderón et al. 2017).

Despite these limitations, this study does provide important insights on the geographical aspects of the itineraries of Ruffs staging in spring in The Netherlands. The dairy farmland of the province of Friesland currently hosts mostly sub-Saharan wintering Ruffs. As the ca. 15% of the males staging in Friesland which had wintered in Europe is much more than the few percent expected on the basis of the ratio of wintering numbers (a few thousand in Europe; Gill et al. 1995; Qninba et al. 2006; Hortas and Masero 2012; Devos et al. 2012; Hornman et al. 2013) and hundreds of thousands in West Africa (Zwarts et al. 2009), Friesland appears especially important for these nearby wintering Ruffs.

Jukema et al. (2001) suggested that Ruffs wintering in Europe arrived earlier than sub-Saharan winterers on the staging site in Friesland. We found no evidence for this. Note, however, that Ruff numbers in spring 2012 represent less than 10% of the numbers observed by Jukema et al. (2001) in 1998 and, therefore, the whole context may be now very different. In particular, Jukema et al. (2001) suggested that European males arrived on the study site by late February, followed by the bigger flocks of African wintering Ruffs coming in by mid-March. Now, the first large flocks of Ruffs in Friesland are rarely observed before 10 March. It could thus be that the early passing European Ruffs are no more. Nevertheless, the Friesland staging area may remain a critical site for a perhaps currently restricted population of males wintering in northwest Europe.

In West Africa, the body mass of Ruffs at departure for spring migration should enable them to reach the Netherlands in a single non-stop flight (OAG Münster 1998). However, unlike red knots reaching the Wadden Sea (the Netherlands), their main stopover area in one non-stop flight of similar length from the Nearctic breeding grounds (Dietz et al. 2010), we found no indication of Ruffs landing on the Frisian staging site directly from sub-Saharan African origins. Only one bird, caught on 20 April, may have arrived non-stop from its sub-Saharan wintering quarters. Instead, $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ plasma values of Ruffs from West Africa suggest that Ruffs stopover in other intensive agricultural areas just before reaching Friesland (Figs. 3, 4). This is perhaps from elsewhere in the Netherlands or in neighbouring countries such as England

and Belgium where Ruffs are also found during the winter. It will be important to identify those intermediate staging areas.

Indeed, all sub-Saharan wintering Ruffs newly arrived on the staging site (i.e. caught early) may have used intensive agricultural areas for 2 weeks, 73% for a month (Fig. 3). However, interestingly, the remaining birds showed more contrasting blood cells values, reflecting the use of habitats dominated by C3 terrestrial or freshwater aquatic primary production, within a month before they arrived in the Netherlands (Fig. 3a, b, group B). This either suggests previous intermediate stopovers in natural wetlands of Europe, where C3 plants dominate, or perhaps in rice fields of the Mediterranean or the Sahel. In the course of the migratory period, the global increase in $\delta^{15}\text{N}$ values in blood cells of Sub-Saharan Ruffs while $\delta^{13}\text{C}$ values remained low (Fig. 4) also suggests habitat switching from natural or man-made wetlands to intensive agricultural areas of Europe. During the pre-fattening period in Africa, Ruffs usually fuel on invertebrates along the receding water line of the floodplains (van der Kamp et al. 2002b; Zwarts et al. 2009), but there were also evidence of Ruffs fuelling on rice kernels. Tréca (1994) showed that from December onwards, the stomach of Ruffs caught in Senegal were filled almost exclusively with rice. In parallel, in Europe, January censuses in of southern provinces of Spain interestingly reported increasing numbers of Ruffs since the 1990s for Cataluña and since 2002 for Extremadura, highlighting the importance of rice fields for the species (Hortas and Masero 2012). Several resightings of Dutch colour-ringed birds in rice field areas in Spain, Portugal, and Italy in early spring (L.E. Schmaltz, unpublished data) eventually confirm the existence of early spring migratory connections between the Mediterranean and Dutch agricultural grasslands.

To conclude, despite the evident isotopic complexity of wetland habitats and habitats strongly modified by anthropogenic activities (Hebert and Wassenaar 2001; Oppel et al. 2011) and the opportunistic feeding behaviour of Ruffs, the use of isotopic threshold values helped us to obtain fair estimates of the non-breeding provenance and the use of spring staging habitats of the remnant population of Ruffs staging in Friesland in spring 2012. Most birds wintered in the sub-Saharan floodplains, while the dairy farmland of Friesland is a particularly important staging area for male Ruffs wintering in Europe. Sub-Saharan wintering Ruffs appear to use in sequence different habitats. Their spring migratory itineraries may comprise the use of ricefields in the Sahel, intermediate stopover(s) in ricefields and freshwater wetlands in Europe, before reaching intensive agricultural areas of Europe and only later the staging site in Friesland. Ruffs use heavily anthropogenic habitats during the non-breeding season,

among which rice fields may be important during migration, which calls for new attention on the use of these habitats north and south of the Sahara, and on the importance of the threatened natural dynamics of the floodplains in the Sahel. The combination of more ground-thruted samples and movement data from satellite tags could tell in the future whether our insights are correct.

Acknowledgements We thank the Frisian “wilsternetters” Albert Anne Mulder, Doede Anne Mulder, Eeltje Anne Mulder, Fons Baarsma, Jappie Boersma, Cees Dekker, Piet Feenstra, Albert Hendrik Mulder, Doede Hendrik Mulder, Douwe de Jager, Bauke de Jong, Joop Jukema, Bauke Kuipers, Willem Louwsma, Catharinus Monkel, Rein Mulder, Jaap Strikwerda, Fokke Tuinstra, Bram van der Veen, F. van der Veen, Arend Veenstra, Sierd Visser, Willem Visser, Piet Vlas, Douwe van der Zee, and Rinkje van der Zee for their dedication to catch Ruffs since 2004. We thank Marta Lomas Vega and Helena Bathala for their help in the field and Idrissa Ndiaye for his precious insights from the field in Senegal. This study was carried out under license of Animal Experimentation Committee (DEC) of the University of Groningen in accordance to the Dutch laws [reference number 6351B]. This work was financially supported by an Ubbo Emmius PhD grant from the University of Groningen, supplemented by the Province of Friesland and a start-up grant of the University of Groningen to TP. Stable isotope analyses were financed by an operating grant to KAH from Environment and Climate Change Canada.

References

- Alerstam T (1990) Bird migration. Cambridge University Press, Cambridge
- Bacetti N, Chelazzi L, Colombini I, Serra L (1998) Preliminary data on the diet of migrating ruffs *Philomachus pugnax* in northern Italy. *Int Wader Stud* 10:361–364
- Baker AJ, Gonzalez PM, Piersma T, Niles LJ, de Lima S, do Nascimento I, Atkinson PW, Collins P, Clark NA, Minton CDT, Peck MK, Aarts G (2004) Rapid population decline in red knots: fitness consequences of decreased refuelling rates and late arrival in Delaware Bay. *Proc R Soc B* 271:875–882. doi:10.1098/rspb.2003.2663
- Bearhop S, Waldron S, Votier SC, Furness RW (2002) Factors that influence assimilation rates and fractionation of nitrogen and carbon stable isotopes in avian blood and feathers. *Physiol Biochem Zool* 75:451–458. doi:10.1086/342800
- Bowen GJ, Wassenaar LI, Hobson KA (2005) Global application of stable hydrogen and oxygen isotopes to wildlife forensics. *Oecologia* 143:337–348. doi:10.1007/s00442-004-1813-y
- Castelijns H (1994) Black-tailed godwit *Limosa limosa islandica* and ruff *Philomachus pugnax* winter in increasing numbers in Zeeuws-Vlaanderen (SW-Netherlands). *Limosa* 67:113–11
- Catry T, Lourenço PM, Lopes RJ, Bocher P, Carneiro C, Alves JP, Delaporte P, Bearhop S, Piersma T, Granadeiro JP (2016) Use of stable isotope fingerprints to assign wintering origin and trace shorebird movements along the East-Atlantic flyway. *Basic Appl Ecol* 17:177–187. doi:10.1016/j.baec.2015.10.005
- Caut S, Angula E, Courchamp F (2009) Variation in discrimination factors ($\Delta^{15}\text{N}$ and $\Delta^{13}\text{C}$): the effect of diet isotopic values and application for diet reconstruction. *J Appl Ecol* 46:443–453. doi:10.1111/j.1365-2664.2009.01620.x
- Christianen MJA, Middelburg JJ, Holthuisen SJ, Jouta J, Compton TJ, van der Heide T, Piersma T, Sinninghe Damsté JS, van der Veer HW, Schouten S, Olff H (2017) Benthic primary producers are key to sustain the Wadden Sea food web: a stable carbon isotope analysis at landscape scale. *Ecology* 98:1498–1512
- Coulton DW, Clark RG, Hebert CE (2010) Determining natal origins of birds using stable isotopes ($\delta^{34}\text{S}$, $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$): model validation and spatial resolution for mid-continent Mallards. *Waterbirds* 33:10–21
- Cramp S, Simmons KEL (1983) The birds of the Western Palearctic, vol III. Oxford University Press, Oxford
- Devos K, T’Jolyn F, Brosens D, Desmet P (2012) Watervogels—Wintering waterbirds in Flanders, Belgium. v3.3. Research Institute for Nature and Forest (INBO). Dataset/Occurrence. doi:10.15468/lj0udq
- Dietz MW, Spaans B, Dekinga A, Klaassen M, Korthals H, van Leeuwen C, Piersma T (2010) Do red knots (*Calidris canutus islandica*) routinely skip Iceland during southward migration? *Condor* 112:48–55. doi:10.1525/cond.2010.090139
- Dietz MW, Piersma T, Dekinga A, Korthals H, Klaassen M (2013) Unusual patterns in ^{15}N blood values after a diet switch in red knot shorebirds. *Isotopes Environ Health Stud* 49:283–292. doi:10.1080/10256016.2013.776045
- Evans Ogden JL, Hobson KA, Lank DB (2004) Blood isotopic ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) turnover and diet-tissue fractionation factors in captive dunlin (*Calidris alpina pacifica*). *Auk* 121:170–177. doi:10.1642/0004-8038(2004)121
- Gill JA, Clark J, Clark N, Sutherland WJ (1995) Sex-differences in the migration, moult and wintering areas of British ringed ruff. *Ring Migr* 16:159–167. doi:10.1080/03078698.1995.9674107
- Gill REJR, Tibbitts TL, Douglas DC, Handel CM, Mulcahy DM, Gottschalck JC, Warnock N, McCaffery BJ, Battley PF, Piersma T (2009) Extreme endurance flights by landbirds crossing the Pacific Ocean: ecological corridor rather than barrier? *Proc R Soc Lond B* 276:447–457. doi:10.1098/rspb.2008.1142
- Gilroy JJ, Gill JA, Buchart SHM, Jones VR, Franco AMA (2016) Migratory diversity predicts population declines in birds. *Ecol Lett* 19:301–317. doi:10.1111/ele.12569
- Gutiérrez-Espósito C, Ramírez F, Afán I, Forero MG, Hobson KA (2015) Toward a deuterium feather isoscape for sub-Saharan Africa: progress, challenges and the path ahead. *PLoS ONE* 10:e0135938. doi:10.1371/journal.pone.0135938
- Hebert CE, Wassenaar LI (2001) Stable nitrogen isotopes in waterfowl reflect agricultural land use in western Canada. *Envir Sci Technol* 35:3482–3487
- Hebert CE, Wassenaar LI (2005) Feather stable isotopes in western north American waterfowl: spatial patterns, underlying factors, and management applications. *Wildl Soc Bull* 33:92–102
- Hobson KA (1999) Tracing origins and migration of wildlife using stable isotopes: a review. *Oecologia* 120:314–326. doi:10.1007/s004420050865
- Hobson KA (2005) Stable isotopes and the determination of avian migratory connectivity and seasonal interactions. *Auk* 122:1037–1048. doi:10.1642/0004-8038(2005)122
- Hobson KA, Clark RG (1992) Assessing avian diets using stable isotopes I: turnover of ^{13}C in tissues. *Condor* 94:181–188. doi:10.2307/1368807
- Hobson KA, Wassenaar LI (2008) Tracking animal migration with stable isotopes. Academic Press, London
- Hobson KA, van Wilgenburg SL, Wassenaar LI, Larson K (2012) Linking hydrogen ($\delta^2\text{H}$) isotopes in feather and precipitation: source of variance and consequences for assignment to isoscapes. *PLoS ONE* 7:e35137. doi:10.1371/journal.pone.0035137
- Hornman M, Hustings F, Koffijberg K, Klaassen O, Kleefstra R, van Winden E, Sovon Ganzen en Zwanenwerkgroep, Soldaat L (2013) Rapport Sovon-Watervogels in Nederland 2011/2012. Sovon Vogelonderzoek Nederland, Nijmegen

- Hortas F, Masero J (2012) Combatiente. In: Atlas de las aves en invierno en España 2007–2010. del Moral JC, Molina B, Ana Bermejo A, Palomino D (eds) Ministerio de Agricultura, Alimentación y Medio Ambiente-SEO/BirdLife, Madrid, pp 254–255
- IAEA/WMO (2001) Global network of isotopes in precipitation: the GNIP database. <http://www.iaea.org/water>
- Iwamura T, Possingham HP, Chadès I, Minton C, Murray NJ, Rogers DI, Treml EA, Fuller RA (2013) Migratory connectivity magnifies the consequences of habitat loss from sea-level rise for shorebird populations. *Proc R Soc B* 280:20130325. doi:10.1098/rspb.2013.0325
- Iwamura T, Fuller RA, Possingham HP (2014) Optimal management of a multispecies shorebird flyway under sea-level rise. *Conserv Biol* 28:1710–1720. doi:10.1111/cobi.12319
- Jukema J, Piersma T (2006) Permanent female mimics in a lekking shorebird. *Biol Lett* 2:161–164. doi:10.1098/rsbl.2005.0416ER
- Jukema J, Wymenga E, Piersma T (2001) Stopping over in SW Friesland: fattening and moulting in ruffs *Philomachus pugnax* during northward migration in the Netherlands. *Limosa* 74:17–26
- Karasov WH, Martinez del Rio C (2007) Physiological ecology: How animals process energy, nutrients and toxins. Princeton University Press, Princeton
- Karlionova N, Meissner W, Pinchuk P (2008) Differential development of breeding plumage in adult and second-year male ruffs *Philomachus pugnax*. *Ardea* 96:39–45. doi:10.5253/078.096.0105
- Kendall C (1998) Tracing nitrogen sources and cycling catchment. In: Kendall C, McDonnell JJ (eds) Isotope tracers in catchment hydrology. Elsevier, New York, pp 519–576
- Klaassen M, Piersma T, Korthals H, Dekinga A, Dietz M (2010) Single-point isotope measurements in blood cells and plasma to estimate the time since diet switches. *Funct Ecol* 24:796–804. doi:10.1111/j.1365-2435.2010.01689.x
- Koopman K (1986) Primary moult and weight changes of ruffs in the Netherlands in relation to migration. *Ardea* 74:69–77
- López-Calderón C, Hobson KA, Marzal A, Balbontín J, Reviriego M, Magallanes S, García-Longoria L, de Lope F, Møller AP (2017) Wintering areas predict age-related breeding phenology in a migratory passerine bird. *J Avian Biol* 48:631–639
- Márquez-Ferrando R, Figuerola J, Hooimeijer JCEW, Piersma T (2014) Recently created man-made habitats in Doñana provide alternative wintering space for the threatened continental European black-tailed godwit population. *Biol Cons* 171:127–135. doi:10.1016/j.biocon.2014.01.022
- Marra PP, Hobson KA, Holmes RT (1998) Linking winter and summer events in a migratory bird by stable-carbon isotope. *Science* 282:1884–1886. doi:10.1126/science.282.5395.1884
- Meissner W, Scebba S (2005) Intermediate stages of age characters create dilemmas in ageing female ruffs *Philomachus pugnax* in spring. *Wader Study Group Bull* 106:30–33
- Melter J, Sauvage A (1997) Measurements and moult of ruffs *Philomachus pugnax* wintering in West Africa. *Malimbus* 19:12–18
- Münster OAG (1998) Mass of ruffs *Philomachus pugnax* wintering in West Africa. *Int Wader Stud* 10:435–440
- Newton I (2008) The migration ecology of birds. Academic Press, Amsterdam
- Onrust J, Loonstra AHJ, Schmaltz LE, Verkuil YI, Hooimeijer JCEW, Piersma T (2017) Detection of earthworm prey by ruffs *Philomachus pugnax*. *Ibis*, in press. doi:10.1111/ibi.12467
- Oppel S, Pain DJ, Lindsell JA, Lachmann L, Diop I, Tegetmeyer C, Donald PF, Anderson G, Bowden GR, Tanneberger F, Flade M (2011) High variation reduces the value of feather stable isotope ratios in identifying new wintering areas for aquatic warblers *Acrocephalus paludicola* in West Africa. *J Avian Biol* 42:342–354. doi:10.1111/j.1600-048X.2011.05252.x
- Ouwehand J, Ahola MP, Aulsems ANMA, Bridge ES, Burgess M, Hahn S, Hewson CM, Klaassen RHG, Laaksonen T, Lampe HM, Velmala W, Both C (2015) Light-level geolocators reveal migratory connectivity in European populations of pied flycatchers *Ficedula hypoleuca*. *J Avian Biol* 47:69–83. doi:10.1111/jav.00721
- Pearson DJ (1981) The wintering and moult of ruffs *Philomachus pugnax* in the Kenyan Rift valley. *Ibis* 123:158–182. doi:10.1111/j.1474-919X.1981.tb00922.x
- Pernollet CA, Guelmami A, Green AJ, Curcú Masip A, Dies B, Bogliani G, Tesio F, Brogi A, Gauthier-Clerc M, Guillemain M (2015) A comparison of wintering duck numbers among European rice production areas with contrasting flooding regimes. *Biol Cons* 186:214–224. doi:10.1016/j.biocon.2015.03.019
- Peterson BJ, Fry B (1987) Stable isotopes in ecosystem studies. *Annu Rev Ecol Syst* 18:293–320. doi:10.1146/annurev.es.18.110187.001453
- Piersma T (2007) Using the power of comparison to explain habitat use and migration strategies of shorebirds worldwide. *J Ornithol* 148:S45–S59. doi:10.1007/s10336-007-0240-3
- Piersma T, Lok T, Chen Y, Hassell CJ, Yang HY, Boyle A, Slaymaker M, Chan YC, Melville DS, Zhang ZW, Ma Z (2016) Simultaneous declines in summer survival of three shorebird species signals a flyway at risk. *J Appl Ecol* 53:479–490. doi:10.1111/1365-2664.12582
- Post DM (2002) Using stable isotopes to estimate trophic position: models, methods, and assumptions. *Ecology* 83:703–718
- Prater AJ (1973) The wintering population of ruffs in Britain and Ireland. *Bird Study* 20:245–250. doi:10.1080/00063657309476388
- Prater AJ, Marchant JH, Vuorinen J (1977) Guide to the identification and ageing of Holarctic waders. British Trust for Ornithology, UK
- Qninba A, Dakki M, Benhoussa A, El-Agbani MA (2006) Rôle de la côte Atlantique marocaine dans l'hivernage des limicoles (Aves, Charadrii). *Ostrich* 78:489–493. doi:10.2989/OSTRICH.2007.78.2.59.173
- R Development Core Team (2016) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>
- Rakhimberdiev E, Verkuil YI, Saveliev AA, Väisänen RA, Karagicheva J, Soloviev MY, Tomkovich PS, Piersma T (2010) A global population redistribution in a migrant shorebird detected with continent-wide qualitative breeding survey data. *Divers Distrib* 17:144–151. doi:10.1111/j.1472-4642.2010.00715.x
- Runge CA, Watson JEM, Butchart SHM, Hanson JO, Possingham HP, Fuller RA (2015) Protected areas and global conservation of migratory birds. *Science* 350:1255–1257. doi:10.1126/science.aac9180
- Sánchez-Guzmán JM, Morán R, Masero JA, Corbacho C, Costillo E, Villegas A, Santiago-Quesada F (2007) Identifying new buffer areas for conserving waterbirds in the Mediterranean basin: the importance of the rice in Extremadura, Spain. *Biodivers Conserv* 16:3333–3344. doi:10.1007/s10531-006-9018-9
- Schmaltz LE, Juillet C, Tinbergen JM, Verkuil YI, Hooimeijer JCEW, Piersma T (2015) Apparent annual survival of staging ruffs during a period of population decline: insights from sex and site-use related differences. *Popul Ecol* 57:613–624. doi:10.1007/s10144-015-0511-4
- Schmaltz LE, Vega ML, Verkuil YI, Hooimeijer JCEW, Piersma T (2016) Use of agricultural fields by ruffs in southwest Friesland in 2003–2013. *Ardea* 104:23–32. doi:10.5253/arde.v104i1.a2
- Skagen SK, Bart J, Andres B, Brown S, Donaldson G, Harrington B, Johnston V, Jones SL, Morrison RIG (2003) Monitoring the

- shorebirds of North America: towards a unified approach. Wader Study Group Bull 100:102–104
- Taylor CM, Norris DR (2010) Population dynamics in migratory networks. *Theor Ecol* 3:65–73. doi:[10.1111/j.1600-048X.2012.05573.x](https://doi.org/10.1111/j.1600-048X.2012.05573.x)
- Thorup K, Korner-Nivergelt F, Cohen EB, Baillie SR (2014) Large-scale spatial analysis of ringing and re-encounter data to infer movement patterns: a review including methodological perspectives. *Methods Ecol Evol* 5:1137–1350. doi:[10.1111/2041-210X.12258](https://doi.org/10.1111/2041-210X.12258)
- Tøttrup AP, Klaassen RHG, Strandberg R, Thorup K, Kristensen MW, Jørgensen PS, Fox J, Afanasyev V, Rahbek C, Alerstam T (2012) The annual cycle of a trans-equatorial Eurasian-African passerine migrant: different spatio-temporal strategies for autumn and spring migration. *Proc R Soc B* 279:1008–1016. doi:[10.1098/rspb.2011.1323](https://doi.org/10.1098/rspb.2011.1323)
- Tréca B (1990) Régimes et préférences alimentaires d'anatidés et de scolopacidés dans le delta du Sénégal, études de leur capacités d'adaptation aux modifications du milieu. Exploitation des milieux cultivés. PhD Thesis
- Tréca B (1994) The diet of ruffs and Black-tailed Godwits in Senegal. *Ostrich* 65:256–263
- Triplet P, Diop I, Sylla SI, Schricke V (2014) Les oiseaux d'eau dans le delta du fleuve Sénégal (rive gauche). Bilan de 25 années de dénombrements hivernaux (mi-janvier 1989–2014). OMPO, ONCFS, DPN, SMBS, p 125
- van der Kamp J, Diallo M, Fofana B (2002a) Dynamique des populations d'oiseaux d'eau. In: Wymenga E, Kone B, van der Kamp J, Zwarts L (eds) Delta Intérieur du Niger: ecologie et gestion durable des ressources naturelles. A&W/Wetlands International/Rijkswaterstaat, Veenwouden, pp 87–140
- van der Kamp J, Zwarts L, Diallo M (2002b) Niveaux de crue, oiseaux d'eau et ressources alimentaires disponibles. In: Wymenga E, Kone B, van der Kamp J, Zwarts L (eds) Delta Intérieur du Niger: Ecologie et gestion durable des ressources naturelles. A&W/Wetlands International/Rijkswaterstaat, Veenwouden, pp 141–161
- van Gils JA, Ahmedou Salem MV (2015) Validating the incorporation of ^{13}C and ^{15}N in a shorebird that consumes an isotopically distinct chemosynthetic bivalve. *PLoS ONE* 10:e0140221. doi:[10.1371/journal.pone.0140221](https://doi.org/10.1371/journal.pone.0140221)
- van Rhijn JG (1991) The ruff: individuality in a gregarious wading bird. T. & A.D. Poyser, London
- Verkuil YI, de Goeij P (2003) Do reeves make different choices? Meadow selection by spring staging ruffs *Philomachus pugnax* in Southwest Friesland. *Limosa* 76:157–168
- Verkuil YI, Wymenga JJ, Hooijmeijer JCEW, Piersma T (2010) Spring migration of ruffs *Philomachus pugnax* in Fryslân: estimates of staging duration using resighting data. *Ardea* 98:21–33. doi:[10.5253/078.098.0104](https://doi.org/10.5253/078.098.0104)
- Verkuil YI, Karlionova N, Rakhimberdiev E, Jukema J, Wymenga JJ, Hooijmeijer JCEW, Pinchuk P, Wymenga E, Baker AJ, Piersma T (2012) Losing a staging area : eastward redistribution of Afro-Eurasian ruffs is associated with deteriorating fuelling conditions along the western flyway. *Biol Conserv* 149:51–59. doi:[10.1016/j.biocon.2012.01.059](https://doi.org/10.1016/j.biocon.2012.01.059)
- Wassenaar LI, Hobson KA (2003) Comparative equilibration and online technique for determination of non-exchangeable hydrogen of keratins for use in animal migration studies. *Isot Environ Health S* 39:211–217
- Webster MS, Marra PP, Haig SM, Bensch S, Holmes RT (2002) Links between worlds: unraveling migratory connectivity. *Trends Ecol Evol* 17:76–83. doi:[10.1016/S0169-5347\(01\)02380-1](https://doi.org/10.1016/S0169-5347(01)02380-1)
- Werner SJ, Hobson KA, Van Wilgenburg S, Fischer JW (2016) Multi-Isotopic ($\delta^2\text{H}$, $\delta^{13}\text{C}$, $\delta^{15}\text{N}$) tracing of moult origin for Red-winged Blackbirds associated with agro-ecosystems. *PLoS ONE* 11:e0165996. doi:[10.1371/journal.pone.0165996](https://doi.org/10.1371/journal.pone.0165996)
- Wilcove DS, Wikelski M (2008) Going, going gone: is animal migration disappearing? *PLoS Biol* 6:e188. doi:[10.1371/journal.pbio.0060188](https://doi.org/10.1371/journal.pbio.0060188)
- Wymenga E (1999) Migrating ruffs, *Philomachus pugnax*, through Europe, spring 1998. Wader Study Group Bull 88:43–48
- Wymenga E, Zwarts L (2010) Use of rice fields by birds in West Africa. *Waterbirds* 33:97–104. doi:[10.1675/063.033.s107](https://doi.org/10.1675/063.033.s107)
- Wymenga E, Kone B, van der Kamp J, Zwarts L (2002) Delta intérieur du fleuve Niger: Ecologie et gestion durable des ressources naturelles. A&W/Wetlands International/Rijkswaterstaat, Veenwouden
- Yerkes T, Hobson KA, Wassenaar LI, MacLeod R, Colucci JM (2008) Stable isotopes ($\delta^2\text{H}$, $\delta^{13}\text{C}$, $\delta^{15}\text{N}$) reveal associations among geographical location and condition of Alaskan northern pintails. *J Wildl Manage* 72:715–725. doi:[10.2193/2007-115](https://doi.org/10.2193/2007-115)
- Zwarts L, Bijlsma RG, van der Kamp J, Wymenga E (2009) Living on the edge: wetlands and birds in a Changing Sahel. KNNV Publishing, Zeist