University of Groningen

Endophytes

Staniek, Agata; Woerdenbag, Herman J.; Kayser, Oliver

Published in:
Journal of plant interactions

DOI:
[10.1080/17429140801886293](https://doi.org/10.1080/17429140801886293)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2008

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Staniek, A., Woerdenbag, H. J., & Kayser, O. (2008). Endophytes: Exploiting biodiversity for the improvement of natural product-based drug discovery. *Journal of plant interactions*, 3(2), 75-93. <https://doi.org/10.1080/17429140801886293>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Endophytes: exploiting biodiversity for the improvement of natural product-based drug discovery

Agata Staniek , Herman J. Woerdenbag & Oliver Kayser

To cite this article: Agata Staniek , Herman J. Woerdenbag & Oliver Kayser (2008) Endophytes: exploiting biodiversity for the improvement of natural product-based drug discovery, Journal of Plant Interactions, 3:2, 75-93, DOI: [10.1080/17429140801886293](https://doi.org/10.1080/17429140801886293)

To link to this article: <https://doi.org/10.1080/17429140801886293>

Copyright Taylor and Francis Group, LLC

Published online: 21 May 2008.

Submit your article to this journal [↗](#)

Article views: 1213

View related articles [↗](#)

Citing articles: 42 View citing articles [↗](#)

ORIGINAL ARTICLE

Endophytes: exploiting biodiversity for the improvement of natural product-based drug discovery

Agata Staniek, Herman J. Woerdenbag and Oliver Kayser*

Pharmaceutical Biology Department, University of Groningen, The Netherlands

(Received 12 November 2007; final version received 28 December 2007)

Endophytes, microorganisms that colonize internal tissues of all plant species, create a huge biodiversity with yet unknown novel natural products, presumed to push forward the frontiers of drug discovery. Next to the clinically acknowledged antineoplastic agent, paclitaxel, endophyte research has yielded potential drug lead compounds with antibacterial, antiviral, antioxidant, insulin mimetic, anti-neurodegenerative and immunosuppressant properties. Furthermore, while being implicated in livestock neurotoxicosis, some endophyte-produced alkaloids have been shown to display insecticidal activity. The endophyte-host relationship is postulated to be a 'balanced antagonism'. Moreover, the plausibility of horizontal gene transfer (HGT) hypothesis is taken into account. Knowledge of the genetic background of endophytic natural product biosynthesis is discussed on the basis of loline alkaloids, ergopeptines, lolitrems and maytansinoids. The current dynamic progress in genomics will contribute to a better understanding of endophytic microbes and to further exploiting them as a source of pharmaceutically relevant compounds.

Keywords: endophyte; natural product; drug discovery; plant-microbe interactions

Introduction

Endophyte (Gr. *endon*, within; *phyton*, plant) – the term was first coined by de Bary (de Bary 1866) and has become deeply embedded in the literature ever since. At present, endophytic organisms are defined as 'microbes that colonize living, internal tissues of plants without causing any immediate, overt negative effects' (Stone et al. 2000). First reports describing these microbes date back to the turn of the nineteenth and twentieth century (Freeman 1904). The most frequently encountered endophytes are representatives of the fungi; however, the existence of many endophytic bacteria has been documented as well.

Of the nearly 300,000 plant species inhabiting our planet, each individual one is host to several to hundreds of endophytes (Tan and Zou 2001), creating an enormous biodiversity: a myriad of undescribed species, a rich source of novel natural products therefrom and an unknown genetic background of all the interdependencies thus implied.

Several reviewers addressed the issue (e.g. Tan and Zou 2001; Strobel and Daisy 2003; Owen and Hundley 2004; Schulz and Boyle 2005; Gunatilaka 2006), shedding light onto selected aspects of the nature of endophytes. The aim of the hereby presented study is to render a comprehensive overview of the current knowledge on the subject. Considerable emphasis is put on the role and potential of metabolic engineering (Tyo et al. 2007) and combinatorial biosynthesis (Floss 2006), as well as on their impact on exploiting the astounding diversity of the endophytic world for its pharmaceutical potential.

Endophytes as biological factories of functional metabolites

Endophytic natural products as drugs and novel drug leads

It seems impossible to underestimate the impact of natural products on the drug discovery process. Even with lack of support from the major pharmaceutical companies, arguing that the natural product screening paradigm established in the late 1980s and early 1990s was no longer compatible with the HTS (high-throughput screening) approach, the significance of secondary metabolites as a source of novel drugs and drug-leads is still alive and well. Over 20 natural-product-derived drugs have been launched onto the worldwide market from 2001–2005 and around 140 have undergone various stages of clinical development in all major therapeutic areas (Butler 2005; Lam 2007).

The recent development and implementation of new technologies offers unique opportunities in the screening of natural products and will re-establish them as a major source for drug discovery. Improved natural product sourcing, aiming at narrowing the focus from all available sources to a single most prolific and reliable one, points to microorganisms as the ultimate, readily renewable, reproducible and inexhaustible source of novel structures bearing pharmaceutical potential. It is believed that up to 99% of microorganisms have yet to be discovered (Davis et al. 2005). This undefined richness of microbial world encompasses the plethora of endophytic entities occupying utterly millions of unique

*Corresponding author. Email: o.kayser@rug.nl

biological niches (higher plants) in various, many a time unusual, environments (Strobel 2006).

Functional metabolites of endophytic origin have already demonstrated a considerable potential to impact the pharmaceutical arena (Tan and Zou 2001; Strobel 2003; Strobel and Daisy 2003; Strobel et al. 2004; Gunatilaka 2006). A few examples are presented in this section, with the focus on their presumed therapeutic significance.

As the anti-infective branch is experiencing a shortage of lead compounds progressing into clinical trials, new antibacterial templates with novel mechanisms of action should have advantages over known antibiotics, especially in the fight against multi-drug resistant bacteria and emerging pathogens. Guanacastepene (Figure 1), a novel diterpenoid produced by a fungus isolated from the branch of *Daphnopsis americana* growing in Guanacaste, Costa Rica, might prove a representative of a potentially new class of antibacterial agents, showing activity against methicillin-resistant *Staphylococcus aureus* and vancomycin-resistant *Enterococcus faecium* (Singh et al. 2000).

Although apparently the potential for the discovery of endophytic compounds having antiviral activity is in its infancy, probably due to absence of appropriate antiviral screening systems in most compound discovery programmes, some promising fungal metabolites have been found. Initially, two novel tridepside human cytomegalovirus (hCMV) protease inhibitors, cytonic acids A and B were isolated from *Cytonaema* sp. inhabiting the internal tissues of *Quercus* sp. (Guo et al. 2000). Further studies of the microbial flora characteristic of oak trees resulted in the isolation of a potentially valuable fungal specimen from the leaves of *Quercus coccifera*. This endophyte proved to be a synthesizer of hinnuliquinone (Figure 2) – a potent inhibitor of the HIV-1 protease (Singh et al. 2004).

Two isobenzofuranones, pestacin and isopestacin (Figure 3), with structural similarity to the flavonoids – a well-established group of free-radical-scavengers, proved to exceed the anti-oxidant activity of trolox (a vitamin E derivative) by at least one order of

Figure 1. Guanacastepene.

Figure 2. Hinnuliquinone.

magnitude, as measured by the total oxyradical scavenging capacity (TOSC) assay. These new potent antioxidants were obtained from *Pestalotiopsis microspora*, an endophyte of *Terminalia morobensis* inhabiting the Sepik River drainage of Papua New Guinea (Strobel et al. 2002; Harper et al. 2003).

Pseudomassaria sp., a fungal endophyte recovered from leaves of an undetermined plant collected near Kinshasa, Democratic Republic of Congo, was shown to be a biofactory of nonpeptidal insulin mimetic, L-783,281 (Figure 4). Its discovery prompted quite a revolutionary notion in the therapy of diabetes, namely: an orally administered activator of the human insulin receptor (Zhang et al. 1999).

Moreover, this intriguing endophytic metabolite was reported capable of stimulating the *Trk* family of tyrosine kinase receptors, leading to the activation of multiple signalling cascades, culminating in neuro-regenerative effects, including neuronal survival and neurite outgrowth (Wilkie et al. 2001). Although the cytotoxicity of the compound seems to preclude its direct therapeutic application, it is a prototype for small molecule insulin and neurotrophin mimetics, and may lead to the development of pharmaceutically significant compounds for the treatment of diabetes and neurodegenerative disorders.

Following the track of search for new immunosuppressants from endophytes resulted in the isolation of subglutinol A (Figure 5) and B. These diterpene pyrones from *Fusarium subglutinans*, harbored by the perennial twining vine *Tripterygium wilfordii*, showed substantial immunosuppressive activity while causing none of the detrimental cytotoxic effects characteristic of cyclosporine A (Lee et al. 1995; US Patent 5648376, 1997).

Amongst the bioactive natural products of endophytic origin described to date, the ones notable for their antitumor activity seem to have drawn quite

Figure 3. Pestacin (a) and Isopestacin (b).

an unequivocal attention, with paclitaxel (Taxol) as the most striking example (Figure 6).

Since its discovery in the 1960s, through structure elucidation completed in 1971 and FDA approval in 1992, this highly functionalized diterpenoid natural product has evolved to become a blockbuster drug with commercial sales of well over \$3 billion in 2004 (Croteau 2005). Since it was primarily obtained from the inner bark of *Taxus brevifolia*, a relatively rare and slow-growing tree, the supply of this potent antineoplastic agent was soon to become the issue of scarcity, as even early estimations indicated that the demand for paclitaxel might exceed 300 kg, which would amount to 750,000 yew trees per year (Stierle et al. 1995). The supply crisis, as well as the ecological implications resulting in the plant-endangered status, prompted the search for paclitaxel-producing microorganisms among the endophytic fungi of *Taxus*. While the 1990s brought quite an abundance of reports on paclitaxel-producing endophytes (e.g. Stierle et al. 1993; Strobel et al. 1996, 1997; Li et al. 1998a, b; Noh et al. 1999; Zhou and Ping 2001), no conclusive follow-up data concerning fungal metabolite profile or genetic background of the biosynthetic pathway leading to paclitaxel in fungi is available at present. While the pursuit of the idea of a microbial paclitaxel source, providing for an inexhaustible supply of this antineoplastic blockbuster and novel taxanes, is being hampered by obtaining disappointingly low yields, the active ingredient is still mainly derived via chemical semisynthesis from the advanced taxoid, 10-deacetylbaccatin III, readily available from

the needles of the European yew tree, *Taxus baccata*, being a renewable source and, to a lesser extent, by means of plant cell culture methods (Leistner 2005; Frense 2007). Nonetheless, the exciting progress that has been made in the elucidation of biosynthetic route leading to paclitaxel *in planta* due to fundamental works of Croteau and his co-workers (Walker and Croteau 2001; Jennewein et al. 2004a, b; Croteau 2005; DeJong et al. 2006; Nims et al. 2006), as well as the recent advances in microbial genomics (Zazopoulos et al. 2003; Stephanopoulos et al. 2004; Keller et al. 2005; van Lanen and Shen 2006) and combinatorial biosynthesis (Floss 2006; Nguyen 2006; Klein-Marcuschamer et al. 2007), might still revive and boost the interest in endophytic paclitaxel synthesizers.

Agricultural aspects: endophytic insecticides vs. tremorgenic mycotoxins

As the world becomes wary of ecological damage done by synthetic insecticides, endophytic research

Figure 4. L-783,281.

Figure 5. Subglutinol A.

Figure 6. Paclitaxel.

continues for the discovery of powerful, selective and safer alternatives (Strobel and Daisy 2003).

From 1930–1970 millions of hectares in the temperate transition zone of the south central and southeastern USA were converted from revegetation shrubs and weeds to pastures of tall fescue (*Festuca arundinacea*). This grass was superior forage to native flora, but it was soon noticed that the expected livestock productiveness was not being met. An associated complex of symptoms episodically suffered by grazing livestock was termed ‘tall fescue toxicosis’ and consequently led to considerable economic losses to the US beef industry (Hoveland 1993; Thompson and Stuedeman 1993). Simultaneously, the livestock grazing on perennial ryegrass (*Lolium perenne*) pastures in New Zealand was afflicted by ‘ryegrass staggers’ (Gallagher et al. 1984; Bush et al. 1997). These problems spurred intense research that led to the isolation and identification of two respective fungal endophytes: *Neotyphodium coenophialum* and *Neotyphodium lolii*, both phylogenetically related to the ergot fungus *Claviceps purpurea* (Kuldau et al. 1997). Thus, the said endophytes were implicated in the associated livestock toxicoses, but both also greatly enhanced fitness of their plant hosts under biotic and abiotic stresses. Further investigation of the genus *Neotyphodium* (sexual state: *Epichloë*) proved its representatives capable of synthesizing four groups of alkaloids. Of these, the ergot alkaloids and tremorgenic lolitrems cause neurotoxic effects on grazing or granivorous vertebrates (Steyn and Vleggaar 1985; Bacon et al. 1986). Peramine is an insect feeding deterrent and protects perennial ryegrass from the Argentine stem weevil, *Listronotus bonariensis*, a highly destructive insect pest in New Zealand (Rowan and Latch 1994; Tanaka et al. 2005). Lolines are potent insecticidal and anti-aphid agents, not being implicated in any mammalian toxicoses (Jackson et al. 1996; Wilkinson et al. 2000). Obviously, the last of the aforementioned groups of alkaloids should provide for the most promising candidates to aid the agricultural arena.

As the forage grasses associated endophytes seem to constitute an extensively investigated and well documented microbial group (see further sections), they might prove to become a platform for the endophyte research to advance to the next level of modern microbial genomics and systems biology (Stephanopoulos et al. 2004; van Lanen and Shen 2006).

Endophyte vs. host: the relationship

A conflict of interests?

Over the years, despite the controversy it seemed to have aroused in the early 1990s (Wennström 1994), the term ‘endophyte’ has evolved from its original suggested definition, which described merely the location, to depict the nature of the association between the microbe and the host-plant (Wilson 1995). To truly define the interaction, however, seems to be quite a task. In fact, it can vary in a seamless manner from mutualism to parasitism, based on a fine-tuned balance between the demands of the invader and the plant response (Kogel et al. 2006).

A very elegant hypothesis has been proposed and extensively documented by Schulz and Boyle (2005), postulating the relationship to be a ‘balanced antagonism’ (Figure 7).

The said notion can be deciphered as equilibrium, under environmental, physiological and genetic control, that results in fitness benefits for both partners. On one hand, the theory depicts fungal endophytes as ‘masters of phenotypic plasticity’, able to infect as endosymbionts, to colonize cryptically, and finally to sporulate as pathogens or saprophytes. This creative variability implies evolutionary potential. On the other hand, it does not exclude the possibility of secondary metabolites being a contribution of the endophytic partner to a mutualistic relationship. A very interesting example of the protective power of an endophyte involves the aquatic plant *Rhynchospora penicillata* collected from a river system in Southwest Venezuela where the harsh aquatic environment subjected the plant to constant beating by virtue of rushing waters, debris, and tumbling rocks and pebbles. Such circumstances should provide ample opportunity for pathogenic oomycetes to invade the plant. Still, the plant population remained quite healthy. Upon extraction and investigation of the endophytic flora of *R. penicillata*, a potent anti-fungal bacterium, *Serratia marcescens*, was identified, which in turn was shown to produce oocydin A (Figure 8), a novel anti-oomycetous compound that obviously provided the plant with the requisite protection from the water molds (Strobel et al. 1999).

Recently, it has been hypothesized that endophytes may protect their host plants by scavenging the damaging reactive oxygen species (ROS) generated by the plant defence mechanisms in response to

Figure 7. Balanced antagonism hypothesis (Schulz and Boyle 2005).

environmental stress (Rodriguez and Redman 2005; Tanaka et al. 2006). Thus, a strong activation of ROS due to biotic and abiotic stresses on part of the host followed by an equally rapid free-radical-scavenging response of the microbe might potentially be a prime mechanism in maintaining the delicate balance between the two.

To address the question asked in the title of this section: there is always a 'conflict of interests' at all stages of relationships between endophytic and plant partners (Smith and Read 1997). However, the development of tools for non-invasive observation of subcellular activities during the establishment of mutualistic interactions will provide a deeper understanding of the mechanism that balances virulence against defence; hostility against hospitality (Kogel et al. 2006).

A relationship within a relationship

While fungal viruses have been shown capable of modulating plant-fungal symbioses by regulating the hypovirulence of pathogenic fungi (e.g. Ahn and Lee 2001; Dawe and Nuss 2001), the effect of mycoviruses on mutualistic fungal endophytes has not been described. But one report of a mycovirus from the well-known endophytic microbe, *Epichloë festucae*, was recorded (Zabalgoatzea et al. 1998). Still, no decisive phenotypic traits have been associated with this virus since.

Figure 8. Oocidin A.

However, a very recent study reports a complex tripartite symbiosis involving a virus, a fungal endophyte and a plant (Márquez et al. 2007). Thus far it was believed that the mutualistic relationship between a tropical panic grass from geothermal soils, *Dichanthelium lanuginosum*, and the fungus *Curvularia protuberata*, allowed both organisms to grow at high soil temperatures in Yellowstone National Park (Redman et al. 2002). The in-depth investigation of this beneficial interaction showed, however, that the ability of the fungus to confer heat tolerance to its host plant is, in fact, related to a presence of yet another kind. To reflect its host of origin and its unique phenotype the responsible mycovirus was named *Curvularia* thermal tolerance virus (CThTV).

A physical interaction merely? The plausibility of horizontal gene transfer (HGT) hypothesis

While enthusiasts call it 'the essence of the phylogenetic process and the driving force in a new paradigm for evolution' (Doolittle 1999), sceptics describe it as no more than one of many phylogenetic anomalies (Kurland et al. 2003). Though highly controversial, the hypothesis of horizontal gene transfer seems quite seductive.

Hereby, two intriguing examples from the endophytic world are discussed. Both instances deal with the occurrence of identical natural products in unrelated taxa, namely: the host and the invader.

The presence of maytansinoids, potent cytotoxic agents, was first noticed in the Ethiopian shrub, *Maytenus serrata* (Kupchan et al. 1972). Further investigation recorded their occurrence not only in higher plants (Wani et al. 1973; Ahmed et al. 1981; Powel et al. 1982), but also in mosses (Sakai et al. 1988; Suwanborirux et al. 1990) and, remarkably, in gram-positive *Actinomycetes* (Higashide et al. 1977; Asai et al. 1978). One could assume that the biosynthesis of these unique natural products has

been repeatedly invented during evolution. However, the fact that approximately 48 genes are involved in the bacterial synthesis of maytansinoids (Yu et al. 2002) makes it highly unlikely. Similarly, the aforementioned ubiquity of paclitaxel occurrence in yews as well as in taxonomically distant fungal microbes raises questions. Therefore, it seems possible that in the course of evolution a lateral (horizontal) gene transfer took place between different, taxonomically unrelated species, thus explaining the distant distribution of the antineoplastic secondary metabolites mentioned above between pro- and eukaryotes.

Nevertheless, before invoking HGT, alternative and often equally plausible, explanations ought to be thoroughly considered. In case of maytansinoids, all evidence seems to point to them being ultimately produced by plant associated microorganisms. Maytansine, the unique parent compound (Figure 9), was found neither in cell suspension cultures from *Maytenus buehnanii* (Kutney et al. 1981) nor in callus cultures raised from *Maytenus wallichiana* (Dymowski and Furmanowa 1990) and *Putterlickia verrucosa* (Pullen et al. 2003).

This is in line with the result of an in-depth search for the unique gene involved in maytansinoid biosynthesis, encoding for 3-amino-5-hydroxybenzoic acid (AHBA) synthase, in *Putterlickia verrucosa* cell cultures. An extensive PCR based homology screen gave negative results only (Pullen et al. 2003). These observations point to the conclusion that plants do not produce maytansinoids *ab initio*. However, an active role of the plant in an overall biosynthesis cannot be excluded, as it seems likely that the host converts a bacterially synthesized precursor into the final, biologically active compound. Secondly, it is possible that maytansine is only produced as a consequence of a pathogen attack on the plant. The plants may contain a biologically inactive bacterially produced precursor, which is only converted into the potent final product in response to a signal resulting from the attack. Alternatively, and more plausibly, the bacterial production of the maytansinoid pre-

cursor could be triggered by a plant signal in response to the pathogen aggression (Cassady et al. 2004).

On the contrary, the bio-formation of paclitaxel seems to be a genuine feature of the yew host, as ample evidence supporting the production of the diterpenoid by sterile cell suspension cultures of *Taxus* has been provided (e.g. Ketchum and Gibson 1996; Ketchum and Croteau 1998; Yukimune et al. 2000; Wu and Lin 2003; Naill and Roberts 2005; Khosroushahi et al. 2006; Vongpaseuth and Roberts 2007). This conclusion is further supported by the aforementioned work of Croteau and his associates who succeeded in the isolation of paclitaxel biosynthetic genes of plant origin. Interestingly, the taxadiene synthase gene has a long N-terminal targeting sequence for localization to and processing in the plastids, indicating that this gene is plant-derived rather than a fungal product (Koepp et al. 1995; Walker and Croteau 2001). Accordingly, an extensive PCR-based screen for taxadiene synthase gene in *Taxomyces andreanae*, the very first presumed endophytic taxane-producer (Stierle et al. 1993), failed to provide for any positive results (Staniek, unpublished data).

To sum up, the evidence for lateral gene transfer in eukaryotes remains largely anecdotal (Rosewich and Kistler 2000). However tempting and attractive, the HGT hypothesis has to give way to a more plausible alternative postulating the endophyte-host co-evolution.

Investigating the genetic background of endophytic biosynthetic pathways: identification, molecular cloning, genetic analysis and manipulation of endophytic gene clusters

Clusters of functionally related genes are a general feature of prokaryotic gene organization, but were believed to be much less prevalent in eukaryotes. However, the advent of recombinant DNA methodologies in the 1980s enabled dramatic progress in the genetics and biochemistry of fungal secondary metabolism. This rapid progress was facilitated by what is now considered a hallmark characteristic of secondary metabolic biosynthetic pathways – the grouping of pathway genes in a contiguous cluster (Keller and Hohn 1997; Keller et al. 2005).

The balansiaceous endophytes and their natural products

The balansiaceous endophytes form a unique group of closely related fungi with ecological requirements and adaptations distinct from those of other endophytes (Petrini 1996). They belong to the clavicipitaceous genera *Epichloë* and *Balansia*, and their anamorphs *Neotyphodium* and *Ephelis* (Schardl et al. 2004).

As previously stated, due to their ecological and economic impact, theirs seems to be the best studied group of endophytes. The following section focuses

Figure 9. Maytansine

on the recent advances in dissecting the molecular mechanisms driving the biosynthesis of their secondary metabolites.

Loline alkaloids (LA)

Unique not only in structure, comprising a saturated 1-aminopyrrolizidine-ring system, with a highly strained ether bridge between C-2 and C-7 (Figure 10) (Petroski et al. 1989), but also due to their potent, broad-spectrum insecticidal activities (Jackson et al. 1996; Wilkinson et al. 2000), loline alkaloids have remained a biosynthetic enigma till very recently (Blankenship et al. 2005).

Establishing that the production of lolines in axenic cultures of *Neotyphodium uncinatum* can be regulated by culture conditions (Blankenship et al. 2001), which in turn suggests differential expression of genes involved in LA biosynthesis, prompted a notion that the isolation of the genes up-regulated during loline production could be a first step in identifying possible enzymes catalyzing their biosynthesis. The recent development of novel highly effective methods for identifying differentially expressed transcripts, enabled the application of suppression subtractive hybridization technique (Diatchenko et al. 1996, 1999) and provided for a successful identification of two genes, *lolA* and *lolC*, associated with endophytic loline synthesis (Spiering et al. 2002).

These encouraging results spurred intense research on the molecular genetics of loline-alkaloid production, originating from a preliminary hypothesis of *lolA* and *lolC* being clustered (Spiering et al. 2005; US Patent 7183098, 2007). The question was tackled with an impressive arsenal of up-to-date techniques and ultimately resulted in the identification of double homologs of nine genes similarly arranged into two clusters, *LOL-1* and *LOL-2*. Long-range (LA: long and accurate) PCR and parallel screening of a partial genomic library of *N. uncinatum* to ascertain the existence of two highly similar, yet distinct gene clusters. Extensive genome walking into unknown regions yielding a total of 25 kb sequenced from *LOL-1* and 16 kb from *LOL-2*. Gene-prediction searches with the FGENESH pro-

gramme (available from: <http://www.softberry.com/berry.phtml>) aided by RACE (rapid amplification of cDNA ends) and confirmed by reverse transcription-PCR based expression verification for identifying and mapping ORFs (open reading frames). Large-scale utilization of Basic Local Alignment Search Tool (NCBI) to ascribe the putative functions of the enzymatic entities encoded by the nine *lol* genes. And, last but definitely not least, the very first attempt of gene silencing by RNAi (RNA interference) performed on a mutualistic fungus, providing direct and indisputable evidence for involvement of *lolC* in loline biosynthesis.

All in all, an unprecedented endeavor leading to a final conclusion that the products of the *lol* genes thus identified may be sufficient for the biosynthesis of the entire loline-alkaloid three-ring structure in *Neotyphodium uncinatum*, from the primary precursor amino acids, L-proline and L-homoserine (Figure 11) (Blankenship et al. 2005; Spiering et al. 2005).

As spectacular as it may seem, the aforementioned accomplishment did not put a closure to solving the riddle of endophytic lolines. On the contrary, it contributed to further identification of single *LOL* gene clusters in *Epichloë festucae*, *Neotyphodium* sp. PauTG-1 and *Neotyphodium coenophialum* (Kutil et al. 2007).

To analyze the genetic architecture and to predict the evolutionary history of *LOL*, the five characterized clusters were compared by means of the coupled powers of phylogenetic approaches and other forms of sequence analysis. All *lol* genes anchored to the map for each species turned out to occur in strictly conserved order and orientation (Figure 12).

Furthermore, PhyloCon-based (Wang and Stormo 2003) comparison of the putative *lol* gene promoter regions yielded the identification of four motifs conserved across the genes in all five clusters, with each motif having significant similarity to known fungal transcription factor binding sites in the TRANSFAC database (Wingender et al. 1996). Conservation of these motifs further supported the hypothesis of the *lol* genes being co-regulated (Spiering et al. 2002, 2005).

Interestingly, the history of asexual *Neotyphodium* spp. includes multiple interspecific hybridization events (Tsai et al. 1994; Schardl 2001; Clay and Schardl 2002; Moon et al. 2004). Thus, comparing the clusters from three seed-transmitted *Neotyphodium* anamorphs and their sexual counterpart, *E. festucae*, allowed determining which *Epichloë* ancestors were the most likely contributors of *LOL* in these asexual loline synthesizers. Perhaps the most striking outcome of this comparative analysis was that three *Neotyphodium* species with a history of *Epichloë typhina* as an ancestor possess nearly identical copies of *LOL* clusters (*N. uncinatum* *LOL-2*, *N. coenophialum*, *N. sp.* PauTG-1). This is most remarkable in light of the fact that there are no known isolates of

Figure 10. Loline.

Figure 11. The loline alkaloid biosynthetic pathway (Blankenship et al. 2005).

the extant species *E. typhina* which express loline alkaloids (Leuchtmann et al. 2000). Thus, these data support a model of evolution in which the polymorphism in loline alkaloid production phenotypes among endophytic species is likely due to the loss of the trait over time (Kutil et al. 2007).

Ergopeptines

Ergot alkaloids are produced by ascomycetous fungi from discontinuous taxonomic groupings, including

plant-associated fungal genera from the family *Clavicipitaceae* and some members of the order *Eurotiales*, including the human pathogen *Aspergillus fumigatus* (Panaccione 2005). Due to their long biotechnological tradition, with manifold applications in therapy (Tudzynski et al. 2001), the physiology and biochemistry of ergopeptine formation have been studied in minute detail (Floss 1976; Socic and Gaberc-Porekar 1992; Gröger and Floss 1998). Nevertheless, an uncomfortable premonition of scar-

Figure 12. Comparative maps of *LOL* gene cluster(s) from four endophyte species; the shaded boxes in the *E. festucae* map indicate regions with homology to *pol* proteins; disconnected contig lines indicate genes that are sequenced, but not anchored to other *lol* genes; the contig of sequence known for this region in *E. festucae* continues beyond the boundaries shown; arrows along the bottom of the diagram indicate locations of putative promoter regions analyzed in the motif analysis (Kutil et al. 2007).

city in knowledge about the genetics of the biosynthetic route leading to these valuable alkaloids seemed to have arisen.

Yet again, the dynamic progress in the field of molecular genetics proved invaluable. The first step towards revealing the molecular background of ergopeptine biosynthesis was taken: the gene coding for dimethylallyl tryptophan synthase (*DMATS*) – the committed enzyme of the synthetic process in question (Figure 13a) – was cloned via reverse genetics (Tsai et al. 1995), identified by a differential cDNA screening approach (Arntz and Tudzynski 1997) and characterized initially for *Claviceps fusiformis*, subsequently for *Claviceps purpurea* (Tudzynski et al. 1999), and finally for the phylogenetically distant *Aspergillus fumigatus* (Coyle and Panaccione 2005; Unsöld and Li 2005; Li and Unsöld 2006).

Fuelled by the conviction of secondary metabolite pathway genes clustering in fungal genomes (Keller and Hohn 1997), large-scale studies have been taken up, resulting in the identification of gene clusters for ergot alkaloid biosynthesis in *C. purpurea* (Tudzynski et al. 1999; Correia et al. 2003; Haarmann et al. 2005, 2006) and *A. fumigatus* (Coyle and Panaccione 2005; Unsöld and Li 2005; Li and Unsöld 2006); they contain 13 and 14 genes, respectively. It has been recently hypothesized that the fundamental ability to synthesize some type of ergot alkaloid was present in the most recent common ancestor of these *Ascomycetes*, accounting for the current phylogenetically discontinuous distribution of these secondary metabolites (Panaccione 2005). Thus, genes shared between the two clusters are presumed to be responsible for the early steps common to the two organisms (Coyle and Panaccione 2005).

The impressive achievements described hitherto, have been successfully conveyed into the realm of clavicipitaceous endophytic ergopeptine producers. *DMATS*, as a gene encoding for the enzymatic entity indispensable for ergot alkaloid production, was characterized for *Neotyphodium lolii* harbored by

perennial ryegrass. Once the putative endophyte homolog was identified by degenerate PCR, its function was tested by gene knockout. Subsequent complementation experiments resulted in the ultimate confirmation of the postulated role of *DMATS* in the biosynthetic process in question (Wang et al. 2004). Analogous experimental procedure allowed for the characterization of two consecutive genes, displaying the capacity to encode nonribosomal peptide synthetases (NRPS), namely: *lpsA* (Pannaccione et al. 2001, 2003) and *lpsB* (Tanaka et al. 2005; Fleetwood et al. 2007), playing a vital role in the ergopeptine formation (Figure 13b) (Riederer et al. 1996; Walzel et al. 1997; Correia et al. 2003). The ultimate objective: to determine whether ergot alkaloid biosynthetic genes were clustered in *Neotyphodium lolii* was pursued by using chromosome walking and Southern blot analysis and successfully attained (Fleetwood et al. 2007).

Interestingly, while gene sequence is relatively highly conserved between each of the three thus far identified ergot alkaloid gene clusters, there are several differences in gene order and the *N. lolii* cluster (*EAS*) seems to be more complex in structure and organization, as compared to its pathogenic counterparts (Figure 14). What is more, the BLASTN (NCBI) and MEME (Multiple EM for Motif Elicitation) (Bailey and Elkan 1994) analyses of the *EAS* cluster revealed that its genes are closely associated with transposon relics, including retrotransposons and autonomous as well as non-autonomous DNA transposons.

Finally, it has been shown that ergot alkaloids are specifically produced during biotrophic growth of *Epichloë* endophytes (Tanaka et al. 2005). Attendant with this, the *eas* genes proved all to be expressed *in planta*, while giving no evidence for expression under axenic culture conditions. These results suggest that specific plant environment may be required for the induction of ergopeptine biosynthetic genes in the fungal endophyte (Fleetwood et al. 2007).

Notably, ergot alkaloids have also been considered to constitute a chemotaxonomic signature of

Figure 13. The ergot alkaloid biosynthetic pathway; biosynthesis of lysergic acid (a) and further ergopeptine formation, as exemplified by ergotamine (b) (Tudzynski et al. 1999, 2001).

Convolvulaceae plants. However, recent studies on *Ipomoea asarifolia*, a representative of this dicotyledonous plant family, indicated that the accumulation of the natural products in question depends, in fact, on the presence of a plant-associated clavicipi-

taceous fungus. Thus, it has been postulated that ergopeptines are not typical of the *Ascomycota*, the *Poaceae* or the *Convolvulaceae* alone, but, in contrast, they are very likely to prove a trait of a clavicipitaceous fungal taxon capable of colonizing certain

Figure 14. Comparative *eas* gene order among *C. purpurea* (Cp), *N. lolii* (Nl) and *A. fumigatus* (Af); black arrows, genes proposed to be required for ergot alkaloid synthesis, but not yet identified for *N. lolii*; light grey arrows, genes found in *N. lolii* and *C. purpurea*, but not in *A. fumigatus*; dark grey arrows, genes found only in *A. fumigatus*; *N. lolii* *dmaW*, *eloA* and *lpsA* are not shown, as their locations relative to *EAS* cluster are not known (Fleetwood et al. 2007).

higher plants in mutualistic symbiosis (Kucht et al. 2004; Steiner et al. 2006).

Lolitrems

Indole-diterpenes are a large, structurally diverse group of natural products commonly found in filamentous fungi of the genera *Penicillium*, *Aspergillus* and *Claviceps* (Mantle 1987; Parker and Scott 2004). Many of these compounds are potent mammalian tremorgenic mycotoxins (Steyn and Vleggaar 1985) and fall into four major structural classes, namely the penitrems, janthitrems, paspalitrems and lolitrems (Figure 15) (Mantle and Weedon 1994). They all share a common structural core comprised of a cyclic diterpene skeleton derived from geranylgeranyl diphosphate (GGPP) and an indole moiety derived from tryptophan, while different patterns of prenylation, hydroxylation, epoxydation and acetylation, and differences in ring stereochemistry around the basic indole-diterpene ring structure, are to be accountable for their structural diversity (Parker and Scott 2004). The best known of these fungal tremorgens are the lolitrems produced by *Epichloë* endophytes in association with temperate forage grasses (Gallagher et al. 1984).

The history of unravelling the molecular genetics underlying endophytic lolitrem production seems to encompass some of the elements of both the aforementioned biosynthetic puzzles of microbial alkaloids.

Initially, employing *Penicillium paxilli* as a model experimental system to dissect the biochemistry of indole-diterpene biosynthesis (Young et al. 2001) (an obvious parallel can be drawn with the ergots and their model synthesizer: *Claviceps purpurea*) allowed for the isolation of three orthologous genes from the grass endophytes, *Neotyphodium lolii* and *Epichloë festucae*, and subsequent confirmation of their contiguous clustering (Young et al. 2005). While further attempts to clone additional linked *ltm* genes in the endophytes by such approaches as inverse PCR were thwarted by the presence of large blocks of highly repetitive retrotransposon sequences flanking both sides of the cluster, an alternative strategy combining suppression subtrac-

tive hybridization (a technique applied successfully in case of *LOL* characterization) with chromosome walking was taken up. As a result, two additional *LTM* gene clusters were identified and consequently proved to be linked with *LTM-1* (Young et al. 2006).

Ultimately, it has been shown that the genes for lolitrem biosynthesis in the endophytic anamorph *Neotyphodium lolii* form a complex genetic locus of at least three *LTM* gene clusters (Figure 16). What is more, the structure and organization of the said endophytic locus, as compared to the orthologous clusters from *Penicillium paxilli* (Young et al. 2001) and *Aspergillus flavus* (Zhang et al. 2004), has yet again proved to be of greater complexity (accordingly: *EAS* clusters of pathogenic vs. endophytic origin).

As sub-telomers are proposed to be 'a workshop of evolutionary genomic experimentation' (Wong and Wolfe 2005), it has been suggested that the elaborate nature of the cluster in question may be accounted for by its location in such a rapidly evolving region of the endophytic genome (Gardner et al. 2002; Kellis et al. 2003; Machida et al. 2005; Nierman et al. 2005), and consequently driven by recombination and mutational processes associated with Type I transposon elements (Kempken and Kück 1998) combined with the phenomenon of repeat induced point mutation (RIP) (Selker et al. 1987).

Thus, consecutive similarities between *LTM* and *EAS* loci seem to be brought to light. Firstly, the association of the genes composing both these endophytic clusters with the transposon relics appears to be of considerable importance (Young et al. 2006; Fleetwood et al. 2007). The abundance of these relic sequences adds to the evolutionary potential of both clusters. It might also suggest that the regulation at the level of chromatin may be of consequence for *ltm* and *eas* genes, since chromatin remodelling as a method of coordinate gene regulation has been proposed as a possible factor causing selection pressure for secondary metabolite genes to be clustered (Martienssen and Colot 2001; Volpe et al. 2002; Bok and Keller 2004; Keller et al. 2005; Lee et al. 2005). Secondly, all ten genes identified at the *LTM* locus, as well as each gene

Figure 16. A physical map of the *N. lolii* LTM locus; the boundaries of the three *ltm* gene clusters are identified by boxes numbered 1–3; the genes, abbreviated to a single letter, are shown as arrows indicating the direction they are transcribed, with the exon structure shown as black blocks underneath each gene; the blocks between *ltm* clusters 2 and 3 show the positions of two imperfect direct repeats; the retrotransposon relics, Tahi and Rua, are identified by blocks above the sequence; the AT content of each region is indicated as a percentage at the bottom of the figure; the distance across the sequence, shown immediately below the genes, is represented in kb; the sequence containing *ltm* clusters 2 and 3 is separated from *ltm* cluster 1 by ~35 kb (Young et al. 2006).

thetic processes seem to emphasize the ultimate uniqueness of lolines.

Maytansinoids from bacterial endophytes

As a matter of principle, hereby the elucidation of molecular background underlying the clustering of prokaryotic genes in the realm of endophytic secondary metabolism is shortly depicted. Thus, encompassing the diversity of microbial endophytic world, as well as acknowledging the early foundations set to boost the gene clustering investigation *per se* will be attempted.

The remarkable antitumor potency of maytansinoids, as well as the mystery surrounding their biosynthetic and evolutionary origins was reflected upon previously. It is noteworthy, however, that these 19-membered macrocyclic lactams are closely related to ansamycin antibiotics of microbial origin, such as rifamycin B and geldanamycin (Rinehart and Shield 1976). In fact, the aforementioned similarity stimulated a search for maytansinoid-producing microorganisms, ultimately leading to the isolation of ansamitocins (Figure 17) from the *Actinomycetes*, *Actinosynnema pretiosum* ssp. *pretiosum* and a mutant strain *Actinosynnema pretiosum* ssp. *auranticum* (Higashide et al. 1977; Asai et al. 1978).

The drive at the elucidation of microbial maytansinoids biosynthesis at the biochemical and genetic level was based upon the coupled forces of heterologous hybridization and PCR. The primary aim was to identify the gene encoding for the key enzyme in the assembly of 3-amino-5-hydroxybenzoic acid (AHBA) as the starter unit. Cloning by reverse genetics of the *AHBA* synthase from *Amycolatopsis mediterranei* (Kim et al. 1998) provided the necessary probe and set the pace for identification, isolation and sequencing of the consecutive entities embedded in the biosynthetic gene cluster in question. A total of 250 kb of contiguous DNA was cloned and mapped. The appropriate gene inactivations were carried out to address the essential role of the AHBA gene homologues and to define the boundary of the

biosynthetic locus (Yu et al. 2002). Notably, the ansamitocin synthesis genes (*asm*) turned out to be contained not in one, but two clusters in the *Actinosynnema pretiosum* genome (Figure 18). This separation of genes encoding the biosynthesis of a given antibiotic into two essential clusters is unprecedented in *Prokaryota*. Thus, it is proposed to be merely an inconsequential accident resulting from an evolutionary gene rearrangement which inserted 30 kb of non-essential DNA into the otherwise contiguous cluster (Yu et al. 2002).

Thus equipped with the essential genetic information and modern tools to manipulate the biosynthetic machinery, as well as urged by the extraordinary therapeutic potential, the research on microbial maytansinoids has recently entered a novel combinatorial stage. Two principal approaches are being explored. On one hand, the expression of all the *asm* genes, assembled into several cassettes in a heterologous host, such as *Streptomyces coelicolor* or *S. lividans* is being considered. This approach seems most promising in terms of elucidating the detailed biosynthetic process, but it has some major practical drawbacks, as the product titres are likely to be very low. On the other hand, it has been proposed to engineer mutations in the *asm* in the parent

Figure 17. Ansamitocin P-3.

Figure 18. The ansamitocin biosynthetic gene cluster from *A. pretiosum*; asmABCD and asm19, polyketide synthase; asm22-24, asm43-45 and asm47, AHBA synthesis; asm13-17, methoxymalonate synthesis; asm7, asm10-12, asm19 and asm21, post-PKS modifications; asm2, asm8, asm18, asm29 and asm31, regulation (Floss 2006).

producer organism. This alternative approach has the advantage of all the regulatory elements and resistance mechanisms required for ansamitocin formation being present and functional (Cassady et al. 2004).

Undoubtedly challenging prospects, with substantial problems to be overcome and much more work needed, yet bearing a considerable potential for future drug discovery, optimization and production processes (Floss 2006).

In summary, the accomplishments thus described might prove to become a vital platform and a blueprint for detailed studies of endophytic microbes *en masse* – their biosynthetic networks, evolutionary implications, host interdependencies, ecological, pharmaceutical and industrial significance – in the modern era of metabolic engineering and combinatorial biosynthesis (Stephanopoulos et al. 2004; Floss 2006).

Concluding remarks and future perspectives

Biodiversity: a precious source of novelty. Not only in terms of unravelling numerous mysteries of nature – discovering a plethora of yet undescribed species, their evolutionary backgrounds, genetics and ecology, as well as the richness of thus implied new, potentially valuable molecules; but also a revolution of thought – an expanded view promising to transform glimpses of reductionist research of the past years into snapshots of a dynamic world of systems biology, where cells grow, divide and produce, or organisms develop, differentiate and begin to deviate from the norm (Kate and Laird 2000; Stephanopoulos et al. 2004; Kayser and Quax 2007). Endophytic microbes seem to fit perfectly into this natural ‘warehouse’, only a small part of which we have been able to tap into so far.

The recent ‘genomics revolution’ has given momentum to considerable progress in the development of new technologies in bioscience, addressing specifically the arena of natural product biosynthesis. Whole-genome sequence mining (Lautru et al. 2005) and genome scanning as an alternative approach, providing an efficient way to discover natural product biosynthetic gene clusters without having the complete genome sequence (Zazopoulos et al. 2003); advances in microbial cell fermentation technology (Zengler et al. 2005; Weuster-Botzl et al. 2007) and metagenomics as a valuable alternative offering a cultivation-independent approach (Schloss and Handelsman 2005); ample successes in heterologous expression and metabolic engineering (among many others: Alper et al. 2005; Schmidt et al. 2005; Wenzel et al. 2005; DeJong et al. 2006; Julsing et al. 2006; Li et al. 2006; Lindahl et al. 2006; Nims et al. 2006; Ro et al. 2006), the latter being in fact perceived as a progenitor of functional genomics and systems biology (Stephanopoulos et al. 2004; Tyo et al. 2007) – to name only a few highlights.

As hereby reviewed, some of the aforementioned novel strategies have enabled penetration into the previously inaccessible natural-product resources: microbial endophytes. While one has to be mindful that the problem we set out to address is several orders of magnitude larger than those with which we are familiar, no one can deny the opportunities that now present themselves.

References

- Ahmed MS, Fong HH, Soejarto DD, Dobberstein RH, Waller DP, Moreno-Azorero RJ. 1981. High-performance liquid chromatographic separation and quantitation of maytansinoids in *Maytenus ilicifolia*. J Chromatogr. 213:340–344.

- Ahn IP, Lee YH. 2001. A viral double-stranded RNA up regulates the fungal virulence of *Nectria radicola*. *Mol Plant-Microbe Interact.* 14:496–567.
- Alper H, Jin YS, Moxley JF, Stephanopoulos G. 2005. Identifying gene targets for the metabolic engineering of lycopene biosynthesis in *Escherichia coli*. *Metab Eng.* 7:155–164.
- Arntz C, Tudzynski P. 1997. Identification of genes induced in alkaloid-producing cultures of *Claviceps* sp. *Curr Genet.* 31:357–360.
- Asai M, Mizuta E, Izawa M, Haibara K, Kishi T. 1978. Isolation, chemical characterization and structure of ansamitocin, a new antitumour antibiotic. *Tetrahedron.* 35:1079–1085.
- Bacon CW, Lyons PC, Porter JK, Robbins JD. 1986. Ergot toxicity from endophyte-infected grasses: a review. *Agronomy J.* 78:106–116.
- Bailey TL, Elkan C. 1994. Fitting a mixture model by expectation maximization to discover motifs in biopolymers. *Proceedings of the 2nd International Conference on Intelligent Systems for Molecular Biology.* Menlo Park, CA: AAAI Press. p. 28–36.
- de Bary A. 1866. *Morphologie und Physiologie der Pilze, Flechten, und Myxomyceten.* Vol. II. Hofmeister's Handbook of Physiological Botany. Leipzig, Germany.
- Blankenship JD, Houseknecht JB, Pal S, Bush LP, Grossman RB, Schardl CL. 2005. Biosynthetic precursors of fungal pyrrolizidines, the loline alkaloids. *Chem Biochem.* 6:1016–1022.
- Blankenship JD, Spiering MJ, Wilkinson HH, Fannin FF, Bush LP, Schardl CL. 2001. Production of loline alkaloids by the grass endophyte, *Neotyphodium uncinatum*, in defined media. *Phytochemistry.* 58: 395–401.
- Bok JW, Keller NP. 2004. LaeA, a regulator of secondary metabolism in *Aspergillus* spp. *Eukaryot Cell.* 3: 527–535.
- Bush LP, Wilkinson HH, Schardl CL. 1997. Bioprotective alkaloids of grass-fungal endophyte symbioses. *Plant Physiol.* 114:1–7.
- Butler MS. 2005. Natural products to drugs: natural product derived compounds in clinical trials. *Nat Prod Rep.* 22:162–195.
- Cassady JM, Chan KK, Floss HG, Leistner E. 2004. Recent developments in the maytansinoids antitumour agents. *Chem Pharm Bull.* 52:1–26.
- Clay K, Schardl CL. 2002. Evolutionary origins and ecological consequences of endophyte symbiosis with grasses. *Am Nat.* 160:s99–127.
- Correia T, Grammel N, Ortel I, Keller U, Tudzynski P. 2003. Molecular cloning and analysis of the ergopeptide assembly system in the ergot fungus *Claviceps purpurea*. *Chem Biol.* 10:1281–1292.
- Coyle CM, Panaccione DG. 2005. An ergot alkaloid biosynthesis gene and clustered hypothetical genes from *Aspergillus fumigatus*. *Appl Environ Microbiol.* 71:3112–3118.
- Croteau R. 2005. Taxol biosynthesis and molecular genetics. *Terpnet 2005, International Conference held at Wageningen University and Research Centre.* 2005 Apr 20–23, Wageningen, The Netherlands.
- Davis KE, Joseph SJ, Janssen PH. 2005. Effects of growth medium, inoculum size, and incubation time on culturability and isolation of soil bacteria. *Appl Environ Microbiol.* 71:826–834.
- Dawe AL, Nuss DL. 2001. Hypoviruses and chestnut blight: exploiting viruses to understand and modulate fungal pathogenesis. *Annu Rev Genet.* 35:1–29.
- DeJong JM, Liu Y, Bollon AP, Jennewein S, Williams D, Croteau R. 2006. Genetic engineering of taxol biosynthetic genes in *Saccharomyces cerevisiae*. *Biotech Bioeng.* 93:212–224.
- Diatchenko L, Lau YC, Campbell AP, Chenchik A, Moqadam F, Huang B, Lukyanov K, Gurskaya N, Sverdlov ED, Siebert PD. 1996. Suppression subtractive hybridization: a method for generating differentially regulated or tissue-specific cDNA probes and libraries. *PNAS.* 93:6025–6030.
- Diatchenko L, Lukyanov S, Lau YF, Siebert P. 1999. Suppression subtractive hybridization: a versatile method for identifying differentially expressed genes. *Meth Enzymol.* 303:349–380.
- Doolittle WF. 1999. Phylogenetic classification and the universal tree. *Science.* 284:2124–2129.
- Dymowski W, Furmanowa M. 1990. Investigating cytostatic substances in tissue of plants *Maytenus molina* in *in-vitro* cultures. Chromatographic test of extracts from callus of *Maytenus wallichiana*. *Acta Pol Pharm.* 47:51–54.
- Fleetwood DJ, Scott B, Lane GA, Tanaka A, Johnson RD. 2007. A complex ergovaline gene cluster in *Epichloë* endophytes of grasses. *Appl Environ Microbiol.* 73:2571–2579.
- Floss HG. 1976. Biosynthesis of ergot alkaloids and related compounds. *Tetrahedron.* 32:873–912.
- Floss HG. 2006. Combinatorial biosynthesis – potential and problems. *J Biotech.* 124:242–257.
- Freeman EM. 1904. The seed fungus of *Lolium temulentum* L. *Phil Trans R Soc Lond (Biol)* 196:1–27.
- Frense D. 2007. Taxanes: perspectives for biotechnological production. *Appl Microbiol Biotechnol.* 73:1233–1240.
- Gallagher RT, Hawkes AD, Steyn PS, Vlegaar R. 1984. Tremorgenic neurotoxins from perennial ryegrass causing ryegrass staggers disorder of livestock: structure elucidation of lolitrem B. *J Chem Soc Chem Commun.* 9:614–616.
- Gardner MJ, Hall N, Fung E, White O, Berriman M, Hyman RW, Carlton JM, Pain A, Nelson KE, Bowman S. 2002. Genome sequence of the human malaria parasite *Plasmodium falciparum*. *Nature.* 419:498–511.
- Gröger D, Floss HG. 1998. Biochemistry of ergot alkaloids – achievements and challenges. *Alkaloids.* 50:171–218.
- Gunatilaka AAL. 2006. Natural products from plant-associated microorganisms: distribution, structural diversity, bioactivity, and implications of their occurrence. *J Nat Prod.* 69:509–526.
- Guo B, Dai J, Ng S, Huang Y, Leong C, Ong W, Carte BK. 2000. Cytonic acids A & B: novel tridepside inhibitors of hCMV protease from the endophytic fungus *Cytospora naema* species. *J Nat Prod.* 63:602–604.
- Haarmann T, Machado C, Lübke Y, Correia T, Schardl CL, Panaccione DG, Tudzynski P. 2005. The ergot alkaloid gene cluster in *Claviceps purpurea*: extension of the cluster sequence and intra species evolution. *Phytochemistry.* 66:1312–1320.
- Haarmann T, Ortel I, Tudzynski P, Keller U. 2006. Identification of the cytochrome P450 monooxygenase

- that bridges the clavine and ergoline alkaloid pathways. *Chem Biochem.* 7:645–652.
- Harper JK, Ford EJ, Strobel GA, Arif A, Grant DM, Porco J, Tomer DP, O'Neill K. 2003. Pestacin: a 1,3-dihydro isobenzofuran from *Pestalotiopsis microspora* possessing antioxidant and antimycotic activities. *Tetrahedron.* 59:2471–2476.
- Higashide E, Asai M, Ootsu K, Tanida S, Kozay Y, Hasegawa T, Kishi T, Sugino Y, Yoneda M. 1977. Ansamitocins, a group of novel maytansinoid antibiotics with antitumour properties from *Nocardia*. *Nature.* 270:721–722.
- Hoveland C. 1993. Importance and economic significance of the *Acremonium* endophytes to performance of animals and grass plants. *Agric Ecosyst Environ.* 44:3–12.
- Jackson JA, Varney DR, Petroski RJ, Powell RG, Bush LP. 1996. Physiological responses of rats fed loline and ergot alkaloids from endophyte-infected tall fescue. *Drug Chem Toxicol.* 19:85–96.
- Jennewein S, Park H, DeJong JM, Long R, Bollon AP, Croteau R. 2004b. Coexpression in yeast of *Taxus* cytochrome P450 reductase with cytochrome P450 oxygenases involved in taxol biosynthesis. *Biotech Bioeng.* 89:588–598.
- Jennewein S, Wildung MR, Chau M, Walker K, Croteau R. 2004a. Random sequencing of an induced *Taxus* cell cDNA library for identification of clones involved in taxol biosynthesis. *PNAS.* 101:9149–9154.
- Julsing MK, Koulman A, Woerdenbag HJ, Quax WJ, Kayser O. 2006. Combinatorial biosynthesis of medicinal plant secondary metabolites. *Biomol Eng.* 23: 265–279.
- Kate KT, Laird SA, editors. 2000. The commercial use of biodiversity: access to genetic resources and benefit sharing. London, UK: Earthscan Publications Ltd.
- Kayser O, Quax WJ, editors. 2007. Medicinal plant biotechnology: from basic research to industrial applications. Weinheim, Germany: Wiley-VCH Verlag GmbH & Co. KGaA.
- Keller NP, Hohn TM. 1997. Metabolic pathway gene clusters in filamentous fungi. *Fung Gen Biol.* 21:17–29.
- Keller NP, Turner G, Bennett JW. 2005. Fungal secondary metabolism – from biochemistry to genomics. *Nat Rev Microbiol.* 3:937–947.
- Kellis M, Patterson N, Endrizzi M, Birren B, Lander ES. 2003. Sequencing and comparison of yeast species to identify genes and regulatory elements. *Nature.* 423:241–254.
- Kempken F, Kück U. 1998. Transposons in filamentous fungi – facts and perspectives. *BioEssays.* 20:652–659.
- Ketchum RE, Croteau R. 1998. Recent progress toward an understanding of taxol biosynthesis in plant cell cultures. In: Ageta H, Aimi N, Ebizuka Y, Fujita T, Honda G, editors. Towards natural medicine research in the 21st century. Proceeding of the International Symposium on Natural Medicines. Elsevier Science, Amsterdam, The Netherlands. p. 339–348.
- Ketchum RE, Gibson DM. 1996. Paclitaxel production in suspension cell cultures of *Taxus*. *Plant Cell Tiss Org Cult.* 46:9–16.
- Khosroushahi AY, Valizadeh M, Ghasempour A, Khosroushahli M, Naghdibadi H, Dadpour MR, Omid Y. 2006. Improved taxol production by combination of inducing factors in suspension cell culture of *Taxus baccata*. *Cell Biol Internat.* 30:262–269.
- Kim CG, Yu TW, Fryhle CB, Handa S, Floss HG. 1998. 3-amino-5-hydroxybenzoic acid (AHBA) synthase, the terminal enzyme in the formation of the precursor of mC₇N units in rifamycin and related antibiotics. *J Biol Chem.* 273:6030–6040.
- Klein-Marcuschamer D, Ajikumar PK, Stephanopoulos G. 2007. Engineering microbial cell factories for biosynthesis of isoprenoid molecules: beyond lycopene. *Trends Biotechnol.* 25:417–424.
- Koepp AE, Hezari M, Zajicek J, Vogel BS, LaFever RE, Lewis NG, Croteau R. 1995. Cyclization of geranylgeranyl diphosphate to taxa-4(5),11(12)-diene is the committed step of taxol biosynthesis in Pacific yew. *J Biol Chem.* 270:8686–8690.
- Kogel KH, Franken P, Hückelhoven R. 2006. Endophyte or parasite – what decides? *Curr Opin Plant Biol.* 9:358–363.
- Kucht S, Groß J, Hussein Y, Grothe T, Keller U, Basar S, König WA, Steiner U, Leistner E. 2004. Elimination of ergoline alkaloids following treatment of *Ipomoea asarifolia* (*Convolvulaceae*) with fungicides. *Planta.* 219:619–625.
- Kuldau GA, Liu JS, White JF, Siegel MR, Schardl CL. 1997. Molecular systematics of *Clavicipitaceae* supporting monophyly of genus *Epichloë* and form genus *Ephelis*. *Mycologia.* 89:431–441.
- Kupchan SM, Komoda Y, Court WA, Thomas GJ, Smith RM, Karim A, Gilmore CJ, Haltiwanger RC, Bryan RF. 1972. Maytansine, a novel antileukaemic ansa macrolide from *Maytenus ovatus*. *J Am Chem Soc.* 94:1355–1356.
- Kurland CG, Canback B, Berg OG. 2003. Horizontal gene transfer: a critical view. *PNAS.* 100:9658–9662.
- Kutil BL, Greenwald C, Liu G, Spiering MJ, Schardl CL, Wilkinson HH. 2007. Comparison of loline alkaloid gene clusters across fungal endophytes: predicting the co-regulatory sequence motifs and evolutionary history. *Fung Genet Biol.* 44:1002–1010.
- Kutney JP, Beale MH, Salisbury PJ, Stuart KL, Worth BR, Townsley PM, Chalmers WT, Nilson K, Jacoli GG. 1981. Isolation and characterization of natural products from plant tissue cultures of *Maytenus buchananii*. *Phytochemistry.* 4:653–657.
- Lam KS. 2007. New aspects of natural products in drug discovery. *Trends Microbiol.* 15:279–289.
- van Lanen SG, Shen B. 2006. Microbial genomics for the improvement of natural product discovery. *Curr Opin Microbiol.* 9:252–260.
- Lautru S, Deeth RJ, Bailey LM, Challis GL. 2005. Discovery of a new peptide natural product by *Streptomyces coelicolor* genome mining. *Nat Chem Biol.* 1:265–269.
- Lee DY, Teyssier C, Strahl BD, Stallcup MR. 2005. Role of protein methylation in regulation of transcription. *Endocr Rev.* 26:147–170.
- Lee J, Lobkovsky E, Pliam NB, Strobel GA, Clardy J. 1995. Subglutinols A & B: immunosuppressive compounds from the endophytic fungus – *Fusarium subglutinans*. *J Org Chem.* 60:7076–7077.
- Leistner E. 2005. Die Biologie der Taxane. *Pharm Unserer Zeit.* 34:98–103.
- Leuchtmann A, Schmidt D, Bush LP. 2000. Different levels of protective alkaloids in grasses with stroma-forming

- and seed-transmitted *Epichloë/Neotyphodium* endophytes. *J Chem Ecol.* 26:1025–1036.
- Li H, Sethuraman N, Stadheim TA, Zha D, Prinz B, Ballew N, Bobrowicz P, Choi BK, Cook WJ, Cukan M, Houston-Cummings NR, Davidson R, Gong B, Hamilton SR, Hoopes JP, Jiang Y, Kim N, Mansfield R, Nett JH, Rios S, Strawbridge R, Wildt S, Gerngross TU. 2006. Optimization of humanized IgGs in glycoengineered *Pichia pastoris*. *Nat Biotechnol.* 24: 210–215.
- Li JY, Sidhu RS, Bollon A, Strobel GA. 1998a. Stimulation of taxol production in liquid cultures of *Pestalotiopsis microspora*. *Mycol Res.* 102:461–464.
- Li JY, Sidhu RS, Ford EJ, Long DM, Hess WM, Strobel GA. 1998b. The induction of taxol production in the endophytic fungus *Periconia* sp. from *Torreya grandifolia*. *J Ind Microbiol Biotech.* 20:259–264.
- Li SM, Unsöld IA. 2006. Post-genome research on the biosynthesis of ergot alkaloids. *Planta Med.* 72: 1117–1120.
- Lindahl AL, Olsson ME, Mercke P, Tollbom O, Schelin J, Brodelius M, Brodelius PE. 2006. Production of the artemisinin precursor amorpha-4,11-diene by engineered *Saccharomyces cerevisiae*. *Biotechnol Lett.* 28:571–580.
- Machida M, Asai K, Sano M, Tanaka T, Kumagai T, Terai G, Kusumoto K, Arima T, Akito O, Kashiwagi Y, et al. 2005. Genome sequencing and analysis of *Aspergillus oryzae*. *Nature.* 438:1157–1161.
- Mantle PG. 1987. Secondary metabolites of *Penicillium* and *Acremonium*. In: Peberdy JF, editor. *Penicillium* and *Acremonium*. New York: Plenum Press. p. 161–243.
- Mantle PG, Weedon CM. 1994. Biosynthesis and transformation of tremorgenic indole-diterpenoids by *Penicillium paxilli* and *Acremonium lolii*. *Phytochemistry.* 36:1209–1217.
- Márquez LM, Redman RS, Rodriguez RJ, Rossinck MJ. 2007. A virus in a fungus in a plant: three-way symbiosis required for thermal tolerance. *Science.* 315:513–515.
- Martienssen RA, Colot V. 2001. DNA methylation and epigenetic inheritance in plants and filamentous fungi. *Science.* 293:1070–1074.
- Moon CD, Craven KD, Leuchtmann A, Clement SL, Schardl CL. 2004. Prevalence of interspecific hybrids amongst asexual fungal endophytes of grasses. *Mol Ecol.* 13:1455–1467.
- Naill MC, Roberts SC. 2005. Cell cycle analysis of *Taxus* suspension cultures at the single cell level as an indicator of culture heterogeneity. *Biotech Bioeng.* 90:491–500.
- National Center for Biotechnology Information (NCBI). 1988. Accessed from the website: <http://www.ncbi.nlm.nih.gov>
- Nguyen KT, Ritz D, Gu JQ, Alexander D, Chu M, Miao V, Brian P, Baltz RH. 2006. Combinatorial biosynthesis of novel antibiotics related to daptomycin. *PNAS.* 103:17462–17467.
- Nierman WC, Pain A, Anderson MJ, Wortman JR, Kim HS, Arroyo J, Berriman M, Abe K, Archer DB, Bermejo C, et al. 2005. Genomic sequence of the pathogenic and allergenic filamentous fungus *Aspergillus fumigatus*. *Nature.* 438:1151–1156.
- Nims E, Dubois CP, Roberts SC, Walker EL. 2006. Expression profiling of genes involved in paclitaxel biosynthesis for targeted metabolic engineering. *Metab Eng.* 8:385–394.
- Noh M, Yang J, Kim K, Yoon Y, Kang K, Han H, Him S, Park H. 1999. Isolation of a novel microorganism *Pestalotia heterocornis*, producing paclitaxel. *Biotech Bioeng.* 64:620–623.
- Owen NL, Hundley N. 2004. Endophytes – the chemical synthesizers inside plants. *Sci Prog.* 87:79–99.
- Panaccione DG. 2005. Origins and significance of ergot alkaloid diversity in fungi. *FEMS Microbiol Lett.* 251:9–17.
- Panaccione DG, Johnson RD, Wang J, Young CA, Damrongkool P, Scott B, Schardl CL. 2001. Elimination of ergovaline from a grass-*Neotyphodium* endophyte symbiosis by genetic modification of the endophyte. *PNAS.* 98:12820–12825.
- Panaccione DG, Tapper BA, Lane GA, Davies E, Fraser K. 2003. Biochemical outcome of blocking the ergot alkaloid pathway of grass endophyte. *J Agric Food Chem.* 51:6429–6437.
- Parker EJ, Scott DB. 2004. Indole-diterpene biosynthesis in ascomycetous fungi. In: An Z, editor. *Handbook of industrial mycology*. New York: Dekker. p. 405–426.
- Petrini O. 1996. Ecological and physiological aspects of host specificity in endophytic fungi. In: Redlin SC, Carris LM, editors. *Endophytic fungi in grasses and woody plants: systemics, ecology and evolution*. St Paul, USA: APS Press. p. 87–100.
- Petroski RJ, Yates SG, Weisleder D, Powell RG. 1989. Isolation, semi-synthesis and NMR spectral studies of loline alkaloids. *J Nat Prod.* 52:810–817.
- Powel RG, Weisleder D, Smith CR, Kozlowski J, Rohwedder WK. 1982. Treflorine, trenudine, and N-methyltrenudone: novel maytansinoids tumour inhibitors containing two fused macrocyclic rings. *J Am Chem Soc.* 104:4929–4934.
- Pullen CB, Schmitz P, Hoffmann D, Meurer K, Boettcher T, von Bamberg D, Pereira AM, de Castro França S, Hauser M, Geertsema H, van Wyk A, Mahmud T, Floss HG, Leistner E. 2003. Occurrence and non-detectability of maytansinoids in individual plants of the genera *Maytenus* and *Putterlickia*. *Phytochemistry.* 62:377–387.
- Redman RS, Sheehan KB, Stout RG, Rodriguez RJ, Henson JM. 2002. Thermotolerance generated by plant/fungal symbiosis. *Science.* 298:1581–1581.
- Riederer B, Han M, Keller U. 1996. D-Lysergyl peptide synthetase from the ergot fungus *Claviceps purpurea*. *J Biol Chem.* 271:27524–27530.
- Rinehart KL, Shield LS. 1976. Chemistry of the ansamycin antibiotics. In: Herz W, Grisebach H, Kirby GW, editors. *Progress in the chemistry of organic natural products*. New York: Springer. p. 232–300.
- Ro DK, Paradise EM, Ouellet M, Fisher KJ, Newman KL, Ndungu JM, Ho KA, Eachus RA, Ham TS, Kirby J, Chang MC, Withers ST, Shiba Y, Sarpong R, Keasling JD. 2006. Production of the antimalarial drug precursor artemisinic acid in engineered yeast. *Nature.* 440:940–943.
- Rodriguez R, Redman R. 2005. Balancing the generation and elimination of reactive oxygen species. *PNAS.* 102:3175–3176.
- Rosewich UL, Kistler HC. 2000. Role of horizontal gene transfer in the evolution of fungi. *Ann Rev Phytopathol.* 38:325–363.

- Rowan DD, Latch GC. 1994. Utilization of endophyte-infected perennial ryegrass for increased insect resistance. In: Bacon CW, White JF, editors. *Biotechnology of endophytic fungi of grasses*. Boca Raton, FL: CRC Press. p. 169–183.
- Sakai K, Ichikawa T, Yamada K, Yamashita M, Tanimoto M, Hikita A, Ijuin Y, Kondo K. 1988. Antitumour principles in mosses: the first isolation and identification of maytansinoids, including a novel 15-methoxyansamitocin P-3. *J Nat Prod*. 51:845–850.
- Schardl CL. 2001. *Epichloë festucae* and related mutualistic symbionts of grasses. *Fung Genet Biol*. 33:59–82.
- Schardl CL, Leuchtman A, Spiering MJ. 2004. Symbioses of grasses with seed-borne fungal endophytes. *Ann Rev Plant Biol*. 55:315–340.
- Schloss PD, Handelsman J. 2005. Metagenomics for studying unculturable microorganisms: cutting the Gordian knot. *Genome Biol*. 6:229–233.
- Schmidt EW, Nelson JT, Rasko DA, Sudek S, Eisen JA, Haygood MG, Ravel J. 2005. Patellamide A and C biosynthesis by a microcin-like pathway in *Prochloron didemni*, the cyanobacterial symbiont of *Lissoclinum patella*. *PNAS*. 102:7315–7320.
- Schulz B, Boyle C. 2005. The endophytic continuum. *Mycol Res*. 109:661–686.
- Selker EU, Cambareri EB, Jensen BC, Haack KR. 1987. Rearrangement of duplicated DNA in specialized cells of *Neurospora*. *Cell*. 51:741–752.
- Singh MP, Janso JE, Luckman SW, Brady SF, Clardy J, Greenstein M, Maiese WM. 2000. Biological activity of guanacastepene, a novel diterpenoid antibiotic produced by an unidentified fungus CR115. *J Antibiot*. 53:256–261.
- Singh SB, Ondeyka JG, Tsipouras N, Ruby C, Sardana V, Schulman M, Sanchez M, Pelaez F, Stahlhut MW, Munshi S, Olsen DB, Lingham RB. 2004. Hinnuliquinone, a C₂-symmetric dimeric non-peptide fungal metabolite inhibitor of HIV-1 protease. *Biochem Biophys Res Commun*. 324:108–113.
- Smith SE, Read DJ, editors. 1997. *Mycorrhizal symbiosis*. 2nd ed. San Diego: Academic Press.
- Socic H, Gaberc-Porekar V. 1992. Biosynthesis and physiology of ergot alkaloids. In: Arora DK, Elander RP, Mukerij KG, editors. *Handbook of applied mycology*. Vol. 4. Fungal biotechnology. New York: Dekker. p. 475–515.
- Spiering MJ, Moon CD, Wilkinson HH, Schardl CL. 2005. Gene clusters for insecticidal loline alkaloids in the grass-endophytic fungus *Neotyphodium uncinatum*. *Genetics*. 169:1403–1414.
- Spiering MJ, Wilkinson HH, Blankenship JD, Schardl CL. 2002. Expressed sequence tags and genes associated with loline alkaloid expression by the fungal endophyte *Neotyphodium uncinatum*. *Fung Gen Biol*. 36:242–254.
- Spiering MJ, Wilkinson HH, Schardl CL. 2007. Loline alkaloid gene clusters of the fungal endophyte *Neotyphodium uncinatum*. US Patent 7183098.
- Steiner U, Ahimsa-Müller MA, Markert A, Kucht S, Groß J, Kauf N, Kuzma M, Zych M, Lamshöft M, Furmanowa M, Knoop V, Drewke C, Leistner E. 2006. Molecular characterization of a seed transmitted clavicipitaceous fungus occurring on dicotyledonous plants (*Convolvulaceae*). *Planta*. 224:533–544.
- Stephanopoulos G, Alper H, Moxley J. 2004. Exploiting biological complexity for strain improvement through systems biology. *Nat Biotech*. 22:1261–1267.
- Steyn PS, Vleggaar R. 1985. Tremorgenic mycotoxins. *Prog Chem Org Nat Prod*. 48:1–80.
- Stierle A, Strobel GA, Stierle DB. 1993. Taxol and taxane production by *Taxomyces andreanae*, an endophytic fungus of Pacific yew. *Science*. 260:214–216.
- Stierle A, Strobel GA, Stierle DB, Grothaus P, Bignami G. 1995. The search for a taxol-producing microorganism among the endophytic fungi of the Pacific yew, *Taxus brevifolia*. *J Nat Prod*. 58:1315–1324.
- Stone JK, Bacon CW, White JF. 2000. An overview of endophytic microbes: endophytism defined. In: Bacon CW, White JF, editors. *Microbial endophytes*. New York: Marcel Dekker Inc. p. 3–29.
- Strobel GA. 2003. Endophytes as sources of bioactive products. *Micr Inf*. 5:535–544.
- Strobel GA. 2006. Harnessing endophytes for industrial microbiology. *Curr Opin Microbiol*. 9:240–244.
- Strobel GA, Daisy B. 2003. Bioprospecting for microbial endophytes and their natural products. *Microbiol Mol Biol Rev*. 67:491–502.
- Strobel GA, Daisy B, Castillo U, Harper J. 2004. Natural products from endophytic microorganisms. *J Nat Prod*. 67:257–268.
- Strobel GA, Ford E, Worapong J, Harper JK, Arif AM, Grant DM, Fung PC, Chan K. 2002. Isopestacin, an isobenzofuranone from *Pestalotiopsis microspora*, possessing antifungal and antioxidant activities. *Phytochemistry*. 60:179–183.
- Strobel GA, Hess WM, Ford E, Sidhu RS, Yang X. 1996. Taxol from fungal endophytes and the issue of biodiversity. *J Ind Microbiol*. 17:417–423.
- Strobel GA, Hess WM, Li JY, Ford E, Sears J, Sidhu RS, Summerell B. 1997. *Pestalotiopsis guepinii*, a taxol-producing endophyte of the Wollemi pine, *Wollemia nobilis*. *Austr J Bot*. 45:1073–1082.
- Strobel GA, Li JY, Sugawara F, Koshino H, Harper JK, Hess WM. 1999. Oocydin A, a chlorinated macrocyclic lactone with potent anti-oomycete activity from *Serratia marcescens*. *Microbiol*. 145:3557–3564.
- Strobel GA, Pliam NB. 1997. Immunosuppressant diterpene compound. US Patent 5648376.
- Suwanborirux K, Chang CJ, Spjut RW, Cassady JM. 1990. Ansamitocin P-3, a maytansinoid from *Claopodium crispifolium* and *Anomodon attenuatus* or associated *Actinomycetes*. *Experientia*. 46:117–120.
- Tan RX, Zou WX. 2001. Endophytes: a rich source of functional metabolites. *Nat Prod Rep*. 18:448–459.
- Tanaka A, Christensen MJ, Takemoto D, Park P, Scott B. 2006. Reactive oxygen species play a role in regulating a fungus – perennial ryegrass mutualistic interaction. *Plant Cell*. 18:1052–1066.
- Tanaka A, Tapper BA, Popay A, Parker EJ, Scott B. 2005. A symbiosis expressed non-ribosomal peptide synthetase from a mutualistic fungal endophyte of perennial ryegrass confers protection to the symbiotum from insect herbivory. *Mol Microbiol*. 57:1036–1050.
- Thompson FN, Stuedemann JA. 1993. Pathophysiology of fescue toxicosis. *Agric Ecosyst Environ*. 44:263–281.
- Tsai HF, Liu J, Staben C, Christensen MJ, Latch GC, Siegel MR, Schardl CL. 1994. Evolutionary diversification of fungal endophytes of tall fescue grass by

- hybridization with *Epichloë* species. PNAS. 91:2542–2546.
- Tsai HF, Wang H, Gebler JC, Poulter CD, Schardl CL. 1995. The *Claviceps purpurea* gene encoding dimethylallyltryptophan synthase, the committed step for ergot alkaloid biosynthesis. Biochem Biophys Res Com. 216:119–125.
- Tudzynski P, Correia T, Keller U. 2001. Biotechnology and genetics of ergot alkaloids. Appl Microbiol Biotechnol. 57:593–605.
- Tudzynski P, Hölter K, Correia T, Arntz C, Grammel N, Keller U. 1999. Evidence for an ergot alkaloid gene cluster in *Claviceps purpurea*. Mol Gen Genet. 261:133–141.
- Tyo KE, Alper HS, Stephanopoulos GN. 2007. Expanding the metabolic engineering toolbox: more options to engineer cells. Trends Biotechnol. 25:132–137.
- Unsöld IA, Li SM. 2005. Overproduction, purification and characterization of FgaPT2, a dimethylallyltryptophan synthase from *Aspergillus fumigatus*. Microbiol. 151:1499–1505.
- Volpe TA, Kidner K, Hall IM, Teng G, Grewal SI, Martienssen RA. 2002. Regulation of heterochromatic silencing and histone H3 lysine-9 methylation by RNAi. Science. 297:1833–1837.
- Vongpaseuth K, Roberts SC. 2007. Advancements in the understanding of paclitaxel metabolism in tissue culture. Curr Pharm Biotechnol. 8:219–236.
- Walker K, Croteau R. 2001. Taxol biosynthetic genes. Phytochemistry. 58:1–7.
- Walzel B, Riederer B, Keller U. 1997. Mechanism of alkaloid cyclopeptide formation in the ergot fungus *Claviceps purpurea*. Chem Biol. 4:223–230.
- Wang J, Machado C, Panaccione DG, Tsai HF, Schardl CL. 2004. The determinant step in ergot alkaloid biosynthesis by an endophyte of perennial ryegrass. Fung Gen Biol. 41:189–198.
- Wang TA, Stormo GD. 2003. Combining phylogenetic data with coregulated genes to identify regulatory motifs. Bioinformatics. 19:2369–2380.
- Wani MC, Taylor HL, Wall ME. 1973. Plant antitumour agents: colubrinol acetate and colubrinol, antileukaemic ansa macrolides from *Colubrina texensis*. J Chem Soc Chem Commun. 12:390–391.
- Wennström A. 1994. Endophyte – the misuse of an old term. Oikos. 71:535–536.
- Wenzel SC, Gross F, Zhang Y, Fu J, Stewart F, Müller R. 2005. Heterologous expression of a myxobacterial natural products assembly line in pseudomonads via Red/ET recombineering. Chem Biol. 12:349–356.
- Weuster-Botzl D, Hekmat D, Puskeiler R, Franco-Lara E. 2007. Enabling technologies: fermentation and downstream processing. Adv Biochem Eng Biotech. 105:205–247.
- Wilkie N, Wingrove PB, Bilsland JG, Young L, Harper SJ, Hefti F, Ellis S, Pollack SJ. 2001. The non-peptidyl fungal metabolite L-783,281 activates TRK neurotrophin receptors. J Neurochem. 78:1135–1145.
- Wilkinson HH, Siegel MR, Blankenship JD, Mallory AC, Bush LP. 2000. Contribution of fungal loline alkaloids to protection from aphids in a grass-endophyte mutualism. Mol Plant-Microbe Interact. 13:1027–1033.
- Wilson D. 1995. Endophyte – the evolution of a term, and clarification of its use and definition. Oikos. 73: 274–276.
- Wingender E, Dietze P, Karas H, Knüppel R. 1996. TRANSFAC: a database on transcription factors and their DNA binding sites. Nucleic Acids Res. 24:238–241.
- Wong S, Wolfe KH. 2005. Birth of a metabolic gene cluster in yeast by adaptive gene relocation. Nat Genet. 37:777–782.
- Wu J, Lin L. 2003. Enhancement of taxol production and release in *Taxus chinensis* cell cultures by ultrasound, methyl jasmonate and *in situ* solvent extraction. Appl Microbiol Biotechnol. 62:151–155.
- Young CA, Bryant MK, Cristensen MJ, Tapper BA, Bryan GT, Scott B. 2005. Molecular cloning and genetic analysis of a symbiosis-expressed gene cluster for lolitrem biosynthesis from a mutualistic endophyte of perennial ryegrass. Mol Gen Genom. 274:13–29.
- Young CA, Felitti S, Shields K, Spangenberg G, Johnson RD, Bryan GT, Saikia S, Scott B. 2006. A complex gene cluster for indole-diterpene biosynthesis in the grass endophyte *Neotyphodium lolii*. Fung Genet Biol. 43:679–693.
- Young CA, McMillan L, Telfer E, Scott B. 2001. Molecular cloning and genetic analysis of an indole-diterpene gene cluster from *Penicillium paxilli*. Mol Microbiol. 39:754–764.
- Yu TW, Bai L, Clade D, Hoffmann D, Toelzer S, Trinh KQ, Xu J, Moss SJ, Leistner E, Floss HG. 2002. The biosynthetic gene cluster of the maytansinoid antitumour agent ansamitocin from *Actinosynnema pretiosum*. PNAS. 99:7968–7973.
- Yukimune Y, Hara Y, Nomura E, Seto H, Yoshida S. 2000. The configuration of methyl jasmonate affects paclitaxel and baccatin III production in *Taxus* cells. Phytochemistry. 54:13–17.
- Zabalgoitia I, Benito EP, Ciudad AG, Criado BG, Eslava AP. 1998. Double-stranded RNA and virus-like particles in the grass endophyte *Epichloë festucae*. Mycol Res. 102:914–918.
- Zazopoulos E, Huang K, Staffa A, Liu W, Bachmann BO, Nonaka K, Ahlert J, Thorson JS, Shen B, Farnet CM. 2003. A genomics-guided approach for discovering and expressing cryptic metabolic pathways. Nat Biotech. 21:187–190.
- Zengler K, Walcher M, Clark G, Haller I, Toledo G, Holland T, Mathur EJ, Woodnutt G, Short J, Keller M. 2005. High-throughput cultivation of microorganisms using microcapsules. Meth Enzymol. 397: 124–130.
- Zhang S, Monahan BJ, Tkacz JS, Scott B. 2004. Indole-diterpene gene cluster from *Aspergillus flavus*. Appl Environ Microbiol. 70:6875–6883.
- Zhang B, Salituro G, Szalkowski D, Li Z, Zhang Y, Royo I, Vilella D, Dez M, Pelaez F, Ruby C, Kendall RL, Mao X, Griffin P, Calaycay J, Zierath JR, Heck JV, Smith RG, Moller DE. 1999. Discovery of small molecule insulin mimetic with antidiabetic activity in mice. Science. 284:974–981.
- Zhou DP, Ping XW. 2001. Study on isolation of taxol-producing fungus. J Microbiol. 21:18–20.