University of Groningen

Severity of Illness and Adaptive Functioning Predict Quality of Care of Children Among Parents with Psychotic Disorders

Campbell, L.; Hanlon, M.--C.; Cherrie, G.; Harvey, C.; Stain, H. J.; Cohen, M.; van Ravenzwaaij, D.; Brown, S. D.

Published in:
Australian and new zealand journal of psychiatry

DOI:
[10.1177/0004867417731526](https://doi.org/10.1177/0004867417731526)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Campbell, L., Hanlon, M.-C., Cherrie, G., Harvey, C., Stain, H. J., Cohen, M., van Ravenzwaaij, D., & Brown, S. D. (2018). Severity of Illness and Adaptive Functioning Predict Quality of Care of Children Among Parents with Psychotic Disorders: A Confirmatory Factor Analysis. *Australian and new zealand journal of psychiatry*, 52(5), 435-445. <https://doi.org/10.1177/0004867417731526>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Severity of illness and adaptive functioning predict quality of care of children among parents with psychosis: A confirmatory factor analysis

Australian & New Zealand Journal of Psychiatry
2018, Vol. 52(5) 435–445
DOI: 10.1177/0004867417731526

© The Royal Australian and
New Zealand College of Psychiatrists 2017
Reprints and permissions:
sagepub.co.uk/journalsPermissions.nav
journals.sagepub.com/home/anp

Linda E Campbell¹, Mary-Claire Hanlon¹,
Cherrie A Galletly^{2,3}, Carol Harvey⁴, Helen Stain⁵, Martin Cohen¹,
Don van Ravenzwaaij⁶ and Scott Brown¹

Abstract

Objective: Parenthood is central to the personal and social identity of many people. For individuals with psychotic disorders, parenthood is often associated with formidable challenges. We aimed to identify predictors of adequate parenting among parents with psychotic disorders.

Methods: Data pertaining to 234 parents with psychotic disorders living with dependent children were extracted from a population-based prevalence study, the 2010 second Australian national survey of psychosis, and analysed using confirmatory factor analysis. Parenting outcome was defined as quality of care of children, based on participant report and interviewer enquiry/exploration, and included level of participation, interest and competence in childcare during the last 12 months.

Results: Five hypothesis-driven latent variables were constructed and labelled psychosocial support, illness severity, substance abuse/dependence, adaptive functioning and parenting role. Importantly, 75% of participants were not identified to have any dysfunction in the quality of care provided to their child(ren). Severity of illness and adaptive functioning were reliably associated with quality of childcare. Psychosocial support, substance abuse/dependence and parenting role had an indirect relationship to the outcome variable via their association with either severity of illness and/or adaptive functioning.

Conclusion: The majority of parents in the current sample provided adequate parenting. However, greater symptom severity and poorer adaptive functioning ultimately leave parents with significant difficulties and in need of assistance to manage their parenting obligations. As symptoms and functioning can change episodically for people with psychotic illness, provision of targeted and flexible support that can deliver temporary assistance during times of need is necessary. This would maximise the quality of care provided to vulnerable children, with potential long-term benefits.

Keywords

Mental health, parents, psychosis, quality of care

Parenthood is a common aspiration; however, this role is difficult to fully prepare for and involves a range of emotions and practical responsibilities. For most people, the steep learning curve accompanying parenthood is a joyous, albeit at times very challenging, part of life. Of people with psychotic disorders, more than half of women and around a quarter of men are parents (Campbell et al., 2012). As can be expected, parents with serious mental illnesses including psychotic disorders have the same aspirations to parent effectively and hold comparable dreams and fears for their children as parents without serious mental illness (Ackerson and Venkataraman, 2003; Fudge et al., 2004). However, people with serious mental illness might find the parental journey even more challenging. It is important then to

¹The University of Newcastle, Newcastle, NSW, Australia

²School of Medicine, The University of Adelaide, Adelaide, SA, Australia

³Discipline of Psychiatry, University of Adelaide Ramsay Health Care (SA) Mental Health Services Northern Adelaide Local Health Network, South Australia

⁴Department of Psychiatry, The University of Melbourne, Melbourne, Victoria, Australia

⁵School of Social and Health Sciences, Leeds Trinity University, Leeds, United Kingdom

⁶University of Groningen, Groningen, The Netherlands

Corresponding author:

Linda Elisabet Campbell, Science Offices, School of Psychology, Faculty of Science, University of Newcastle, 10 Chittaway Road, Ourimbah, NSW 2258, Australia.

Email: linda.e.campbell@newcastle.edu.au

identify factors that increase the likelihood of positive child and parent outcomes.

Parents who experience a psychotic disorder become parents on average at 20 years of age, with 65% of those yet to experience their first episode of psychosis (Nicholson and Miller, 2008; Morgan et al., 2005). For those 35% whose psychotic illness precedes parenthood, managing symptoms and meeting parental role obligations place immense pressure on them, and on their family and friends (Caton et al., 1999). Women who develop serious mental illness, including psychotic disorders, around the time of their child's birth have worse parenting outcomes compared to those who had their children before mental illness onset (Mowbray et al., 2005). Yet, mothers with psychosis who have higher levels of marital and social support, and are of higher socio-economic status, tend to cope better with crises and have better parenting outcomes (Landeem et al., 2007; Morgan et al., 2012). Recent Australian clinical guidelines highlight the need for specialised care for mothers with psychosis (Galletly et al., 2016). However, in reality, many mothers with psychosis report wanting more support and often experience loneliness (Blegen et al., 2010; Hearle et al., 1999; Howard et al., 2001).

Individuals with serious mental illness report a great deal of pleasure from engagement in the parenting role (Bassett et al., 1999), although many also report considerable stress, making it challenging for them to parent well. This is unsurprising considering many non-parents and parents alike with psychotic disorders live in poverty, experience insecure living circumstances, and have difficulties completing education, maintaining employment and lack satisfying social relationships (Campbell et al., 2012; Harvey et al., 2012; Plant et al., 2002). Challenges associated with low levels of education, and subsequent problems with obtaining well-paid and flexible employment (Campbell et al., 2012), can also make it more difficult to parent, regardless of diagnosis. These stressors have been linked to poor parenting outcomes such as difficulties in forming attachments to their children and nurturing them appropriately (Oyserman et al., 2000). Indeed, mothers with psychosis appear particularly vulnerable to poor parenting outcomes compared to parents with, for example, mood disorders (Landeem et al., 2007).

For parents with psychosis, having access to support and respite in times of crisis or hospitalisation is critical (Morgan et al., 2012). However, for many people with psychosis, social isolation is a significant problem (Stain et al., 2012). Underlying the social isolation is often a pervasive sense of fear that their illness will lead to stigmatisation and custodial loss of their children to protective services or, if separated, to the other parent (Bassett et al., 1999; Zemencuk et al., 1995). Unfortunately, fear of custodial loss is not unfounded. Between 30% and 80% of mothers with psychotic disorders report having lost custody of their children (Nicholson et al., 2001). Poor insight,

a feature common among individuals with psychotic disorders, can also exacerbate parenting difficulties (Landeem et al., 2007; Pini et al., 2001) and impair many parents' ability to evaluate how well the family unit is functioning and seeking help when necessary (Plant et al., 2002).

Alcohol and other substance use is another factor known to negatively impact parenting. Among parents with psychotic disorders, substance use is common: around 70% of fathers and 45% of mothers with psychotic disorders have a lifetime substance use comorbidity (Campbell et al., 2012). Substance use increases the risk of treatment noncompliance, exacerbates pre-existing illness and impairs overall functioning and parenting capacity (Nicholson and Miller, 2008; Seeman, 2002). Additionally, substance use disorders are strongly associated with social disadvantage, violence, disruptive behaviour, and child abuse and neglect (Henshaw et al., 2011).

This study aimed to identify predictors of positive parenting outcomes for adults with psychotic disorders, using data from the second Australian national survey of psychosis (Morgan et al., 2012). The key question was which characteristics predict good versus poor parenting outcomes? The unique study design permitted the examination of quality of care provided by both mothers and fathers to their child(ren) over the last 12 months (outcome variable) and the investigation of the relationships between parenting outcomes as well as parental clinical, demographic and current psychosocial characteristics. We hypothesised that better adaptive functioning (defined as how well a person handles common demands in life; Table 3 lists the variables included in this construct), less symptomatology, low levels of substance abuse, parenting role (e.g. contact and responsibility) and strong psychosocial support would be predictive of better parenting outcomes.

Method

Design and recruitment

Data analysis was based on information collected as part of the second Australian national survey of psychosis – the Survey of High Impact Psychosis (SHIP). Briefly, a two-phase design was utilised to design a representative, population-based study covering a catchment population of ~1.5 million people aged 18–64 years (~10% of Australia's population in this age group). The survey was carried out at seven mental health service sites across five Australian states, covering a total area of roughly 62,000 km². Individuals who presented to mental health services and non-government organisations, and who screened positive for psychosis in the previous month using a specially developed psychosis screener, were included (Jablensky et al., 1999, 2000; Morgan et al., 2012). Screen items involved having hallucinations or delusions, being on antipsychotic medication or having been given a diagnosis by a doctor.

Table 1. Family composition.

	Mothers, N = 174		Fathers, N = 60		F	Alpha-value
	Mean (SD)	Range	Mean (SD)	Range		
Biological children	2.41 (1.39)	0–7	2.22 (1.33)	0–7	0.87	0.35
Stepchildren	0.22 (1.01)	0–11	0.43 (0.98)	0–4	2.04	0.16
Dependent children (including stepchildren)	1.91 (1.07)	1–8	2.23 (1.35)	1–6	3.6	0.06
Dependent children living at home	1.70 (0.96)	1–7	1.87 (0.98)	1–5	1.32	0.25
Children living elsewhere	0.21 (0.52)	0–3	0.37 (0.78)	0–3	3.20	0.08

SD: standard deviation.

The screen-positive cut-off was a positive rating on two or more of these items. The inclusion criteria identified 7955 individuals. Potential participants were randomised and stratified by age group. The final sample consisted of 1825 people who undertook interviews and assessments between April and December 2010. The study was granted ethics approval from the relevant institutional human research ethics committee at each site, and all participants provided written, informed consent. See Morgan et al. (2012) for detailed information pertaining to the study design and participants.

Participants

We report data from a subsample of the participants in the second Australian national survey of psychosis, namely, 234 (60 = men, 174 = women) parents with psychotic illness aged between 20 and 62 years (mean age = 37 years, standard deviation [SD] = 8.0 years) resident within the catchment areas. Inclusion criteria comprised living with at least one dependent child and having a role in the upbringing of their children. Exclusion criteria included the following: (1) insufficient English language abilities to complete an interview, (2) impaired cognition hindering capacity for informed consent or (3) residence in a nursing home or prison. As per the design, all participants met screening criteria for psychosis. At the time of full assessment, parents were diagnosed as per International Classification of Diseases, 10th Revision (ICD-10; World Health Organization, 2008) with non-affective (e.g. schizophrenia, non-organic psychosis [$n=138$]) or affective psychosis (e.g. bipolar disorder, schizoaffective disorder, depressive psychosis [$n=94$]). Additional participant family composition, and demographic and diagnostic information are provided in Tables 1 and 2.

Materials

Assessment included the Diagnostic Interview for Psychosis to determine ICD-10 diagnosis (Castle et al., 2006; World Health Organization, 2008). The remaining survey consisted

of 33 modules including questions about education, accommodation, income, substance use, stigma, social and occupational function, service utilisation, past and present symptoms, and intellectual ability. The Multidimensional Scale of Independent Functioning (MSIF) which has good criterion, discriminative, inter-rater and construct validity, assessed global independent functioning and reflected the participant's overall level of adaptive functioning while controlling for the level of performance in each category (Jaeger et al., 2003). Participants were asked about the number of biological- and step-children they had, how many of these were less than 18 years of age, the living and shared-care arrangements, childcare needs and contact with child protection services. Quality of care of child(ren) in the last 12 months was rated based on participant report and interviewer enquiry/exploration using a number of probes and other relevant interview information. The ratings were made by trained clinicians with regular discussions within each local site team and across the national site team to ensure high reliability of ratings. There were three quality of care ratings: no dysfunction (defined as participating in childcare about as much as the average carer of same sex/age group would under similar circumstances), obvious dysfunction (defined as having little interest in childcare or reduced competence for up to half the time) and severe dysfunction (defined as a total lack of interest in childcare or reduced competence for more than half the time or evidence of neglect). These three quality of care ratings provided the operational definition of 'parenting outcome'.

Analysis

Descriptive statistics were extracted using IBM SPSS statistical software (version 22.0; SPSS, Chicago, IL, USA). Initially, mothers and fathers were compared on a number of demographic variables (see Table 1) using one-way analyses of variance (ANOVAs) with gender as the between-group variable. Chi-square tests assessed univariate associations between nominal variables of interest (variables of interest included current housing, fortnightly

Table 2. Demographic and diagnostic characteristics of parents living with dependent children by gender.

Variables	Mothers N (%)	Fathers N (%)	Cramer's V
Marital status			0.24***
Single, never married	40 (23) ^a	6 (10) ^a	
Married or de facto	78 (44.8) ^a	43 (71.7) ^a	
Separated, divorced, widowed	56 (32.2) ^a	11 (18.3) ^a	
Education			0.14
No qualifications	52 (29.9)	17 (28.3)	
Leaving certificate	22 (12.6)	8 (13.3)	
Higher education	76 (43.7)	25 (41.7)	
Tertiary education	18 (10.3)	4 (6.7)	
Other	6 (3.4)	6 (10)	
Current housing			0.16
Own or public/private rental	108 (62)	33 (55)	
Family home (with own parents)	17 (9.8)	7 (11.8)	
Supported group accommodation	0 (0)	1 (1.7)	
Homeless	2 (1.2)	1 (1.7)	
Other including hospital, institution	2 (1.2)	1 (1.7)	
Fortnightly income			0.27**
Less than \$300 or NK	6 (3.7)	3 (5.2) ^a	
\$300–499	17 (10.6) ^a	13 (22.4)	
\$500–799	61 (37.9)	25 (43.1)	
\$800–1000	47 (29.2)	6 (10.3)	
More than \$1000	30 (18.6) ^a	11 (19.0) ^a	
Age group of onset (years)			0.03
Less than 20	50 (28.9)	16 (26.7)	
20–34	96 (55.5)	35 (58.3)	
More than 34	27 (15.6)	9 (15.0)	
Course of disorder			0.12
Single episode	16 (9.2)	2 (3.3)	
Multiple episodes	126 (72.4)	46 (76.7)	
Continuous chronic illness	32 (18.4)	12 (2)	
Overall global independent functioning			0.22**
Normal functioning	31 (17.8)	11 (18.3)	
Very mild/somewhat disabled	118 (67.8) ^a	29 (48.3) ^a	
Moderately/severe disabled	25 (14.4) ^a	20 (33.3) ^a	
Substance use disorder			–0.12*
LT alcohol abuse/dependence	75 (43.1)	34 (56.7)	
LT cannabis abuse/dependence	82 (47.1)	34 (56.7)	
LT other abuse/dependence	54 (31)	20 (33.3)	
Any children living elsewhere	28 (16.1)	13 (22)	–0.06
Childcare role			0.45***
Primary	121 (69.5) ^a	13 (21.7) ^a	
Shared equally	37 (21.3) ^a	27 (45) ^a	

(Continued)

Table 2. (Continued)

Variables	Mothers	Fathers	Cramer's V
	N (%)	N (%)	
Less than equal but substantial	13 (7.5) ^a	11 (18.3) ^a	
Minimal	3 (1.7) ^a	9 (15) ^a	
Sharing care responsibility with partner or child's other parent	46 (86.8)	44 (93.6)	0.11
Children with special needs	41 (23.6)	12 (20)	0.04
Contact with child protection services	45 (25.9)	14 (23.3)	0.03
Quality of childcare			
No dysfunction	135 (78.5)	42 (71.2)	0.14
Obvious dysfunction	33 (19.2)	12 (20.3)	
Severe dysfunction	4 (2.3)	5 (8.5)	

NK: not known; LT: lifetime.

^aAdjusted residuals equal or exceed threshold.

Significance values from chi-square analyses: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Table 3. Loadings of observed (indicator) variables on latent variables.

Latent variables	Observed variables	Loading (standard error)	p-value
Psychosocial support ^a	Overall socialising	0.759 (0.055)	<0.001
	Number of friends	-0.676 (0.047)	<0.001
	Loneliness	0.855 (0.034)	<0.001
	Personal relationships	0.529 (0.071)	<0.001
	Relationships with other people	0.929 (0.03)	<0.001
	Relationships with family	0.678 (0.07)	<0.001
	Satisfaction with help and support from case worker	0.328 (0.086)	<0.001
	Experiences of stigma or discrimination	0.445 (0.099)	<0.001
Illness severity	Age of onset	-3.305 (0.976)	<0.001
	LT symptoms of hallucinations	0.694 (0.137)	<0.001
	LT symptoms of delusions	0.421 (0.159)	<0.001
	LT symptoms of subjective thought disorder	0.669 (0.116)	<0.001
	Duration of illness	1.69 (0.849)	0.05
	Self-harm	0.538 (0.145)	<0.001
Substance abuse/dependence	LT diagnosis of alcohol abuse/dependence	0.792 (0.099)	<0.001
	LT diagnosis of cannabis abuse/dependence	0.949 (0.097)	<0.001
	LT diagnosis of other abuse/dependence	0.744 (0.093)	<0.001
Adaptive functioning	Ability to do household activities	3.158 (0.536)	<0.001
	Self-care	0.636 (0.09)	<0.001
	Residential performance	0.851 (0.029)	<0.001
	Overall global work rating	0.870 (0.028)	<0.001
	Overall global independent functioning	0.976 (0.021)	<0.001
Parenting role	Childcare involvement	-0.771 (0.176)	<0.001
	Contact with children over last 12 months	0.503 (0.159)	<0.001

LT: lifetime.

^aHigher score signifies lower levels of support.

Figure 1 is a conceptual model diagram. At the top is a box labeled "Quality Care". Below it are five ovals labeled "Parenting role", "Disability", "Substance use", "Illness", and "PayStip". At the bottom is a long row of 25 boxes representing various factors. Arrows point from "Quality Care" to each of the five middle ovals. Arrows also point from each of the five middle ovals to each of the 25 bottom boxes, indicating a direct influence of each middle variable on all bottom variables.

A structural equation model (SEM) tested hypothesised associations between groups of indicator variables and parenting outcome, operationalised as quality of care of children (last 12 months). Associations were mediated by five hypothesis generated latent variables. These were labelled as follows: psychosocial support, illness severity, substance abuse/dependence, adaptive functioning and parenting role. (Further details on the full set of variables that were included and the different models that were fit can be found in the Supplementary Material.) The observed variables that operationalised these latent variables are

The SEM model structure is illustrated in Figure 1. Each observed variable was used as an indicator for only one latent variable. All latent variables were allowed to correlate with one another and to directly associate with the outcome measure. The parameters of the model were estimated using the SEM routine in the package ‘lavaan’ for the statistical programming language R. We report parameters estimated on the scale defined by standardising at the level of latent variables. Both the R language and the lavaan package are freely available open-source software (R Development Core Team, 2004; Rosseel, 2012).

Participant characterisation

The mean number of children was 2.41 (SD, 1.39) for mothers and 2.22 (SD, 1.33) for fathers. Most parents had no children living elsewhere (78% of fathers; 84% of mothers). It was more common for mothers to rate themselves as primary caregivers compared to fathers, and fathers were more likely to have a less than equivalent level of caregiving responsibility (χ^2 (3, $n=234$)=47.51, $p<0.0005$, Cramer's $V=0.45$). When care responsibilities were shared, they were most frequently shared with a partner or the child's other parent (94% of fathers; 87% of mothers). A smaller proportion relied on the child's grandparents for help (6% of fathers; 13.2% of mothers). Of the parents who received childcare support, 82% reported that they would have some or a lot of difficulties if childcare support or help was no longer available. In total, 20% of fathers and 24% of mothers reported caring for children who had psychological or physical special needs.

Table 4. Correlation coefficients (standard error) between latent variables.

	Illness severity	Substance abuse/dependence	Adaptive functioning	Parenting role
Psychosocial support	0.407 (0.093)	0.404 (0.096)	0.430 (0.071)	0.042 (0.126)
Illness severity		0.291 (0.137)	0.430 (0.094)	-0.059 (0.156)
Substance abuse/dependence			0.141 (0.107)	0.110 (0.166)
Adaptive functioning				-0.616 (0.148)

Table 5. Loadings of latent variables onto outcome variable (quality of care).

	Regression coefficient (standard error)	z-Score	p-Value
Psychosocial support	0.103 (0.161)	0.637	0.524
Illness severity	-0.488 (0.157)	3.112	0.002
Substance abuse/dependence	0.020 (0.156)	0.128	0.899
Adaptive functioning	0.852 (0.225)	3.664	<0.001
Parenting role	0.022 (0.217)	0.103	0.981

Importantly, around three quarters of the sample were rated as having no dysfunction in their quality of care for their child(ren) (last 12 months). No relationship between quality of care and ratings of insight was identified ($p > 0.05$). Three quarters of parents had not been in contact with child protection services in the past 12 months. In total, 70% of those who had been in contact with child protection services were in the group rated as providing adequate care. In all, 17 of 40 parents, who reported that they were in contact with non-governmental organisations for various types of support, had received specific advice or support with parenting in the last 12 months and 17% of all parents reported that they had received family support from other services.

SEM

The SEM approach allows one to simplify the complex patterns of association naturally present in such a large, variable dataset. Four variables were hypothesised to influence parenting outcome. The two latent variables were reliably associated with this outcome measure, and strongly so: illness severity ($est = -0.72$, $z = -3.1$, $p < 0.01$) and adaptive functioning ($est = 1.09$, $z = 5.4$, $p < 0.001$). The two manifest variables were not reliably associated with parenting outcome (both $ps > 0.44$). The indicators that loaded onto the latent variable measuring illness severity were age of onset, duration of illness, self-harm and lifetime symptoms of hallucinations, delusions and subjective thought disorder, and the latent variable measuring substance abuse. Indicators that loaded onto the

latent variable measuring substance abuse were alcohol abuse, cannabis abuse and other abuse. Meanwhile, the ability to perform household activities, self-care, overall residential performance, overall global work rating, global independent functioning, the latent variable measuring psychosocial support and the latent variable measuring parenting role loaded onto the latent variable measuring adaptive functioning. Indicators that loaded onto the latent variable measuring psychosocial support were overall socialising, number of friends, experienced stigma or discrimination due to mental illness, perceived loneliness, personal relationships, relationships with other people, relationships with family and satisfaction with case manager. Indicators that loaded onto the latent variable measuring parenting role were responsibility of childcare role and any form of contact with (step) children.

The influence of the remaining three latent variables on parenting outcomes was mediated by the illness severity and adaptive functioning variables. These associations were detailed in the SEM's estimated correlations between the latent variables (see Table 4). The two predictive latent variables (illness severity and adaptive functioning) covaried significantly ($est = 0.65$, $z = 6.6$, $p < 0.001$). The latent variable measuring strength of psychosocial support loaded significantly on adaptive functioning ($est = 0.51$, $z = 4.9$, $p < 0.001$). The latent variable measuring parenting role loaded significantly on adaptive functioning ($est = 0.59$, $z = 2.7$, $p < 0.01$). Finally, the latent variable measuring substance abuse loaded significantly on illness severity ($est = 0.49$, $z = 2.9$, $p < 0.01$) (Table 5).

Discussion

Parents with psychotic illness experience a high cumulative rate of adversities that makes parenting very challenging. Due to the long-term consequences of poor parenting, not only for the child but also the parent, it is important to identify parents at risk of providing inadequate quality of care to their children. The most pertinent finding in this study was that the majority of parents were functioning adequately, with 75% having no reported dysfunction in the quality of care of their child(ren) over the past 12 months. Wide dissemination of these findings could reduce the effect of stigma from members of the community, health-care professionals and parents with psychotic illness. The knowledge that others in the same situation find strategies to cope and even flourish could bestow a sense of hope on parents with psychotic illness. This research provides some information about what predicted adequate parenting in the current sample.

We hypothesised that better adaptive functioning, less mental health symptomatology, low levels of substance abuse/dependence and strong psychosocial support would significantly be associated with, and predict, better quality of care of child(ren) in the last 12 months. Our hypotheses were partially supported with the two latent variables 'illness severity' and 'adaptive functioning' being significantly associated with quality of care of child(ren) in the last 12 months. These findings are important since they allow us to propose that attention in the clinical setting should continue to focus on ameliorating symptom severity and the concomitant experience of stigma, but should be fortified by a further concentration on adaptive strategies to improve everyday functioning. Each of these will be discussed in turn below.

Illness severity is a significant risk factor for poor parenting and has been identified in many studies (e.g. Mowbray et al., 2002). Consistent with the literature, the latent variable of illness severity in this study included the age at which the illness emerged (with younger illness onset associated with lower quality of care), while a longer duration of illness and lifetime presence of symptoms such as hallucinations, delusions, subjective thought disorder and self-harm were also associated with lower quality of care of children. Psychosis symptoms (and medication side effects), particularly negative symptoms, include a range of cognitive and emotional problems including lethargy, irritability, poor concentration, blunted affect, withdrawal and amotivation motivation (Evenson et al., 2008; Nicholson et al., 1998). The presence of such symptoms can make it difficult for parents to interact well with their children and to form appropriate relationships. Positive symptoms can include misinterpretation of information, and erratic and bizarre behaviours which may also adversely impact parenting (Ostler and Ackerson, 2009). It is not only important to recognise the impact that symptoms of mental illness

have on child-rearing but also imperative to be cognisant of their frequently episodic and fluctuating nature. A well-designed longitudinal study showed that when mental illness symptoms decreased, parenting stress decreased and was associated with an increase of parental adaptive behaviours including nurturance (Kahng et al., 2008).

Meanwhile, the ability to perform household activities and carry out other residential obligations, to look after oneself appropriately, to function at work and overall global independent functioning loaded onto the latent variable adaptive functioning. Parents with poor functioning are likely to have difficulties caring for a child when not provided with appropriate support. Indeed, neglect is a risk factor for removal of children from the home (Ostler and Ackerson, 2009). Negative symptoms and cognitive impairments in particular make it difficult not only to plan but also to be motivated to accomplish daily parental tasks, such as providing meals and ensuring the children have clean clothes (Nicholson and Miller, 2008). As expected, the level of adaptive functioning was moderately associated with severity of illness, suggesting that parents with psychotic illness experience more problems in performing everyday activities when in a more active or severe phase of the illness. This then has a cumulative negative effect on the responsibilities associated with child-rearing.

The literature suggests that it is important to introduce a focus on the parent-child relationship as part of adaptive functioning. Forman et al. (2007) reported that while psycho-therapeutic interventions for mothers with postnatal depression could reduce mental health symptomatology and reduce parenting stress, no improvement in the parent-child relationship was found (Forman et al., 2007). In fact, early parental negative attitudes towards the child predicted more negative child outcomes at 18 months later. This combined with findings of an increased risk of poor developmental outcomes of children of parents with psychosis strongly suggests that intervention strategies focussed on adaptive functioning need to be family-centred (Liu et al., 2015). Family-centred approaches to improve adaptive functioning can include, for example, pre- and postnatal care, parenting skills such as building parent-child relationships and effective discipline and also social cognitive skills to facilitate greater parental sensitivity and responsiveness (Liu et al., 2015). In addition, a family-centred approach can also include support to other family members with the aim of helping parents cope more effectively not only with their illness but also in their role as parents. This is important because despite strong evidence for a link between psychosocial support and parenting outcomes in the literature (Chernomas and Clarke, 2003; Evenson and Simon, 2005), our results did not show a direct association between psychosocial support and quality of care for children. Instead, psychosocial support was associated with severity of illness and adaptive functioning, suggesting psychosocial support indirectly

influences quality of parenting in the context of severity and adaptive functioning. Specifically and alarmingly, the results show that parents with the highest needs reported the lowest levels of psychosocial support. Indeed, parents with better psychosocial networks, including marital and social relationships, are more likely to have good parenting outcomes (Abel et al., 2005) and have been found to function better in times of crisis compared to socially isolated parents and to be less likely to lose custody of their children (Ackerson, 2003). Conversely, parenting stress is reportedly higher for mothers when they experience less psychosocial support and more social contextual risk factors such as frequent daily hassles (Kahng et al., 2008). Research on the general population suggests parents who have good psychosocial support have more adaptive interactions with their children and stronger parenting skills. Taken together with our findings, the growing recognition of the importance of psychosocial support for parents suggests greater need for longitudinal studies that model interrelationships between psychosocial support and other factors relevant to parenting. An interesting, potentially related finding was that the majority of parents who had been in touch with child protection services over the past year were rated as providing adequate care for their children. Unfortunately, the nature of the available data did not allow for a closer examination of this finding, but there are several different explanations that might be possible. First, it is possible that the self-ratings are biased and that parents failed to recognise their own inadequacies, although careful interviewing of the parents using the established protocol should have minimised this bias. Second, it might also be so that contact with the child protection services either directly or indirectly (through partner organisations) provides the parent with help and assistance necessary to provide better care for their child, that is, the child protection services act as a psychosocial support structure.

The study also included other variables shown to be of interest to parenting outcomes in the general population. For example, substance misuse is generally recognised as a strong risk factor for poor parenting outcomes in the general population (Ostler and Ackerson, 2009) and occurred at a high rate in the current sample; however, abuse/dependence was only indirectly linked to parenting outcome through weak association with illness severity. Substance misuse potentially influences parenting outcomes for people with substance use disorders through other pathways. Indeed, the relationship between substance abuse/dependence and psychosocial support was moderate, indicating a risk of poor psychosocial networks for people with substance use disorders may be a significant factor to consider. Finally, the participants' parenting role, that is, the amount of contact with children over the past 12 months and their childcare involvement, was also not associated directly with quality of care. However, it was significantly associated with adaptive functioning such that parents with lower

levels of functioning did not interact as much with their children as parents with a higher level of functioning. This underscores the importance of parental functioning, indicating that treatment efforts aimed at improving parents' functioning are also likely to benefit their children.

One of the strengths of this study of parenting is that it involved a large representative sample of people, receiving services or known to service providers, with psychotic disorders. However, due to the cross-sectional nature and diverse range of enquiry within the second Australian national survey of psychosis, the items mapping specifically onto the parenting section lacked sufficient detail to comprehensively explore the parenting experiences of the participants. For instance, it would have been preferable to have more detailed information regarding the parent and child and more than a single outcome to measure quality of parenting. Although the ratings were based on participant self-report and interviewer enquiry/exploration using probes and other relevant interview information, a more objective measure of parenting such as observational home-visits or dyadic interactions would have added value to this study.

Our findings underscore a growing body of literature stating that parents with psychotic illness are in high need of specifically targeted interventions addressing issues such as practical and social support and stress reduction related to parenting challenges, which commonly trigger illness relapse for many parents (Nicholson et al., 1998). Some clinical services may not consider the parenting status of clients and care is often designed to a 'one-size-fits-all' model (Fudge et al., 2004). Greater understanding of predictive factors for parenting difficulties and poor parenting outcomes would assist to improve targeting of appropriate and effective supports for parents and their children. Interventions should aim to include parenting techniques and also work on improving the parent-child relationship directly (Forman et al., 2007). However, while the emphasis on interventions and, by extension, research is most often centred of looking at the experiences of parents who are not doing well, our findings of a large cohort of parents who are doing well suggest that we also need to further explore their experiences.

To conclude, regardless of formidable challenges, parents with psychotic illness value their role as parents and being a parent is a critical part of their identity and daily lives. Our results indicate that there are many parents with psychotic illnesses who are doing well; however, the quality of their parenting is directly affected by their illness severity and daily functioning. Psychosocial support and substance abuse/dependence may indirectly influence the parenting ability of an adult with serious mental illness. In addition, financial hardship is present for many of these parents making it even more challenging to parent well. It is important to acknowledge that the support needs of parents with psychotic illness are likely to change episodically,

and therefore, treatment plans should be flexible and targeted to times of greater needs. Consensus guidelines on care for people with psychosis including specific antenatal care recommendations can assist clinicians (Galletly et al., 2016; McCauley-Elsom and Kulkarni, 2007). However, it is important to not only consider mothers but also fathers. Indeed, parenthood, regardless of gender, needs to be viewed as more central to recovery from mental illness, along with connectedness, empowerment, future-centred hope and optimism as part of finding meaning and purpose in life (Tew et al., 2011). This will entail providing more effective and better-targeted treatment and support to parents with psychotic illness.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship and/or publication of this article.

References

- Abel K, Webb R, Salmon M, et al. (2005) Prevalence and predictors of parenting outcomes in a cohort of mothers with schizophrenia admitted for joint mother and baby psychiatric care in England. *Journal of Clinical Psychiatry* 66: 781–789.
- Ackerson B (2003) Coping with the dual demands of severe mental illness and parenting: The parents' perspective. *Families in Society: The Journal of Contemporary Social Services* 84: 109–118.
- Ackerson B and Venkataraman M (2003) Issues in parenting by clients with severe and persistent mental illness: A survey of experts. *Journal of Family Social Work* 7: 35–52.
- Bassett H, Lampe J and Lloyd C (1999) Parenting: Experiences and feelings of parents with a mental illness. *Journal of Mental Health* 8: 597–604.
- Blegen N, Hummelvoll J and Severinsson E (2010) Mothers with mental health problems: A systematic review. *Nursing & Health Sciences* 12: 519–528.
- Campbell LE, Hanlon M-C, Poon A, et al. (2012) The experiences of Australian parents with psychosis: The second Australian national survey of psychosis. *Australian and New Zealand Journal of Psychiatry* 46: 890–900.
- Castle D, Jablensky A, Mcgrath J, et al. (2006) The Diagnostic Interview for Psychosis (DIP): Development, reliability and applications. *Psychological Medicine* 36: 69–80.
- Caton C, Courmos F and Dominguez B (1999) Parenting and adjustment in schizophrenia. *Psychiatric Services* 50: 239–243.
- Chernomas W and Clarke D (2003) *Social Support and Women Living with Serious Mental Illness*. Winnipeg, MB: Prairie Women's Health Centre of Excellence.
- Evenson E, Rhodes J, Feigenbaum J, et al. (2008) The experiences of fathers with psychosis. *Journal of Mental Health* 17: 629–642.
- Evenson R and Simon R (2005) Clarifying the relationship between parenthood and depression. *Journal of Health and Social Behavior* 46: 341–358.
- Forman DR, O'Hara MW, Stuart S, et al. (2007) Effective treatment for postpartum depression is not sufficient to improve the developing mother-child relationship. *Development and Psychopathology* 19: 585–602.
- Fudge E, Falkov A, Kowalenko N, et al. (2004) Parenting is a mental health issue. *Australasian Psychiatry* 12: 166–171.
- Galletly C, Castle D, Dark F, et al. (2016) Royal Australian and New Zealand College of Psychiatrists clinical practice guidelines for the management of schizophrenia and related disorders. *Australian and New Zealand Journal of Psychiatry* 50: 410–472.
- Harvey C, Killackey E, Groves A, et al. (2012) A place to live: Housing needs for people with psychotic disorders identified in the second Australian national survey of psychosis. *Australian and New Zealand Journal of Psychiatry* 46: 840–850.
- Hearle J, Plant K, Jenner L, et al. (1999) A survey of contact with offspring and assistance with child care among parents with psychotic disorders. *Psychiatric Services* 50: 1354–1356.
- Henshaw C, Adshead G, Bende B, et al. (2011) *Parents as Patients: Supporting the Needs of Patients Who Are Parents and Their Children*. London: Royal College of Psychiatrists.
- Howard L, Kumar R and Thornicroft G (2001) Psychosocial characteristics and needs of mothers with psychotic disorders. *The British Journal of Psychiatry* 178: 427–432.
- Jablensky A, McGrath J, Herrman H, et al. (1999) *People Living with Psychotic Illness: An Australian Study 1997–1998*. Canberra, ACT, Australia: Commonwealth of Australia.
- Jablensky A, McGrath J, Herrman H, et al. (2000) Psychotic disorders in urban areas: An overview of the Study on Low Prevalence Disorders. *Australian and New Zealand Journal of Psychiatry* 34: 221–236.
- Jaeger J, Berns S and Czobor P (2003) The multidimensional scale of independent functioning. *Schizophrenia Bulletin* 29: 153–168.
- Kahng S, Oyserman D, Bybee D, et al. (2008) Mothers with serious mental illness: When symptoms decline does parenting improve? *Journal of Family Psychology* 22: 162.
- Landeen J, Seeman M, Goering P, et al. (2007) Schizophrenia: Effect of perceived stigma on two dimensions of recovery. *Clinical Schizophrenia & Related Psychoses* 1: 64–68.
- Liu CH, Keshavan MS, Tronick E, et al. (2015) Perinatal risks and childhood premorbid indicators of later psychosis: Next steps for early psychosocial interventions. *Schizophrenia Bulletin* 41: 801–816.
- McCauley-Elsom K and Kulkarni J (2007) Managing psychosis in pregnancy. *Australian and New Zealand Journal of Psychiatry* 41: 289–292.
- Morgan V, Mitchell P and Jablensky A (2005) The epidemiology of bipolar disorder: Sociodemographic, disability and service utilization data from the Australian National Study of Low Prevalence (Psychotic) Disorders. *Bipolar Disorder* 7: 326–337.
- Morgan V, Waterreus A, Jablensky A, et al. (2012) People living with psychotic illness in 2010: The second Australian national survey of psychosis. *Australian and New Zealand Journal of Psychiatry* 46: 735–752.
- Mowbray C, Bybee D, Oyserman D, et al. (2005) Timing of mental illness onset and motherhood. *Journal of Nervous and Mental Disease* 193: 369–378.
- Mowbray C, Oyserman D, Bybee D, et al. (2002) Parenting of mothers with a serious mental illness: Differential effects of diagnosis, clinical history, and other mental health variables. *Social Work Research* 26: 225–240.
- Nicholson J and Miller L (2008) Parenting. In: Meuser KT and Jeste DV (eds) *Clinical Handbook of Schizophrenia*. New York: The Guildford Press. 471–480.
- Nicholson J, Biebel K, Hinden B, et al. (2001) *Critical Issues for Parents with Mental Illness and Their Families*. Rockville, MD: Center for Mental Health Services, Substance Abuse and Mental Health Services Administration.
- Nicholson J, Sweeney E and Geller J (1998) Mothers with mental illness: I. The competing demands of parenting and living with mental illness. *Psychiatric Services* 49: 635–642.

- Ostler T and Ackerson B (2009) Parental chronic mental illness. In: Haith MM and Benson JB (eds) *Diseases and Disorders in Infancy and Early Childhood*. Amsterdam: Academic Press, pp. 293–304.
- Oyserman D, Mowbray C, Meares P, et al. (2000) Parenting among mothers with a serious mental illness. *American Journal of Orthopsychiatry* 70: 296–315.
- Pini S, Cassano G, Dell’Osso L, et al. (2001) Insight into illness in schizophrenia, schizoaffective disorder, and mood disorders with psychotic features. *American Journal of Psychiatry* 158: 122–125.
- Plant K, Byrne L, Barkla J, et al. (2002) Parents with psychosis: A pilot study examining self-report measures related to family functioning. *Advances in Mental Health* 1: 38–48.
- R Development Core Team (2004) *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing.
- Rosseel Y (2012) lavaan: An R package for structural equation modeling. *Journal of Statistical Software* 48: 1–36.
- Seeman MV (2002) Women with schizophrenia as parents. *Primary Psychiatry* 9: 39–42.
- Stain H, Galletly C, Clark S, et al. (2012) Understanding the social costs of psychosis: The experience of adults affected by psychosis identified within the second Australian national survey of psychosis. *Australian and New Zealand Journal of Psychiatry* 46: 879–889.
- Tew J, Ramon S, Slade M, et al. (2011) Social factors and recovery from mental health difficulties: A review of the evidence. *British Journal of Social Work* 42: 443–460.
- World Health Organization (2008) *ICD-10: International Statistical Classification of Diseases and Related Health Problems*. New York: World Health Organization.
- Zemencuk J, Rogosch F and Mowbray C (1995) The seriously mentally ill woman in the role of parent: Characteristics, parenting sensitivity, and needs. *Psychosocial Rehabilitation Journal* 18: 77–92.