University of Groningen

Multiple observations of the sea anemone *Phyllodiscus semoni* Kwietniewski, 1897 (Actiniaria: Aliciidae) from Sabah, Borneo represent first records for Malaysia

van der Meij, Sancia E. T.; Draisma, Stefano G. A.; Waheed, Zarinah

Published in:
Regional studies in marine science

DOI:
[10.1016/j.rsma.2018.02.003](https://doi.org/10.1016/j.rsma.2018.02.003)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

van der Meij, S. E. T., Draisma, S. G. A., & Waheed, Z. (2018). Multiple observations of the sea anemone *Phyllodiscus semoni* Kwietniewski, 1897 (Actiniaria: Aliciidae) from Sabah, Borneo represent first records for Malaysia. *Regional studies in marine science*, 18, 135-138. <https://doi.org/10.1016/j.rsma.2018.02.003>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Multiple observations of the sea anemone *Phyllodiscus semoni* Kwietniewski, 1897 (Actiniaria: Aliciidae) from Sabah, Borneo represent first records for Malaysia

Sancia E.T. van der Meij^{a,b,c,*}, Stefano G.A. Draisma^d, Zarinah Waheed^{b,e}

^a Oxford University Museum of Natural History, University of Oxford, Parks Road, Oxford OX1 3PW, United Kingdom

^b Naturalis Biodiversity Center, PO Box 9517, 2300 RA, Leiden, The Netherlands

^c Groningen Institute for Evolutionary Life Sciences, University of Groningen, PO Box 11103, 9700 CC Groningen, The Netherlands

^d Excellence Center for Biodiversity of Peninsular Thailand, Prince of Songkla University, Hat Yai, Songkhla 90110, Thailand

^e Endangered Marine Species Research Unit, Borneo Marine Research Institute, Universiti Malaysia Sabah, Jalan UMS, 88400 Kota Kinabalu, Sabah, Malaysia

ARTICLE INFO

Article history:

Received 13 November 2017

Received in revised form 8 February 2018

Accepted 8 February 2018

Available online 14 February 2018

Keywords:

Coral reefs

Distribution

Marine biodiversity

Morphotype

ABSTRACT

The stinging sea anemone *Phyllodiscus semoni* is recorded from Malaysia for the first time. This species exhibits high morphological variation, and morphotypes resembling dead coral rock, soft and stony corals and algae have been documented. Correct identification and information on local occurrences of *Phyllodiscus* is important, since this species is armed with venom-laden nematocysts that are hazardous to humans. In situ photographs of the specimens encountered in Malaysian Borneo are provided and their morphotypes are discussed.

© 2018 Elsevier B.V. All rights reserved.

1. Introduction

The sea anemone *Phyllodiscus semoni* Kwietniewski, 1897 is an iconic inhabitant of Indo-West Pacific coral reefs. This species is a master of camouflage and mimicry, not only in colour and pattern, but also in shape and form. Many morphotypes have been recorded for this species including morphs resembling dead coral rock, soft and stony corals, and algae (Hoeksema and Crowther, 2011). The large variety in morphotypes complicates identification of this species, and records of *P. semoni* are still rather sparse (Den Hartog, 1997; Hoeksema and Crowther, 2011). Information about the regional occurrence of this species is important; *Phyllodiscus* possess some of the most dangerous venoms for humans. Their sting can induce severe dermatitis and, rarely, renal failure (Mizuno et al., 2007).

Known records of *P. semoni* in the Indo-West Pacific include: Ari Atoll in the Maldives (Erhardt and Knop, 2005), various locations in the Philippines (Gosliner et al., 1996) and Indonesia (Gosliner et al., 1996; Den Hartog, 1997; Coleman, 2000; Erhardt and Knop, 2005; Hoeksema and Crowther, 2011), Milne Bay in Papua New Guinea (Halstead, 2000), Okinawa in Japan (Mizuno et al., 2007),

off Townsville in Australia (Shick et al., 1991), and New Caledonia (Laboute and de Forges, 2004; listed as *Alicia rhadina* Haddon and Shackleton, 1893). In addition there is a possible record from southern Vietnam (see discussion in Hoeksema and Crowther, 2011). Missing from this list of records is Malaysia, a country home to large, diverse coral reefs and part of the so-called Coral Triangle (Affendi and Rosman, 2011). Here we report the first observations of *P. semoni* in Malaysian waters and discuss their morphotypes (Fig. 1).

2. Material and methods

The East Malaysia state of Sabah contains more than 75% of Malaysian reefs (Burke et al., 2002), and the highest concentration of reefs can be found in the north and southeast of Sabah. Observations of *P. semoni* were made during the Semporna Marine Ecological Expedition (SMEE) 2010, and a reconnaissance trip to Semporna in 2009 prior to SMEE 2010 in southeast Sabah, as well as during the Tun Mustapha Park Expedition (TMPE) 2012 in north Sabah (Fig. 2). Background information on the expeditions, including detailed maps and localities visited can be found in Van der Meij and Hoeksema (2013), Waheed and Hoeksema (2013) and Waheed et al. (2015). *P. semoni* specimens were photographed when encountered during the research dives, but no specimens were collected.

* Corresponding author.

E-mail address: Sancia.vanderMeij@naturalis.nl (S.E.T. van der Meij).

Fig. 1. *Phyllodiscus semoni* observations in Malaysian Borneo. **A** a *P. semoni* morphotype resembling *Briareum* soft coral at Darby Bank (Semporna area), **B** an algae covered dead coral rock morphotype at Simaggot (Kudat area), **C–D** *P. semoni* morphotypes resembling *Litophyton* soft coral at Belaruan (Kudat area) (note the blue *Porites* coral in the centre of the image C (arrowed)), **E** a *Sinularia* soft coral mimic at Tanjung Pantau Pantau (Semporna area), **F** a *P. semoni* specimen resembling a green algae morph at Sibaliu N (Kudat area). Photo credits: **A** Z Waheed, **B–C, F** SET van der Meij, **D** BT Reijnen, **E** SGA Draisma.

3. Results and discussion

We recorded *P. semoni* from five localities in Sabah, Borneo, which represent the first published accounts of this species in Malaysia. Three new records of *P. semoni* come from the Kudat area at the northern tip of Borneo in the Sulu Sea (Fig. 2(A)). The waters north of the town of Kudat are part of a marine park, Tun Mustapha Park (TMP), gazetted in 2016. TMP is home to the second largest coral reef area in Sabah, and has high levels of marine biodiversity (Waheed et al., 2015). New species have been discovered in the area in recent years (Chen, 2015; Van der Meij, 2017), highlighting the importance of the marine park.

In TMP *P. semoni* was encountered at three localities: Simaggot (07°02'34"N 117°27'58"E), Sibaliu N (07°11'33"N 117°23'38"E) in the east of the park, and Belaruan (07°01'50"N 117°00'41"E) within the South Banggi Channel in between Banggi Island and the mainland. The single specimen at Simaggot resembles algae covered dead coral rock (Fig. 1(B), compare fig. 7E in Hoeksema and Crowther, 2011), whereas the single specimen at Sibaliu N is a green algae morph (Fig. 1(F), compare p. 229 in Erhardt and Knop, 2005). The latter is reminiscent of a dead branching coral covered in green turf algae. The more common green turf algae found on reefs include *Bryopsis* J.V. Lamouroux (1809) (usually *B. pennata* J.V. Lamouroux, 1809), *Caulerpa verticillata* J. Agardh (1847), and

Chlorodesmis W.H. Harvey and J.W. Bailey (1851) species (most notably *C. fastigiata* (C. Agardh) S.C. Ducker (1969) and *C. hildebrandtii* A. Gepp and E.S. Gepp (1911)). On damaged reefs blooms have been reported for the genus *Trichosolen* Montagne (1861) as well (Pauly et al., 2011). The *P. semoni* morph at Sibaliu N is similar in appearance to these common turf algae species. At Belaruan three different sized, bluish *P. semoni* specimens were observed (Fig. 1(C)–(D)), which could be mistaken for *Litophyton* Forskål (1775) soft coral. The blue coral in the centre of the picture (Fig. 1(C), arrowed) is a *Porites* Link (1807) stony coral. All three TMP sites at which *P. semoni* was discovered are exposed to the dominant wind direction. Simaggot and Sibaliu N are rather shallow with a maximum depth of ca. 20 m depth, whereas Belaruan reaches at least 30 m depth. The latter locality is located in the channel between Banggi Island and the mainland where strong currents can be present.

Two of the new records of *P. semoni* are observations from the Semporna area in the southeast of Sabah (Fig. 2(B)). Semporna is located in the Sulawesi Sea and is known for its remarkably high marine biodiversity (Waheed and Hoeksema, 2013). Parts of the Semporna reefs are located in the Tun Sakaran Marine Park (TSMP), which was gazetted in 2004. A specimen of *P. semoni* with a morphotype resembling a *Briareum*

Fig. 2. Map of Sabah, Malaysian Borneo (inset). **A** Kudat and **B** Semporna area showing the localities where *Phyllo-discus semoni* morphotypes were encountered.

de Blainville (1834) soft coral (Fig. 1(A), compare fig. 6E in Hoeksema and Crowther, 2011) was observed on Darby Bank (south-east of Tawau, 04°06'45"N 118°13'30"E) in 2009, and another *P. semoni* specimen resembling a *Sinularia* May, 1898 soft coral (Fig. 1(E)) was observed at Tanjung Pantau Pantau on Bumbun Island (04°26'54"N 118°46'31"E). Darby Bank is part of a barrier reef system south of Semporna, and Tanjung Pantau Pantau is south of TSMP. Both localities are exposed, offshore sites.

There are no published records of *P. semoni* from Peninsular Malaysia, despite several research dives on both coasts by the authors. In an overview of the anemone fauna of Singapore, just south of Peninsular Malaysia, *P. semoni* is also not recorded (Fautin et al., 2015). The fieldwork in Singapore was, however, based around intertidal areas, where it is uncommon to find *Phyllo-discus*. In order to better understand the distribution of this enigmatic sea anemone further observations are needed, especially from localities west of Malaysian Borneo.

Acknowledgments

The reconnaissance trip to Semporna in 2009 was sponsored by WWF-Malaysia. Both expeditions (SMEE 2010 and TMPE 2012) were co-organised by WWF-Malaysia, Universiti Malaysia Sabah and Naturalis Biodiversity Center, the Netherlands (and Universiti Malaya for SMEE 2010 and Sabah Parks for TMPE 2012). Research permission was granted by Sabah Parks, Sabah Fisheries Department and the Economic Planning Unit, Malaysia. SMEE 2010 was funded through WWF-Malaysia, whereas TMPE 2012 was funded

by the Ministry of Science, Technology and Innovation (MOSTI) and USAID CTSP (Coral Triangle Support Partnership). The first author received additional funding from the Jan Joost ter Pelkwijk-Fonds (Naturalis) for SMEE 2010 and from the TREUB-maatschappij (Society for the Advancement of Research in the Tropics) for TMPE 2012. We appreciate discussing the soft coral look-alikes and photograph contribution from Bastian Reijnen (Naturalis Biodiversity Center). We thank two anonymous reviewers for their constructive comments, which helped to improve the manuscript.

References

- Affendi, Y.A., Rosman, F.R., 2011. Current knowledge on scleractinian coral diversity of Peninsular Malaysia. In: Kamarruddin, I., Mohamed, C.A.R., Rozaimi, M.J., Kee Alfian, A.A., Fitra, A.Z., Lee, J.N. (Eds.), *Malaysia's Marine Biodiversity: Inventory and Current Status*. Department of Marine Park Malaysia, Putrajaya, pp. 21–31.
- Agardh, J.G., 1847. Nya alger från Mexico. Öfversigt af Kongl Vetenskaps-Akademiens Förhandlingar, Stockholm, pp. 5–17.
- Burke, L., Selig, E., Spalding, M., 2002. *Reefs at risk in Southeast Asia*. World Resources Institute, Washington DC.
- Chen, C.A., 2015. A new marine nematode genus *Pseudoplatycoma* with a new species from the Sulu Sea and revision of the subfamily Platycominae (Enoplida: Leptosomatidae). *Zootaxa* 3905, 233–244.
- Coleman, N., 2000. *Marine life of the Maldives*. Atoll Editions, Apollo Bay.
- de Blainville, H.M., 1834. *Manuel d'actinologie ou de zoophytologie*. 100 pls. Paris, p. 688.
- Den Hartog, J.C., 1997. The sea anemone fauna of Indonesian coral reefs. In: Tomascik, T., Mah, A.J., Nontji, A., Moosa, M.K. (Eds.), *The Ecology of the Indonesian Seas 1*. Periplus Editions, Singapore, pp. 351–370.

- Ducker, S.C., 1969. Additions to the genus *Chlorodesmis* (Chlorophyta). *Phycologia* 8, 17–20.
- Erhardt, H., Knop, D., 2005. Corals: Indo-Pacific field guide. IKAN Unterwasserarchiv, Frankfurt.
- Fautin, D.G., Tan, R., Liang, W., Yap, N., Hee, T.S., Crowther, A., Goodwill, R., Sanpanich, K., Chieh, T.Y., 2015. Sea anemones (Cnidaria: Actiniaria) of Singapore: shallow-water species known also from the Indian subcontinent. *Raff. Bul. Zool. Suppl.* 31, 44–59.
- Forskål, P., 1775. Descriptiones animalium, avium, amphibiorum, piscium, insectorum, vermium/quae in itinere orientali observavit Petrus Forskål. Post mortem auctoris edidit Carsten Niebuhr. Adjuncta est materia medica kahirina atque tabula maris Rubri geographica. Hauniæ [Copenhagen]:ex officina Mölleri.
- Gepp, A., Gepp, E.S., 1911. The Codiaceae of the Siboga Expedition Including a Monograph of Flabellariae and Udoteae. In: *Siboga-Expeditie Monographie*, vol. 62, E.J. Brill, Leiden, pp. 1–150.
- Gosliner, T.M., Behrens, D.W., Williams, G.C., 1996. Coral Reef Animals of the Indo-Pacific. Sea Challengers, Monterey.
- Halstead, B., 2000. Coral Sea Reef Guide. Sea Challengers, Danville.
- Harvey, W.H., Bailey, J.W., 1851. Descriptions of seventeen new species of algae collected by the United States Exploring Expedition. *Proc. Boston Soc. Nat. Hist.* 3, 370–373.
- Hoeksema, B.W., Crowther, A.L., 2011. Masquerade, mimicry and crypsis of the polymorphic sea anemone *Phyllodiscus semoni* and its aggregations in South Sulawesi. *Contrib. Zool.* 80, 251–268.
- Laboute, P., de Forges, B.R., 2004. Lagons et récifs de Nouvelle-Calédonie. Catherine Ledru.
- Lamouroux, J.V.F., 1809. Observations sur la physiologie des algues marines, et description de cinq nouveaux genres de cette famille. *Nouv. Bull. Sci. Soc. Philomathique Paris* 1, 330–333.
- Link, D.H.F., 1807–1808. Beschreibung der Naturalien-Sammlung der Universität zu Rostock. Rostock, Adlers Erben. 1 Abt. [Part 1], pp. 1–50; 2 Abt. [Part 2], pp. 51–100; 3 Abt. [Part 3], pp. 101–165; 4 Abt. [Part 4], pp. 1–30; 5 Abt. [Part 5], pp. 1–38 [1808]; 6 Abt. [Part 6], pp. 1–38.
- May, W., 1898. Die von Dr. Stuhlmann im Jahre 1889 gesammelten ostafrikanischen Alcyonaceen des Hamburger Museums. *Jahrb. Hamburg Wissensch. Anstalten* 15 (Beiheft 2), 1–38.
- Mizuno, M., Nozaki, M., Morine, N., Suzuki, N., Nishikawa, K., Morgan, B.P., Matsuo, S., 2007. A protein toxin from the sea anemone *Phyllodiscus semoni* targets the kidney and causes a severe renal injury with predominant glomerular endothelial damage. *Am. J. Path.* 171, 402–414.
- Montagne, C., 1861. Neuvième centurie de plantes cellulaires nouvelles, tant indigènes qu'exotiques. *Décades I et II. Ann. Sci. Nat. Bot. Ser.* 4 14, 167–185.
- Pauly, K., Jupp, B.P., De Clerck, O., 2011. Modelling the distribution and ecology of *Trichosolen* blooms on coral reefs worldwide. *Mar. Biol.* 158, 2239–2246.
- Shick, J.M., Lesser, M.P., Stochaj, W.R., 1991. Ultraviolet radiation and photooxidative stress in zooxanthellate Anthozoa: the sea anemone *Phyllodiscus semoni* and the octocoral *Clavularia* sp. *Symbiosis* 10, 145–173.
- Van der Meij, S.E.T., 2017. The coral genus *Caulastrea* Dana, 1846 (Scleractinia, Merulinidae) as a new host for gall crabs (Decapoda, Cryptochiridae), with the description of *Lithoscaptus tuerkayi* sp. nov. *Crustaceana* 90, 1027–1038.
- Van der Meij, S.E.T., Hoeksema, B.W., 2013. Distribution of gall crabs inhabiting mushroom corals on Semporna reefs, Malaysia. *Mar. Biodiv.* 43, 53–59.
- Waheed, Z., Hoeksema, B.W., 2013. A tale of two winds: species richness patterns of reef corals around the Semporna peninsula, Malaysia. *Mar. Biodiv.* 43, 37–51.
- Waheed, Z., van Mil, H.G.J., Syed Hussein, M.A., Jumin, R., Golam Ahad, B., Hoeksema, B.W., 2015. Coral reefs at the northernmost tip of Borneo: an assessment of scleractinian species richness patterns and benthic reef assemblages. *Plos One* 10, e0146006.