University of Groningen

Multiplex Ligation-Dependent Probe Amplification of Uveal Melanoma

Damato, Bertil; Dopierala, Justyna; Klaasen, Annelies; van Dijk, Marcory; Sibbring, Julie; Coupland, Sarah E.

Published in:
Investigative ophthalmology & visual science

DOI:
[10.1167/iovs.08-3165](https://doi.org/10.1167/iovs.08-3165)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2009

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Damato, B., Dopierala, J., Klaasen, A., van Dijk, M., Sibbring, J., & Coupland, S. E. (2009). Multiplex Ligation-Dependent Probe Amplification of Uveal Melanoma: Correlation with Metastatic Death. *Investigative ophthalmology & visual science*, 50(7), 3048-3055. <https://doi.org/10.1167/iovs.08-3165>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Multiplex Ligation-Dependent Probe Amplification of Uveal Melanoma: Correlation with Metastatic Death

Bertil Damato,¹ Justyna Dopierala,² Annelies Klaasen,³ Marcory van Dijk,⁴ Julie Sibbring,⁵ and Sarah E. Coupland²

PURPOSE. To evaluate multiplex ligation-dependent probe amplification (MLPA) of uveal melanoma as a predictive tool for metastatic death.

METHODS. Uveal melanoma specimens of 73 patients treated between 1998 and 2000 were included. DNA samples were analyzed with MLPA evaluating 31 loci on chromosomes 1, 3, 6 and 8, and the results were correlated with metastatic death.

RESULTS. The patients (27 women; 46 men) had a median age of 60.6 years and a median follow-up of 6.2 years. Metastatic death occurred in 28 patients, correlating most strongly with chromosome 3 losses and gains on 8q (Cox univariate analysis, $P < 0.001$). Chromosome 6, region p25, gains correlated with good survival (Cox univariate analysis, $P = 0.003$). Prediction of metastatic death was improved by considering equivocal chromosome 3 losses as abnormal and by taking account of multiple risk factors, such as 8q gains, tumor diameter, and histologic features indicative of high-grade malignancy.

CONCLUSIONS. MLPA analysis of uveal melanoma predicts metastatic death if statistically insignificant losses of chromosome 3 are considered together with gains in 8q as well as clinical stage and histologic grade of malignancy. (*Invest Ophthalmol Vis Sci.* 2009;50:3048–3055) DOI:10.1167/iovs.08-3165

Approximately 40% to 50% of uveal melanomas are fatal, usually as a result of metastatic disease, which almost always involves the liver.¹ Estimation of risk of metastatic death would help reassure patients with a good prognosis while enabling any screening for metastasis, systemic adjuvant therapy, and other special measures to be targeted at patients with a poor survival probability.

From the ¹Liverpool Ocular Oncology Centre, St. Paul's Eye Clinic, Royal Liverpool University Hospital, Liverpool, United Kingdom; the ²Department of Pathology, University of Liverpool, Liverpool, United Kingdom; the ³Department of Pathology, University Medical Centre Nijmegen, Nijmegen, The Netherlands; the ⁴Department of Pathology, University Hospital Groningen, Groningen, The Netherlands; and the ⁵Molecular Genetic Department, Liverpool Women's Hospital, Liverpool, United Kingdom.

Supported by the Eye Tumour Research Fund of the Royal Liverpool University Hospital. The Ocular Oncology Service is funded by the National Commissioning Group of the National Health Service, which had no role in the design or conduct of the research.

Submitted for publication November 16, 2008; revised January 10, 2009; accepted May 13, 2009.

Disclosure: B. Damato, None; J. Dopierala, None; A. Klaasen, None; M. van Dijk, None; J. Sibbring, None; S.E. Coupland, None

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.

Corresponding author: Bertil Damato, Ocular Oncology Service, Royal Liverpool University Hospital, Prescot Street, Liverpool L7 8XP, UK; bertil@damato.co.uk.

The main risk factors for metastasis in uveal melanoma include large basal diameter (LBD) of the tumor, ciliary body involvement, extraocular spread, epithelioid cellularity, closed connective tissue loops, high mitotic rate, and certain karyotypic abnormalities, particularly those involving chromosomes 1, 3, 6, and 8.^{2–10}

The most important chromosomal abnormality associated with metastatic death is monosomy 3, which usually involves the loss of an entire chromosome.⁶ Gains in chromosome 8 also indicate a poor prognosis and can consist of trisomy 8, isochromosome 8q, and additions in the *C-MYC* region.⁷ Metastatic mortality also correlates with 1p loss.¹⁰ Conversely, 6p gains are associated with a good prognosis, these abnormalities usually being mutually exclusive with monosomy 3.^{7,9}

Since 1999, we have offered cytogenetic analysis to all patients with uveal melanoma treated by local resection or enucleation, and we have used fluorescence in situ hybridization (FISH) for this purpose. An audit of 356 patients showed a high correlation between monosomy 3 and metastatic death; however, in the absence of monosomy 3, the actuarial 5-year rate of metastatic death ranged from 31% to 6%, depending respectively on whether or not there were gains in chromosome 8.¹¹ We presumed that this was because of partial deletions that were missed by FISH.^{12,13} To address this problem, we developed an artificial neural network that estimated survival prognosis according to age, sex, clinical tumor stage, histologic grade of malignancy, and presence or absence of monosomy 3.¹⁴ To some extent, this program compensated for missed chromosome 3 deletions by estimating the likelihood of these being present, according to the other risk factors present. Although the artificial neural network alleviated the problem caused by the low sensitivity of FISH, we decided to deploy more sensitive methods of genomic typing, which could be applied in routine clinical practice.

We therefore adopted multiplex ligation-dependent probe amplification (MLPA), which comprises a set of 43 probes, each hybridizing to a specific genomic sequence.¹⁵ The probes have different molecular weights and lengths so that after amplification by polymerase chain reaction (PCR), they can be separated by gel electrophoresis and quantified in comparison with control sequences. We used the Salsa P027 (MRC-Holland, Amsterdam, The Netherlands), a kit specifically designed for uveal melanoma. The kit comprises 12 control probes and test probes directed at 7 loci on chromosome 1, 13 on chromosome 3, 6 on chromosome 6, and 5 on chromosome 8.

The purpose of this study was to identify the MLPA results that correlate best with metastatic death from uveal melanoma so that estimation of the prognosis for survival can be enhanced.

METHODS

Patients

DNA from frozen tumor specimens remaining from a previous study on uveal melanoma were retrieved from storage.¹⁶ The patients had been

referred to the Liverpool Ocular Oncology Center between January 1998 and February 2000. They underwent full preoperative ophthalmic and systemic examinations that included slit lamp and ophthalmoscopic examination and echographic measurement of largest and smallest basal tumor dimensions and tumor thickness. Treatment was by enucleation or transscleral local resection. As soon as possible after surgery, a fresh tumor sample was collected for experimental purposes. Part of each tumor was snap frozen in liquid nitrogen, and the remainder was formalin fixed and embedded in paraffin. Histologic features were recorded from hematoxylin and eosin and PAS-stained sections of paraffin-embedded tumors.

Clinical, histologic, and chromosomal data were recorded in a computerized database. Patients residing in mainland Britain were flagged in the National Cancer Registry, which automatically informed us of any deaths, together with the date and cause of death. Follow-up information on overseas patients was obtained whenever possible by writing to the referring ophthalmologist or family doctor. The time until death was calculated from the date of local resection or enucleation.

The research adhered to the tenets of the Declaration of Helsinki. Informed consent was obtained from the patients after explanation of the nature and possible consequences of the study. Ethics committee approval was obtained.

MLPA Technique

DNA extraction, DNA quality assessment and quantification, and MLPA were performed as previously reported but with an improved protocol (Dopierala J, et al., unpublished data, 2008).¹⁵

DNA Extraction

Briefly, the frozen uveal melanoma samples and control specimens consisting of normal choroidal tissue were collected in microfuge tubes. Tissue lysis and protein digestion were performed in a mix of 87% TSE buffer (10 mM Tris/HCl [pH 7.5], 0.4 M NaCl, 2 mM EDTA [pH 8.0]), 0.9% of SDS, and proteinase K (1.64 mg/ μ L; Qiagen GmbH, Hilden, Germany). Samples were incubated overnight at 56°C (Eppendorf Thermomixer Comfort; Sigma-Aldrich, St. Louis, MO). Fresh proteinase K was added after 16 hours of incubation, and samples were incubated for an additional 4 hours. Genomic DNA was isolated by using a high salt concentration (5 M NaCl) and ethanol precipitation. DNA was dissolved in 20 to 100 μ L of TE buffer (10 mM Tris [pH 8.5], 1 mM EDTA [pH 8.0]), depending on pellet size.

DNA Quantification and Quality Assessment

DNA concentration and absorbance at 280 and 260 nm were measured (Nano Drop System; NanoDrop, Minneapolis, MN). FSH-receptor PCR (forward primer CTA CCC TGC ACA AAG ACA GTG; reverse primer GTG TAC GTC ATG TCA AAT CCT CTG C) was performed to assess DNA quality in a thermal cycler (G-Storm GS1; GRI, Ltd., Braintree, UK), with a DNA polymerase kit (GoTaq Flexi; Promega, Madison WI). FSH-receptor PCR products were run on stained 2% agarose gel (SYBR Safe stain; Invitrogen; Groningen, The Netherlands). Electrophoresis was run at 120 V for 30 minutes, and the results were visualized with a transilluminator (Benchtop 2UV; Bio Doc-It Imaging System; UVP, Upland, CA).

MLPA Procedure and Sequencing

An MLPA kit (Salsa P027; MRC-Holland) was used to identify chromosomal imbalances in the uveal melanoma samples. The kit consists of 31 probes selected on chromosomes 1 (short arm), 3, 6, and 8, which often show aberrant copy numbers in uveal melanoma cells, as well as 12 control probes on chromosomes 5, 7, 12, 14, 18, 19, 20, and 21. Two-hundred nanogram aliquots of DNA extracted from uveal melanoma cells as well as four to six normal choroidal tissue control samples were dissolved in 5 μ L of TE buffer and transferred to PCR tubes. TE buffer (5 μ L) was used as a negative control in each run. The

MLPA reaction was performed in the thermal cycler (G-Storm GS1; GRI Ltd.). After the MLPA PCR reaction, 4 μ L of these products were mixed with loading buffer and were run on a stained 2% agarose gel (SYBR Safe stain; Invitrogen) to confirm amplification of the ligation products. After amplification was confirmed, 1 μ L of MLPA products was mixed with 8.8 μ L of highly deionized HI-DI formamide (Applied Biosystems, Inc. [ABI], Foster City, CA) and 0.2 μ L of an internal lane size standard (GeneScan-500 LIZ Size Standard; ABI). Samples were separated by electrophoresis on a 16-capillary genetic analyzer (model 3130XL, analyzed by GeneMapper software; ABI) in the Molecular Genetics Department of the Liverpool Women's Hospital. Data were received as peak heights, as a measure of peak intensity, for each of 43 probes for each sample. Each tumor sample was tested using MLPA at least three independent times.

Data Analysis

Raw data obtained after sequencing were analyzed in a spreadsheet-based approach (Excel; Microsoft, Redmond, WA) designed by the National Genetics Reference Laboratory, Manchester, UK (<http://www.ngrl.org.uk/Manchester/mlpapubs.html>). The results from the uveal melanoma samples were compared with a group of five normal choroid tissue controls, run simultaneously. Data from each test and control sample were normalized by summing the total control probes' peak height and by dividing each ligation product's peak height by this figure. The control and test data were then equalized by dividing the normalized peak height by mean peak height of all five controls. Linear regression was performed to correct for sloping. Regression coefficients were calculated for each of five controls and the test samples, according to the probe lengths. Equalized control and test sample's ligation products were corrected for sloping, and the average results from five control samples for each ligation product were calculated. The Dosage Quotients (DQs) were calculated by dividing the test sample matrix by the control mean matrix. The DQ was categorized as: low, <0.65; equivocally low, 0.65–0.84; normal, 0.85–1.14; equivocally high, 1.15–1.34; and high, >1.34. The standard deviation of mean DQs of control probes for normal control and test samples were calculated. Samples were included in this analysis only if the DQs of the control probes had an SD < 0.2 (higher values indicating poor DNA quality or DNA fragmentation). This amounted to a total of 73 uveal melanomas; 30 samples were excluded.

Correlations between MLPA results and survival were performed (ver.11.0; SPSS Science, Chicago, IL), and Kaplan-Meier estimates were used to draw survival curves for time to metastasis-related death. Associations between metastasis-related death and risk factors were assessed with log rank analysis for categorical variables and with Cox univariate analysis for MLPA DQs.

RESULTS

The 73 patients (46 men and 27 women) had a mean age of 60.6 years (range, 25.2–90.5). The tumors had a mean LBD of 15.0 mm (range, 5.1–21.2), with 31 involving ciliary body (i.e., pars plana and/or pars plicata) and 2 extending extraocularly. Histologic examination showed epithelioid cells in 59 tumors and closed loops in 51 tumors, with the mitotic rate exceeding 5 per 40 high-power fields in 34 tumors. MLPA showed no chromosome 3 abnormalities in 31 patients, 1 abnormal locus in 4 patients, 2 to 9 abnormal loci in 12 patients, and 10 to 13 abnormal loci in 26. The follow-up times had a median of 6.2 years in 35 survivors, exceeding 5 years in 33 patients and 8 years in 31. A total of 38 patients died, the cause of death being metastatic uveal melanoma in 28, other malignancy in 2, and other causes in 8 (cardiac failure, 2; myocardial infarction, 2; cerebrovascular disease, 1; old age, 1; pneumonia, 1; and uncertain, 1). One patient developed hepatic metastasis after treatment of a monosomy 3 melanoma and was alive and well at the close of the study, having undergone a successful partial hepatectomy.

Nine patients had FISH analysis of their tumors, which showed monosomy 3 in four cases and disomy 3 in five. In the four tumors with monosomy 3 on FISH testing, MLPA showed unequivocal loss of 12 or 13 chromosome 3 loci in three cases and equivocal loss of eight loci in one. In the five tumors with disomy 3 on FISH testing, MLPA showed equivocal loss of one chromosome 3 locus in two tumors. Eight tumors had FISH data on chromosome 8. In the three tumors with no chromosome 8 gains on FISH testing, MLPA showed equivocal gain of one locus in one case and unequivocal gains in three loci in another. FISH indicated chromosome 8 gains in five cases, and in these MLPA showed unequivocal gains of all four 8q loci in four tumors and equivocal gains of three adjacent 8q loci in one. MLPA indicated the 8p locus to be normal in seven tumors and lost in one case of monosomy 3.

Table 1 shows the correlations between MLPA results and metastatic death, using Cox univariate analysis. All chromosome 3 loci (i.e., losses) and all chromosome 8q loci (i.e., gains) correlated strongly with metastatic death. Gains in 6p25 showed a strong inverse correlation with metastasis (i.e., strongly indicated a good prognosis). Loss of 8p showed a borderline correlation with worse prognosis. Figure 1 shows boxplots of the MLPA results according to survival.

Table 2 shows the results of Cox multivariate analysis including 3q12, 8q11.23, and 6p25.2 in the model predicting

metastatic death. Only loci in chromosomes 3 and 8 showed significant correlations.

Figure 2 shows Kaplan-Meier survival curves according to whether chromosome 3 loss was considered abnormal if the MLPA results were equivocal (Fig. 2A) or unequivocal (Fig. 2B). Figure 2A indicates that (1) when only unequivocal losses of chromosome 3 loci were considered abnormal, 4 (13%) of 31 patients without such definite abnormalities died of metastatic disease; and (2) high mortality occurred irrespective of the number of abnormal chromosome 3 loci. In Figure 2B, when even equivocal losses of chromosome 3 loci were considered abnormal, only 1 (4.5%) of 22 patients with normal chromosome 3 results died of metastatic disease (this was an unusual tumor, see Fig. 3A). Mortality was higher if all 13 chromosome 3 loci examined using MLPA showed equivocal loss than if fewer loci were abnormal. There were 21 patients with equivocal losses involving only part of chromosome 3, and in these patients the 10-year survival probability was 60%. In Figure 2C, tumors were categorized as having no abnormalities in chromosome 3, only equivocal abnormality and definite abnormality (i.e., unequivocal loss of any locus). Only nine patients showed a moderate survival prognosis (a result that is clinically unhelpful). Figure 2D shows survival according to whether there was loss of any chromosome 3 loci and whether there was gain in 8q11.23. Such abnormalities were considered sig-

TABLE 1. Cox Univariate Analysis Correlating MLPA Results with Metastatic Death

Probe	Locus	B	SE	Wald	Sig.	Exp(B)	95% CI for Exp(B)	
							Lower	Upper
<i>MFN2</i>	1p36.22-	-1.017	0.980	1.076	0.300	0.362	0.053	2.470
<i>NBL1</i>	1p36.13-	-1.379	0.955	2.085	0.149	0.252	0.039	1.637
<i>PTAFR</i>	1p.34-	-0.606	0.934	0.421	0.517	0.546	0.088	3.402
<i>MYCBP</i>	1p.34-	-0.194	0.945	0.042	0.838	0.824	0.129	5.255
<i>MUTYH</i>	1p33-	-0.554	0.910	0.371	0.543	0.575	0.097	3.417
<i>RPE65</i>	1p31-	-1.457	1.081	1.818	0.178	0.233	0.028	1.937
<i>NOTCH2</i>	1p11.2-	-2.093	1.172	3.189	0.074	0.123	0.012	1.227
<i>FANCD2(i)</i>	3p25.3-	-4.042	1.104	13.417	<0.001	0.018	0.002	0.153
<i>FANCD2(ii)</i>	3p24.3-	-2.981	0.877	11.542	0.001	0.051	0.009	0.283
<i>VHL</i>	3p25.3-	-3.093	0.876	12.474	<0.001	0.045	0.008	0.252
<i>MLH1</i>	3p22.1-	-3.093	0.876	12.474	<0.001	0.045	0.008	0.252
<i>CTNNB1</i>	3p22-	-3.163	0.943	11.246	0.001	0.042	0.007	0.269
<i>SEMA3B</i>	3p21.3-	-4.105	1.086	14.283	<0.001	0.016	0.002	0.139
<i>FHIT(i)</i>	3p14.2-	-3.250	0.931	12.198	<0.001	0.039	0.006	0.240
<i>FHIT(ii)</i>	3p14.2-	-3.420	0.998	11.738	0.001	0.033	0.005	0.231
<i>ROBO1</i>	3p12.2-	-3.131	0.868	13.001	<0.001	0.044	0.008	0.240
<i>CPO</i>	3q12-	-4.774	1.175	16.495	<0.001	0.008	0.001	0.085
<i>RHO</i>	3q21.3-	-3.589	0.988	13.186	<0.001	0.028	0.004	0.192
<i>MME</i>	3q25.1-	-4.620	1.210	14.580	<0.001	0.010	0.001	0.106
<i>OPA1</i>	3q29-	-4.594	1.178	15.210	<0.001	0.010	0.001	0.102
<i>FOXC1</i>	6p25+	-1.874	0.692	7.326	0.007	0.153	0.039	0.596
<i>SERPINB9</i>	6p25.2+	-2.097	0.705	8.850	0.003	0.123	0.031	0.489
<i>CDKN1A</i>	6p21.2+	-1.497	0.732	4.175	0.041	0.224	0.053	0.941
<i>RUNX2</i>	6p21+	-1.964	0.868	5.121	0.024	0.140	0.026	0.769
<i>CTGF</i>	6q23.1-	-0.332	0.858	0.150	0.698	0.717	0.133	3.856
<i>IGF2R</i>	6q26-	-0.306	0.894	0.117	0.732	0.736	0.128	4.243
<i>NRG1</i>	8p12-	-1.822	0.883	4.260	0.039	0.162	0.029	0.912
<i>RP1</i>	8q11.23+	1.666	0.304	29.932	<0.001	5.291	2.913	9.609
<i>MYC(i)</i>	8q24.12+	1.415	0.337	17.674	<0.001	4.118	2.129	7.966
<i>MYC(ii)</i>	8q24.12+	1.330	0.328	16.433	<0.001	3.779	1.987	7.188
<i>DDEF1</i>	8q24.2+	1.500	0.328	20.861	<0.001	4.480	2.354	8.527

MFN2, mitofusin2; *NBL1*, neuroblastoma, suppressor of tumorigenicity 1; *PTAFR*, platelet activating factor receptor; *MYCBP*, c-Myc binding protein; *MUTYH*, mutY homolog; *RPE65*, retinal pigment epithelium specific protein 65; *NOTCH2*, notch homolog 2; *FANCD2*, Fanconi anemia complementation group D2; *VHL*, von Hippel Lindau; *MLH1*, MutL homolog 1; *CTNNB1*, catenin, beta-1; *SEMA3B*, semaphoring 3B; *FHIT*, fragile histidine triad gene; *ROBO1*, roundabout, axon guidance receptor, homolog 1; *CPO*, coproporphyrinogen oxidase; *RHO*, rhodopsin; *MME*, membrane metallo-endopeptidase; *OPA1*, optic atrophy 1; *FOXC1*, forkhead box C1; *SERPINB9*, serpin peptidase inhibitor, clade B, member 9; *CDKN1A*, cyclin dependent kinase inhibitor 1A; *RUNX2*, runt related transcription factor; *CTGF*, connective tissue growth factor; *IGF2R*, insulin-like growth factor 2 receptor; *NRG1*, neuregulin 1; *RP1*, retinitis pigmentosa 1; *DDEF1*, *DDEF* intronic transcript 1; B, B statistic; Sig., significance; Exp, exponent; CI, confidence interval.

FIGURE 1. Dosage quotients of each MLPA probe, according to survival, with nonfatal and fatal tumors indicated by *hatched* and *gray* boxes, respectively. *Dotted lines*: separate normal from equivocal abnormality; *dashed lines*: demarcate equivocal from definite abnormality.

nificant, even if equivocal. None of six patients with no abnormalities died of metastasis compared with 80% of patients with abnormality of both chromosomes.

Figure 3A shows the MLPA results of the above-mentioned single patient with no chromosome 3 loss (i.e., equivocal or unequivocal). This patient showed apparent gains in 3p25.3 and 3p22.1 and low normal values of 3p21.3 and 3q21.3 as well as gains in 8q. Three patients showed equivocal loss of one or more chromosome 3 loci (Figs. 3B–D). The patient in Figure 3B showed equivocal loss of all four 3q loci with equivocal gains in 8q. The patient in Figure 3C showed equivocal losses of several 3p loci and 3q29 and gains in 8q. Case 3D showed equivocal losses in 3p22 and 3p14.2 as well as gains in 6p and 8q.

Figure 4 shows survival in patients with lethal melanomas (i.e., tumors with equivocal or definite abnormality of chromosomes 3 and/or 8q). Survival correlated strongly with tumor diameter (log rank, $P = 0.002$; Fig. 4A) and presence of closed loops (log rank, $P = 0.002$; Fig. 4C), weakly with the presence of epithelioid cells (probably because few lethal melanomas were purely of spindle cell type) (log rank, $P = 0.024$) (Fig. 4B), and nonsignificantly with mitotic rate (log rank, $P = 0.14$; Fig. 4D).

DISCUSSION

To our knowledge, this is the first study to correlate metastatic death with MLPA results obtained with the Salsa P027 uveal melanoma kit (MRC-Holland). We found that several fatal tumors showed only equivocal MLPA abnormalities of chromosome 3 loci. Correlation with metastatic disease improved when we considered other risk factors, such as chromosome 8 gains, basal tumor diameter, and histologic features suggestive of high-grade malignancy, such as cell type, and closed loops 1p and 6p abnormalities were relatively uninformative.

We selected MLPA to examine the chromosomal abnormalities in uveal melanomas because it provides more information than FISH, the sensitivity of which has been questioned by us and others.^{11,17} MLPA also requires smaller samples than FISH and can be used on DNA extracted from formalin-fixed, paraffin-embedded tissues.¹⁵ Furthermore, it is relatively inexpensive, so that it is affordable for routine clinical use in a large number of patients. MLPA has been used to investigate uveal melanomas (deploying an aneuploidy kit, Salsa P095; MRC-Holland) and cutaneous melanocytic tumors.^{18,19} We are currently using MLPA as the first-line investigational procedure for chromosomal abnormalities in uveal melanomas recently diag-

TABLE 2. Cox Multivariate Analysis Correlating MLPA Results with Metastatic Death

Locus	B	SE	Wald	Sig.	Exp(B)	95% CI for Exp(B)	
						Lower	Upper
3q12–	–4.085	1.525	7.176	0.007	0.017	0.001	0.334
8q11.23+	1.220	0.339	12.980	<0.001	3.388	1.745	6.581
6p25.2+	0.292	0.833	0.123	0.726	1.339	0.262	6.852

B, B statistic; Sig., significance; Exp, exponent; CI, confidence interval.

FIGURE 2. Kaplan-Meier curves showing rates of metastatic death according to (A) unequivocal (i.e., definite) losses in some or all chromosome 3 loci; (B) equivocal losses in some or all chromosome 3 loci; (C) equivocal and definite losses in chromosome 3; and (D) equivocal or definite abnormality of any chromosome 3 locus and/or 8q11.23.

nosed at the Liverpool Ocular Oncology Center and have analyzed more than 250 tumors with this method. With small biopsy samples, MLPA often succeeded when FISH failed, whereas the converse only rarely occurred (Damato B, et al., unpublished data, 2009).

Despite including 73 samples, the present study is weakened by the relatively small number of patients. In addition, the median follow-up of 6.2 years was probably not long enough to identify all patients destined to die of metastatic disease. Furthermore, several patients died of unrelated causes, possibly masking metastatic deaths. Another limitation is that postmortem examinations were not performed, so that there is uncertainty about the exact cause of death in some patients. Some overseas patients were lost to follow-up and may have died of metastasis. Finally, the survival data may have been influenced, albeit slightly, by apparently successful resection of liver metastasis in one patient.

It was not a purpose of this study to compare the MLPA results with those of FISH, which we started performing only toward the end of the period when these patients were treated. MLPA confirmed all abnormal FISH results. In addition, MLPA revealed equivocal abnormalities in chromosomes 3 and 8 and unequivocal gains in chromosome 8 that were not detected with FISH. We are presently conducting a more in-depth comparison of MLPA with FISH in a large cohort of patients treated since 2006. We are also comparing MLPA with

microsatellite analysis (MSA) and will publish the results in another article. Briefly, MLPA detected equivocal and unequivocal chromosome 3 abnormalities in 7 and 10 of 32 tumors showing no loss of heterozygosity (LOH) with MSA. Conversely, all 41 tumors with LOH on MSA showed abnormalities with MLPA (Damato B, et al., unpublished data, 2008).

The results shown in Figure 2B suggest that equivocal losses in chromosome 3 using MLPA are associated with metastatic death, whether these involve part or all of the chromosome. It must be emphasized that only uveal melanoma samples with good quality MLPA data (i.e., control probes having $SD < 0.2$) were included for interpretation in this study. If only definite chromosome 3 losses were considered abnormal (i.e., a DQ of < 0.65), then more than 10% of patients with a so-called normal MLPA result would have died of metastatic disease. Such a high false-negative rate would be unacceptable in the clinical arena, essentially preventing any patients from receiving reassurance about a good prognosis. If even equivocal chromosome 3 losses were considered abnormal (i.e., a DQ between 0.65 and 0.84), then this false-negative rate would be reduced, to the extent that patients could be confidently reassured of a good prognosis. Categorization of tumors according to the number of chromosome 3 loci with equivocal loss resulted, however, in a high proportion of patients with a 60% 10-year survival probability. Such an uncertain prognosis is not clinically useful because, on the one hand, it does not provide reassurance to

FIGURE 4. Kaplan-Meier survival curves showing rates of metastatic death in patients with lethal uveal melanoma (i.e., equivocal or definite loss of any chromosome 3 loci), according to (A) basal tumor diameter, (B) presence of epithelioid melanoma cells, (C) closed laminin loops, and (D) mitotic rate.

would enable estimation of sensitivity and specificity of MLPA in predicting metastatic death, as well as positive and negative predictive values. It is also necessary to investigate specifically the predictive value of MLPA when performed on tiny tumor specimens, obtained by fine needle aspiration biopsy or with a vitreous cutter. This is because in mosaic tumors, such tumor sampling may miss the most aggressive parts of the tumor.

MLPA and other methods of genomic tumor typing only determine whether the uveal melanoma has any metastatic potential. If the tumor is indeed lethal, then to estimate the likely survival time, it is necessary to take account of (1) clinical tumor stage, which indicates how long any metastases are likely to have been growing (i.e., lead-time bias); and (2) histologic grade, which reflects the rate of growth of such metastases (i.e., tumor doubling time).

In conclusion, MLPA is useful for estimating the probability of metastatic death after treatment of uveal melanoma, but requires multivariate analysis of chromosomes 3 and 8, also taking into account the clinical stage and histologic grade of tumor malignancy.

Acknowledgments

The authors thank Andrew Wallace (National Genetics Reference Laboratory) for the spreadsheet that was used to analyze the MLPA data.

References

- Kujala E, Mäkitie T, Kivelä T. Very long-term prognosis of patients with malignant uveal melanoma. *Invest Ophthalmol Vis Sci.* 2003; 44:4651–4659.
- McLean IW, Foster WD, Zimmerman LE, et al. Modifications of Callender's classification of uveal melanoma at the Armed Forces Institute of Pathology. *Am J Ophthalmol.* 1983;96:502–509.
- Coupland SE, Campbell I, Damato B. Routes of extraocular extension of uveal melanoma: risk factors and influence on survival probability. *Ophthalmology.* 2008;115:1778–1785.
- Folberg R, Pe'er J, Gruman LM, et al. The morphologic characteristics of tumor blood vessels as a marker of tumor progression in primary human uveal melanoma: a matched case-control study. *Hum Pathol.* 1992;23:1298–1305.
- Kivelä T, Mäkitie T, Al-Jamal RT, et al. Microvascular loops and networks in uveal melanoma. *Can J Ophthalmol.* 2004;39:409–421.
- Prescher G, Bornfeld N, Hirsch H, et al. Prognostic implications of monosomy 3 in uveal melanoma. *Lancet.* 1996;347:1222–1225.
- White VA, Chambers JD, Courtright PD, et al. Correlation of cytogenetic abnormalities with the outcome of patients with uveal melanoma. *Cancer.* 1998;83:354–359.
- Aalto Y, Eriksson L, Seregard S, et al. Concomitant loss of chromosome 3 and whole arm losses and gains of chromosome 1, 6, or 8 in metastasizing primary uveal melanoma. *Invest Ophthalmol Vis Sci.* 2001;42:313–317.

9. Parrella P, Sidransky D, Merbs SL. Allelotype of posterior uveal melanoma: implications for a bifurcated tumor progression pathway. *Cancer Res.* 1999;59:3032-3037.
10. Kilic E, Naus NC, van Giles W, et al. Concurrent loss of chromosome arm 1p and chromosome 3 predicts a decreased disease-free survival in uveal melanoma patients. *Invest Ophthalmol Vis Sci.* 2005;46:2253-2257.
11. Damato B, Duke C, Coupland SE, et al. Cytogenetics of uveal melanoma: a seven-year clinical experience. *Ophthalmology.* 2007;114:1925-1931.
12. Tschentscher F, Prescher G, Horsman DE, et al. Partial deletions of the long and short arm of chromosome 3 point to two tumor suppressor genes in uveal melanoma. *Cancer Res.* 2001;61:3439-3442.
13. Parrella P, Fazio VM, Gallo AP, Sidransky D, Merbs SL. Fine mapping of chromosome 3 in uveal melanoma: identification of a minimal region of deletion on chromosomal arm 3p25.1-p25.2. *Cancer Res.* 2003;63:8507-8510.
14. Damato B, Eleuteri A, Fisher AC, et al. Artificial neural networks estimating survival probability after treatment of choroidal melanoma. *Ophthalmology.* 2008;115:1598-1607.
15. van Dijk MC, Rombout PD, Boots-Sprenger SH, et al. Multiplex ligation-dependent probe amplification for the detection of chromosomal gains and losses in formalin-fixed tissue. *Diagn Mol Pathol.* 2005;14:9-16.
16. Scholes AG, Damato BE, Nunn J, et al. Monosomy 3 in uveal melanoma: correlation with clinical and histologic predictors of survival. *Invest Ophthalmol Vis Sci.* 2003;44:1008-1011.
17. Young TA, Burgess BL, Rao NP, et al. High-density genome array is superior to fluorescence in-situ hybridization analysis of monosomy 3 in choroidal melanoma fine needle aspiration biopsy. *Mol Vis.* 2007;13:2328-2333.
18. Mensink HW, Kilic E, Vaarwater J, et al. Molecular cytogenetic analysis of archival uveal melanoma with known clinical outcome. *Cancer Genet Cytogenet.* 2008;181:108-111.
19. Takata M, Suzuki T, Ansai S, et al. Genome profiling of melanocytic tumors using multiplex ligation-dependent probe amplification (MLPA): its usefulness as an adjunctive diagnostic tool for melanocytic tumors. *J Dermatol Sci.* 2005;40:51-57.
20. White VA, McNeil BK, Thiberville L, et al. Acquired homozygosity (isodisomy) of chromosome 3 during clonal evolution of a uveal melanoma: association with morphologic heterogeneity. *Genes Chromosomes Cancer.* 1996;15:138-143.
21. Sandinha T, Farquharson M, McKay I, et al. Correlation of heterogeneity for chromosome 3 copy number with cell type in choroidal melanoma of mixed-cell type. *Invest Ophthalmol Vis Sci.* 2006;47:5177-5180.
22. Maat W, Jordanova ES, van Zelderen-Bhola SL, et al. The heterogeneous distribution of monosomy 3 in uveal melanomas: implications for prognostication based on fine-needle aspiration biopsies. *Arch Pathol Lab Med.* 2007;131:91-96.
23. Ehlers JP, Worley L, Onken MD, et al. Integrative genomic analysis of aneuploidy in uveal melanoma. *Clin Cancer Res.* 2008;14:115-122.