University of Groningen

Angular momentum calculations in coronene mononegative ions

Zandstra, P. J.; van Zorge, B. C.

Published in:
Journal of Chemical Physics

DOI:
[10.1063/1.1680741](https://doi.org/10.1063/1.1680741)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1973

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Zandstra, P. J., & van Zorge, B. C. (1973). Angular momentum calculations in coronene mononegative ions. *Journal of Chemical Physics*, 59(9), 5201-5202. <https://doi.org/10.1063/1.1680741>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Angular momentum calculations in coronene mononegative ions

P. J. Zandstra, and B. C. van Zorge

Citation: [The Journal of Chemical Physics](#) **59**, 5201 (1973); doi: 10.1063/1.1680741

View online: <https://doi.org/10.1063/1.1680741>

View Table of Contents: <http://aip.scitation.org/toc/jcp/59/9>

Published by the [American Institute of Physics](#)

PHYSICS TODAY
WHITEPAPERS

ADVANCED LIGHT CURE ADHESIVES

READ NOW

Take a closer look at what these
environmentally friendly adhesive
systems can do

PRESENTED BY
MASTERBOND
ADHESIVES | SEALANTS | COATINGS

Angular momentum calculations in coronene mononegative ions

P. J. Zandstra and B. C. van Zorge

Department of Physical Chemistry, University of Groningen, Groningen, The Netherlands

(Received 6 July 1973)

The orbital angular momentum of some of the π -electron eigenstates of coronene mononegative and dinegative ions were recently calculated with an HMO method.¹ The m.c.d. measurements and some of the conclusions merit further attention. We have done some configuration interaction (CI) calculations for the mononegative ion to check these former preliminary conclusions which are the following:

(a) The ${}^2E_{1g}$ ground-state orbital angular momentum is well described by the symmetry orbital of the unpaired electron. The effect of CI presumably is very small.

(b) The ${}^2E_{1g} \rightarrow {}^2A_{1,2u}$ transitions are well interpreted by a simple HMO calculation insofar as the angular momentum is concerned. Again, one might be inclined to assume that CI is not important.

(c) The excited ${}^2E_{2u}$ states present a problem in that the agreement between calculated and experimentally determined values is rather poor. The use of an orthogonalized set of orbitals leads to an improvement for some parameter values, for others the discrepancies only increase or stay the same.

Using Mataga parameters for the repulsion and exchange integrals a CI calculation for the ${}^2E_{1g}$ ground state and the ${}^2A_{1,2u}$ and ${}^2E_{2u}$ excited states was performed. The number of configurations used for each symmetry was about 50–70 (with increasing energies), however, as one might expect this number is not too critical, higher energies contributing less in general. In view of earlier results the assumption has been made that the ion has D_{6h} symmetry. The β parameter has been adjusted in such a fashion that the transition energies fall within the proper region ($\beta = -1.8$ eV).

In a subsequent calculation the angular momentum for the configurations of interest was calculated in a straightforward way: For each degenerate state the real configurations, which were + or – with respect to reflection in the xz plane, were written as complex conjugate combinations. The angular momentum can finally be expressed in

the individual MO contributions each with a factor giving its weight in the configuration.¹

The results have been collected in Table I. More details of the calculation are also given in the table. The bond lengths were taken equal 1.41 Å and for the Slater orbital parameter (see Ref. 1) $p = 1.625/0.528$ was chosen.

From a comparison with the experimental results we conclude the following:

(a) The ground state of the mononegative ion is well described by the symmetry orbital. CI for this state is very small. The contribution of this orbital to the groundstate is about 0.96, the angular momentum is $1.716 \hbar$ (as compared to $1.700 \hbar$ computed previously¹).

(b) In the region of the m.c.d. measurements one observes¹ two transitions with negative A/D and positive C/D values. These are due to ${}^2E_{1g} \rightarrow {}^2A_{1u}$, ${}^2A_{2u}$ transitions. For a parameter value $\beta = -1.80$ eV four transitions with reasonably

TABLE I. Experimental and calculated m.c.d. quantities for coronene mononegative ion. The values calculated according to the PPP method are given in parenthesis. See text.

A.							
${}^2E_{1g} \rightarrow {}^2A_{1,2u}$ transitions. $A/D^c = -0.858$, $C/D^c = +0.858$							
Symm	$\beta = -1.80$ (–1.30) eV		experimental (Ref. 1)				
	ν^a	D^b	ν^a	D^b	A/D^c	C/D^c	
${}^2A_{1u}$	6.36(9.22)	1.24					
${}^2A_{2u}$	11.03(14.64)	0.49					
${}^2A_{2u}$	23.97(29.85)	0.69	26.94	0.39	–0.90	+0.60	
${}^2A_{1u}$	26.96(32.68)	1.38	28.48	0.52	–0.50	+0.60	
${}^2A_{1u}$	28.84(37.92)	0.63					
${}^2A_{2u}$	30.49(39.55)	0.35					
${}^2A_{2u}$	38.31	0.09					
${}^2A_{1u}$	38.94	0.01					
B.							
${}^2E_{1g} \rightarrow {}^2E_{2u}$ transitions. $C/D^c = -0.858$							
Symm	$\beta = -1.80$ (–1.30) eV		experimental (Ref. 1)				
	ν^a	D^b	ν^a	D^b	A/D^c	C/D^c	
${}^2E_{2u}$	8.27(10.93)	0.22					
	14.06(20.30)	0.61					
	21.60(27.16)	0.01					
	23.57(32.15)	0.02					
	29.33(36.27)	0.94	+1.110	21.10	0.28	+0.73	–0.42
	31.28(40.30)	1.29	+0.193	32.87	1.39	+0.42	–0.24
a In kK. b In 10^{-16} cm ² . c In (Bohr magneton).							

large intensities of these types have been calculated (see Table I). For $\beta = -2.40$ eV only two remain within this region, the other ones shift strongly towards the uv. We are therefore inclined to correlate the observed ${}^2E_{1g} \rightarrow {}^2A_{1,2u}$ transitions at $\nu = 26.94$ kK and 28.48 kK with the transitions ${}^2E_{1g} \rightarrow {}^2A_{2u}$ at 23.97 kK and ${}^2E_{1g} \rightarrow {}^2A_{1u}$ at 26.96 kK, respectively. A more negative value than $\beta = 1.80$ eV might be more appropriate.

(c) Restricting our discussion to the value of $\beta = -1.80$ eV two strong and two weak ${}^2E_{1g} \rightarrow {}^2E_{2u}$ transitions have been calculated within the experimental region. The strong absorptions are located at 29.33 and 31.28 kK, the weak ones at 21.60 and 23.57 kK (see Table I). It therefore seems very possible to assign the two experimentally observed strong absorption bands with ${}^2E_{2u}$ character (positive A/D and negative C/D) to the bands calculated at 29.33 kK and 31.28 kK. More careful m.c.d. measurements, using a line shape analysis, by Scholtens² have revealed the presence of a weak absorption band at about 30.0 kK of the type ${}^2E_{1g} \rightarrow {}^2E_{2u}$.

Upon taking a more negative value for β the two strong ${}^2E_{1g} \rightarrow {}^2E_{2u}$ transitions move to higher energies, eventually outside the experimental region.

Although one might argue¹ that vibronic interactions are important, it seems that MO theory in its present form does not provide a certain

answer for these transitions.

It may be noticed from Table I that changing β does not lead to a change in order of the ${}^2E_{1g} \rightarrow {}^2E_u$ transitions.

It is striking that the values A/D are positive in agreement with experiment and contrary to our previous HMO results.¹

Finally we would like to mention the fact that SCF CI calculations were made with a PPP approximation. The same magnitudes for the energies can be found using a parameter $\beta = -1.3$ eV. The lower energy levels become more crowded than in the other treatment and we therefore prefer the Mataga method which gives better experimental agreement.

We would like to thank Mr. G. v. d. Velde for many helpful discussions concerning the calculations. The PPP program was provided by Professor E. M. Evleth of the University of Santa Cruz, California. The present investigation has been supported by the Netherlands foundation for Chemical Research (SON) with financial aid from the Netherlands Organization for the Advancement of Pure Research ZWO.

¹P. J. Zandstra, D. J. Scholtens, and R. E. Koning, *J. Chem. Phys.* **57**, 3821 (1972).

²D. J. Scholtens (private communication).