University of Groningen

Comment on "Conductance fluctuations in mesoscopic normal- metal/superconductor samples"

den Hartog, S.G.; van Wees, B.J.

Published in:
Physical Review Letters

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1998

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

den Hartog, S. G., & van Wees, B. J. (1998). Comment on "Conductance fluctuations in mesoscopic normal- metal/superconductor samples". *Physical Review Letters*, 80(22), 5023 - 5023.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

University of Groningen

Comment on “Conductance Fluctuations in Mesoscopic Normal-Metal/Superconductor Samples”

Hartog, S.G. den; van Wees, Bart

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:
1998

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Hartog, S. G. D., & Wees, B. J. V., (1998). Comment on “Conductance Fluctuations in Mesoscopic Normal-Metal/Superconductor Samples”, 1 p.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Comment on “Conductance Fluctuations in Mesoscopic Normal-Metal/Superconductor Samples”

Recently, Hecker *et al.* [1] experimentally studied magnetoconductance fluctuations in a mesoscopic Au wire connected to a superconducting Nb contact. They compared the rms magnitude of these conductance fluctuations in the superconducting state [$\text{rms}(G_{\text{NS}})$] to that in the normal state [$\text{rms}(G_{\text{N}})$] by increasing the magnetic field above the critical field of 2.5 T. It was reported that $\text{rms}(G_{\text{NS}})$ was about 2.8 ± 0.4 times larger than $\text{rms}(G_{\text{N}})$, which should confirm the theoretical predicted enhancement factor of $2\sqrt{2} \approx 2.8$.

In this Comment, we show that their claim is not justified. Although not explicitly mentioned in Ref. [1], we have to assume that the $\text{rms}(G)$ was calculated according to $\text{rms}(G) = \text{rms}(R)/R^2$, where $\text{rms}(R)$ denotes the rms magnitude of the measured resistance fluctuations and R the total measured resistance. The point we want to make is that the authors did not take into account the presence of an incoherent series resistance R_{series} from the contacts, which is different when the Nb is in the superconducting or normal state. Since the measured $\text{rms}(R)$ originates only from the phase-coherent part of the disordered conductor, with resistance R_{φ} , the correct procedure is to calculate $\text{rms}(G)$ according to $\text{rms}(G) = \text{rms}(R)/R_{\varphi}^2 = \text{rms}(R)/(R - R_{\text{series}})^2$. As shown below, when we correct for the presence of this series resistance, we find that $\text{rms}(G_{\text{NS}})$ is *not* significantly larger than $\text{rms}(G_{\text{N}})$.

Their device consists of a narrow Au wire (Au^w , length $L = 1.0 \mu\text{m}$, width $W = 0.13 \mu\text{m}$) connected at its ends to a macroscopic Nb and Au contact (Nb^c or Au^c) via a rectangular shaped contact (Nb^r or Au^r , $L = 0.8 \mu\text{m}$, $W = 1.6 \mu\text{m}$). The total resistance is the sum of these five contributions: $R = R_{\text{Nb}}^c + R_{\text{Nb}}^r + R_{\text{Au}}^w + R_{\text{Au}}^r + R_{\text{Au}}^c$, where $R_{\text{Nb}}^c + R_{\text{Nb}}^r$ are zero in the superconducting state.

Since the series resistances of the Au contact ($R_{\text{Au}}^c + R_{\text{Au}}^r \approx 1.2R_{\square}^{\text{Au}} \approx 1.1 \Omega$) are small compared to phase-coherent resistance of the Au wire (10.5Ω), we will correct only for the series resistances of the Nb contact ($R_{\text{Nb}}^c + R_{\text{Nb}}^r \approx 1.2R_{\square}^{\text{Nb}} \approx 4.8 \Omega$). This series resistance is present only in the normal state and is exactly equal to the increase in resistance when the magnetic field exceeds B_c (see Fig. 1(a), in Ref. [1]). We note that not only the macroscopic Nb contact is regarded to be incoherent but the rectangular shaped Nb contact as well. Namely, the phase-breaking length $L_{\varphi} \equiv \sqrt{D\tau_{\varphi}}$ for Nb is expected to be reduced compared to $L_{\varphi} \approx 0.6 \mu\text{m}$

TABLE I. The measured resistance R_{NS} and uncorrected conductance fluctuations $\text{rms}(G_{\text{NS}})$ in the superconducting state at $T = 50 \text{ mK}$ and $B = 1 \text{ T}$, and the measured resistance R_{N} and the *corrected* conductance fluctuations $\text{rms}(G_{\text{N}})$ in the normal state at $T = 50 \text{ mK}$ and $B = 4 \text{ T}$.

	Sample 1	Sample 2
$R_{\text{NS}} (\Omega)$	11.60	9.72
$R_{\text{N}} (\Omega)$	15.87	14.34
$\text{rms}(G_{\text{NS}}) (e^2/h)$	0.16 ± 0.02	0.14 ± 0.02
$\text{rms}(G_{\text{N}}) (e^2/h)$	0.109 ± 0.006	0.109 ± 0.009
$\text{rms}(G_{\text{NS}})/\text{rms}(G_{\text{N}})$	1.5 ± 0.2	1.3 ± 0.2

for Au by $\sqrt{D_{\text{Au}}/D_{\text{Nb}}} \approx 2.5$, which implies that the resistance fluctuations from this Nb rectangle are strongly suppressed due to ensemble averaging as well.

In Table I we have reproduced the measured (average) resistance of the two studied samples in the normal state and in the superconducting state. We did not correct $\text{rms}(G_{\text{NS}})$ [2]. The $\text{rms}(G_{\text{N}})$ has been corrected as described above. As a result, the $\text{rms}(G_{\text{N}})$ are a factor of $(R_{\text{N}}/R_{\text{NS}})^2 \approx 2$ larger than reported in Ref. [1] and, consequently, the ratio $\text{rms}(G_{\text{NS}})/\text{rms}(G_{\text{N}})$ becomes about 1.4 ± 0.2 . We doubt, however, that the remaining difference from 1 is significant, since the statistical error could well be larger than 0.2 due to the fact that only a few large fluctuations determine $\text{rms}(G_{\text{NS}})$ (see Figs. (1b) and 2, in Ref. [1]).

In conclusion, we have argued that the measured $\text{rms}(G_{\text{NS}})$ is not significantly enhanced compared to $\text{rms}(G_{\text{N}})$, and it remains an experimental challenge to observe the predicted enhancement factor of $2\sqrt{2}$.

S. G. den Hartog and B. J. van Wees

Department of Applied Physics and Materials Science
Centre, University of Groningen
Nijenborgh 4, 9747 AG Groningen, The Netherlands

Received 1 October 1997

[S0031-9007(98)06217-6]

PACS numbers: 73.23.-b, 73.50.Jt, 74.80.-g

- [1] K. Hecker, H. Hegger, A. Altland, and K. Fiegle, *Phys. Rev. Lett.* **79**, 1547 (1997).
- [2] The reported values for $\text{rms}(G_{\text{NS}})$ are considerably smaller than the rms magnitude of the sample-specific conductance fluctuations of about $\text{rms}(G_{\text{NS}}) \approx 1.0e^2/h$ observed in both a cross-shaped and a T-shaped two-dimensional electron gas coupled to superconductors. S. G. den Hartog *et al.*, *Phys. Rev. Lett.* **77**, 4954 (1996); S. G. den Hartog *et al.*, *ibid.* **76**, 4592 (1996). A comparison with the normal state values was not made in these experiments.