University of Groningen

Germinal Centers in Diagnostic Biopsies of Patients With Primary Sjogren's Syndrome Are Not a Risk Factor for Non-Hodgkin's Lymphoma but a Reflection of High Disease Activity

Haacke, Erlyn A.; van der Vegt, Bert; Vissink, Arjan; Spijkervet, Fred K. L.; Bootsma, Hendrika; Kroese, Frans G. M.

Published in:
Arthritis & Rheumatology

DOI:
[10.1002/art.40715](https://doi.org/10.1002/art.40715)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Final author's version (accepted by publisher, after peer review)

Publication date:
2019

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Haacke, E. A., van der Vegt, B., Vissink, A., Spijkervet, F. K. L., Bootsma, H., & Kroese, F. G. M. (2019). Germinal Centers in Diagnostic Biopsies of Patients With Primary Sjogren's Syndrome Are Not a Risk Factor for Non-Hodgkin's Lymphoma but a Reflection of High Disease Activity: Comment on the Article by Sene et al. *Arthritis & Rheumatology*, 71(1), 170-171. <https://doi.org/10.1002/art.40715>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Germinal centres in diagnostic biopsies of pSS patients are not a risk factor for non-Hodgkin's lymphoma but a reflection of high disease activity.

1) Erlin A Haacke, *MD*. Dept. of Rheumatology and Clinical Immunology AND ²Dept. of Pathology and Medical Biology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.

2) Bert van der Vegt, *MD, PhD*. Dept. of Pathology and Medical Biology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.

3) Arjan Vissink, *DMD, MD, PhD*. Dept. of Oral and Maxillofacial Surgery, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.

4) Fred K. L. Spijkervet, *DMD, PhD*. Dept. of Oral and Maxillofacial Surgery, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.

5) Hendrika Bootsma, *MD, PhD*. Dept. of Rheumatology and Clinical Immunology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.

6) Frans G.M. Kroese, *PhD*. Dept. of Rheumatology and Clinical Immunology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.

Key words: Sjögren's syndrome, germinal centres, salivary gland biopsies, Non-Hodgkin's lymphoma, MALT lymphoma

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/art.40715

This article is protected by copyright. All rights reserved.

Correspondence to:

Frans G.M. Kroese.

Department of Rheumatology & Clinical Immunology, University of Groningen, University Medical

Center Groningen, HPC AA21, Hanzeplein 1, 9713 GZ, Groningen, The Netherlands

Email: f.g.m.kroese@umcg.nl

Dear Editor,

With great interest we have read the article of Sène et al. (1) in which ectopic germinal centres (GCs) in labial gland biopsies of primary Sjögren syndrome (pSS) patients were found predictive for Non-Hodgkin's lymphoma (NHL) development later in the disease. In the univariate analysis the presence of GCs in these biopsies was not significantly different between pSS patients that developed NHL or pSS patients that did not. However, multivariate analysis revealed that presence of GCs in biopsies was an independent predictor for NHL development (1). This study adds to the ongoing discussion about the presence of ectopic GCs in pSS diagnostic salivary gland biopsies as a risk factor for subsequent NHL development. The study of Theander et al. (2) showed that GCs in diagnostic labial gland biopsies was predictive for NHL development in pSS patients, whereas we (3) and others (4,5) did not detect such an association. As discussed extensively, (3,6) a major reason for the apparent discrepancy in the different studies was the variation in NHL-subtypes that were included.

Remarkably, half of the pSS patients in the study of Sène et al. (1) were male, whereas in the other studies that evaluated the presence of GCs as risk factor, the majority of patients (>81%) were females (2–5). Whether the predictive value of presence of GCs in diagnostic biopsies differs between males and females is not known.

A very unusual observation in the study of Sène et al. (1) was that **all** pSS patients who developed NHL, had a monoclonal gammopathy (MG). In pSS patients in general, the presence of MG is 4-22% (1,7,8). We observed MG at time of lymphoma diagnosis in 4/8 (50%) pSS patients with a

parotid MALT lymphoma (Haacke et al., unpublished data). In another recent study only 3/7 (43%) pSS patients with pulmonary MALT lymphomas exhibited MG (9). Presence of MG is known as a risk factor for NHL development, but is also associated with higher disease activity (8,10). The presence of MG in pSS pre-lymphoma patients in the study of Sène et al. (1), could thus also be a reflection of high disease activity (ESSDAI) which is an independent predictor for NHL development (11).

Also the presence of GCs in biopsies is associated with high disease status (12). Since disease activity as measured by ESSDAI can change over time, (13) the time point when the diagnostic pSS biopsy is taken, is of crucial importance. If a labial gland biopsy is taken during a period of relatively low disease activity, the likelihood of presence of GCs might consequently be low. On the other hand, when a diagnostic pSS salivary gland biopsy is taken in a period of high disease activity, the chance of finding GCs is higher.

Thus, both GCs and MG are associated with higher disease status, but there is no clear indication that presence of *ectopic* GCs is a prerequisite for MALT lymphoma development. There are many pSS patients with GCs in their salivary gland biopsies that do not develop NHL. Of note, presence of GCs is usually assessed in labial salivary glands, which are not the sites where MALT lymphoma preferentially develop. We conclude that for prediction of pSS patients who are at risk for NHL development, clinical and laboratory factors such as low C4, RF-positivity, presence of cryoglobulin, highly active disease, purpura, lymphadenopathy and especially persistent parotid enlargement (5,11,14) are more important than presence of GCs in diagnostic salivary gland biopsies.

References

1. Sène D, Ismael S, Forien M, Charlotte F, Kaci R, Cacoub P, et al. Ectopic germinal centre-like structures in minor salivary gland biopsy predict lymphoma occurrence in patients with primary Sjögren syndrome. *Arthritis Rheumatol* 2018. [Epub ahead of print]
2. Theander E, Vasaitis L, Baecklund E, Nordmark G, Warfvinge G, Liedholm R, et al. Lymphoid organisation in labial salivary gland biopsies is a possible predictor for the development of malignant lymphoma in primary Sjögren's syndrome. *Ann Rheum Dis* 2011;161:1363–1368.
3. Haacke EA, Vegt B van der, Vissink A, Spijkervet FKL, Bootsma H, Kroese FGM. Germinal centres in diagnostic labial gland biopsies of patients with primary Sjögren's syndrome are not predictive for parotid MALT lymphoma development. *Ann Rheum Dis* 2017;76:1781–1784.
4. Johnsen SJ, Gudlaugsson E, Skaland I, Janssen E, Jonsson M V, Helgeland L, et al. Low Protein A20 in Minor Salivary Glands is Associated with Lymphoma in Primary Sjögren's Syndrome. *Scand J Immunol* 2016;83:181–187.
5. Fragkioudaki S, Mavragani CP, Moutsopoulos HM. Predicting the risk for lymphoma development in Sjogren syndrome: An easy tool for clinical use. *Medicine (Baltimore)* 2016;95:e3766.
6. Kroese FGM, Haacke EA, Bombardieri M. The role of salivary gland histopathology in primary Sjögren's syndrome: promises and pitfalls. *Clin Exp Rheumatol* 2018 [In press].

- Accepted Article
7. Baimpa E, Dahabreh IJ, Voulgarelis M, Moutsopoulos HM. Hematologic manifestations and predictors of lymphoma development in primary Sjögren syndrome: clinical and pathophysiologic aspects. *Medicine (Baltimore)* 2009;88:284–93.
 8. Brito-Zerón P, Kostov B, Fraile G, Caravia-Durán D, Maure B, Rascón F-J, et al. Characterization and risk estimate of cancer in patients with primary Sjögren syndrome. *J Hematol Oncol* 2017;10:90.
 9. Yachoui R, Leon C, Sitwala K, Kreidy M. Pulmonary MALT Lymphoma in Patients with Sjögren's Syndrome. *Clin Med Res* 2017;15:6–12.
 10. Yang Y, Chen L, Jia Y, Liu Y, Wen L, Liang Y, et al. Monoclonal gammopathy in rheumatic diseases. *Clin Rheumatol*. 2018;37:1751-1762.
 11. Nocturne G, Virone A, Ng W-F, Guern V Le, Hachulla E, Cornec D, et al. Rheumatoid Factor and Disease Activity Are Independent Predictors of Lymphoma in Primary Sjögren's Syndrome. *Arthritis Rheumatol (Hoboken, NJ)* 2016;68:977–85.
 12. Risselada AP, Looije MF, Kruize AA, Bijlsma JWW, Roon JAG van. The role of ectopic germinal centers in the immunopathology of primary Sjögren's syndrome: a systematic review. *Semin Arthritis Rheum* 2013;42:368–76.
 13. Gottenberg J-E, Seror R, Saraux A, Devauchelle V, Labous E, Dieudé P, et al. Evolution of Disease Activity over a 5-Year Period in the 395 Patients with Primary Sjögren's Syndrome of the Assess Prospective Cohort [Abstract]. *Arthritis Rheumatol* 2016;68.
 14. Nishishinya MB, Pereda CA, Muñoz-Fernández S, Pego-Reigosa JM, Rúa-Figueroa I, Andreu J-L, et al. Identification of lymphoma predictors in patients with primary Sjögren's syndrome: a systematic literature review and meta-analysis. *Rheumatol Int* 2015;35:17–26.