University of Groningen

Social participation of adult patients with spinal muscular atrophy

Kruitwagen-van Reenen, Esther Th; van der Pol, Ludo; Schröder, Carin; Wadman, Renske I; van den Berg, Leonard H; Visser-Meily, Johanna M A; Post, Marcel W M

Published in:
MUSCLE & NERVE

DOI:
[10.1002/mus.26201](https://doi.org/10.1002/mus.26201)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Final author's version (accepted by publisher, after peer review)

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Kruitwagen-van Reenen, E. T., van der Pol, L., Schröder, C., Wadman, R. I., van den Berg, L. H., Visser-Meily, J. M. A., & Post, M. W. M. (2018). Social participation of adult patients with spinal muscular atrophy: Frequency, restrictions, satisfaction, and correlates. *MUSCLE & NERVE*, 58(6), 805-811.
<https://doi.org/10.1002/mus.26201>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Social participation of adult patients with spinal muscular atrophy: frequency, restrictions, satisfaction and correlates.

Esther Th. Kruitwagen-van Reenen, MD (1,2); Ludo van der Pol, MD, PhD (4); Carin Schröder, PhD (1,2); Renske I. Wadman, MD (4); Leonard H. van den Berg, MD, PhD (4); Johanna M.A. Visser-Meily, MD, PhD (1,2); Marcel W.M. Post, PhD (2,3)

1. Department of Rehabilitation, Physiotherapy and Sports, Brain Center Rudolf Magnus, University Medical Center Utrecht, Postbus 85500, 3508 GA Utrecht. The Netherlands

2. Center of Excellence for Rehabilitation Medicine, Brain Center Rudolf Magnus, University Medical Center Utrecht, Utrecht University and De Hoogstraat Rehabilitation, Utrecht, The Netherlands

3. Department of Rehabilitation Medicine, University of Groningen, University Medical Center Groningen, The Netherlands

4. Department of Neurology and Neurosurgery, Brain Center Rudolf Magnus, University Medical Center Utrecht, Postbus 85500, 3508 GA Utrecht, The Netherlands

Corresponding author/guarantor and address:

Dr. E. Th. Kruitwagen - van Reenen

Department of Rehabilitation, Physiotherapy and Sports

University Medical Center Utrecht

The Netherlands

Tel.: +31 (0)88 7558831

E-Mail: E.T.Kruitwagen@umcutrecht.nl

Running title: Social participation of adult SMA

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/mus.26201

Financial Disclosure statement:

E. Th. Kruitwagen-van Reenen reports no disclosures.

W.L. van der Pol receives research support from the Prinses Beatrix Spierfonds, Netherlands ALS Foundation and Stichting Spieren voor Spieren; he serves as a member of scientific advisory boards of Biogen and Avexis and of a data monitoring committee of Novartis for which his employer receives financial compensation.

C. Schröder reports no disclosures

R.I. Wadman reports no disclosures.

J.M.A. Visser-Meily reports no disclosures.

L.H. van den Berg serves on scientific advisory boards for ARISLA the Thierry Latran Foundation, Biogen, Cytokinetics and Orion; serves on the editorial board of Amyotrophic Lateral Sclerosis, The Journal of Neurology, Neurosurgery and Psychiatry; and receives research support from the Prinses Beatrix Fonds, Netherlands ALS Foundation, and the Netherlands Organization for Scientific Research VICI Grant.

M.W.M. Post reports no disclosures

Acknowledgments:

The authors wish to thank all participating patients.

Ethical publication statement: “We confirm that we have read the Journal’s position on issues involved in ethical publication and affirm that this report is consistent with those guidelines.”

ABSTRACT

Social participation of adult patients with spinal muscular atrophy: frequencies, restrictions, satisfaction and correlating factors.

Introduction We assessed social participation in 62 adults with spinal muscular atrophy (SMA) types 1c-4.

Methods Outcome measure: Utrecht Scale of Evaluation Rehabilitation-Participation (USER-P) with Frequency, Restrictions and Satisfaction scores. Hierarchical regression analysis.

Results Early (type 1,2,3a) and late onset (type 3b,4) SMA patients reported similar frequency and satisfaction scores. ‘Age’, ‘motor skills’, ‘pain’ and ‘feelings of depression’ correlated with frequency; ‘motor skills’ and ‘feelings of depression’ correlated with restrictions and ‘level of education’, ‘fatigue’ and ‘feelings of depression’ correlated with satisfaction. Motor skills and feelings of depression explained 33% of variance in *frequency* of participation. Motor skills explained 26% of variance of *restrictions* in participation. Fatigue and feelings of depression explained 50% of variance in *satisfaction* with participation.

Discussion Motor skills, feelings of depression and fatigue are correlates of participation in daily life. This knowledge can be used to optimize care for SMA patients.

Key words: social participation, satisfaction, social engagement, activities of daily living, SMA, adults

INTRODUCTION

Hereditary proximal spinal muscular atrophy (SMA) is an important genetic cause of disability in childhood and adult life.¹ The antisense oligonucleotide nusinersen has been approved and has shown efficacy in altering the natural history of the disease.² Other gene therapy trials are ongoing. However, there is currently no cure for SMA.^{3,4,5}

The variety and severity of impairments and disabilities that accompany SMA have been described extensively.⁶⁻¹⁰ The consensus guidelines for supportive care outlines management for the most common medical complications (e.g. pulmonary problems, scoliosis) that occur primarily in childhood, but focus strongly on symptom management.¹¹ Irrespective of SMA type, all patients will encounter moderate to severe disability in life. There is, however, a striking lack of literature on how adult patients participate in daily life activities and how multidisciplinary care can optimize their participation. The few existing studies on participation among adult patients with a broad range of neuromuscular disorders have focused mainly on work, and the results of SMA patients were not reported separately.¹²⁻¹⁵ In another qualitative study, SMA patients rated their quality of life as 'fine', but experienced the serious impact of progressive functional limitations on their daily activities.¹⁶ Patients emphasized their need to live a normal life and fully participate in social activities, including an active family role, work and maintaining optimism.¹⁷ Increasing our knowledge of participation among adult patients with SMA and factors associated with participation may help to optimize supportive care.

Therefore, the first objective of this study was to describe the frequency of, and perceived restrictions in, participation and the related satisfaction levels of adult patients with SMA. The second objective was to determine whether selected subjective complaints (pain, fatigue, anxiety, feelings of depression) and coping style are associated with participation in adult patients with SMA, adjusting for demographic factors and disease severity. These possible

correlates were selected based on studies showing associations between pain and fatigue, emotional distress, mood and coping strategies with participation and quality of life among patients with neuromuscular diseases.¹⁸⁻²¹

METHODS

Subjects and Procedures

Adult patients with genetically confirmed SMA, regardless of type of SMA, aged 18 years or older at inclusion, were recruited for this study between September 2010 and December 2012. Patients were informed about the study and recruited through the Dutch patient organization for neuromuscular diseases (www.spierziekten.nl), through patient communities on the internet, through pediatricians, (pediatric) neurologists, rehabilitation physicians and the four Dutch Centers for Chronic Respiratory Ventilation. The only exclusion criterion was the inability to read Dutch. All patients were retrieved from the Dutch SMA database.²² For the purpose of this study, all patients were seen at the outpatient clinic of the Department of Neurology and Neurosurgery at the University Medical Center Utrecht for a structured interview and neurological examination. They were asked to complete questionnaires. This study was registered in the Dutch registry for clinical trials (study no. NL29692.041.09/29692). The Medical Ethics Committee of the University Medical Center Utrecht approved the research protocol. All patients gave informed consent prior to inclusion.

Measures

The SMA classification system was used to define SMA types 1-4^{6,23,24}

Participation was measured using the Utrecht Scale for Evaluation Rehabilitation-

Participation (USER-P), a self-report instrument with 32 items. Validity and reproducibility of the USER-P scale are good.²⁵⁻²⁹ Sum scores are calculated for the Frequency, Restrictions and

Satisfaction scales, and each sum score is converted to a score on a scale ranging from 0 to 100. Higher scores indicate more favourable levels of participation (higher frequency of activities, fewer restrictions, greater satisfaction). An example of a question for the Frequency scale is: “How many hours do you spend on household activities?”

An example of a question for the Restrictions scale is: “Does your illness or condition currently limit your daily life concerning outdoor mobility?”. An example of a question for the Satisfaction scale is: “How satisfied are you with your current daily life concerning ‘going out’ (eating out, visiting a cafe, the cinema, a concert, alone or with others).”

Correlates

We used a standardized questionnaire to document disease characteristics (age at diagnosis, age at loss of ambulation) and demographic characteristics (sex, age, relationship status, job status, level of education, whether living independently).

To document motor skills we used the Expanded Hammersmith Functional Motor Scale (HFMSE), a validated test consisting of 33 items to assess motor skills of patients with SMA.^{30,31} The maximum score is 66 points. Higher scores indicate better motor skills.

To assess pain and fatigue we used the sub-domain scores ‘pain’ (two items) and ‘vitality’ (four items) of the Short Form 36-item Health Survey (SF-36).^{32,33} The vitality score is a valid instrument to measure fatigue, as shown by a high correlation between the SF-36 vitality score and scores on the Fatigue Symptom Inventory.³⁴ Higher scores in the domain ‘pain’ indicate that patients experience less pain; higher scores in the domain ‘vitality’ indicate that patients experience less fatigue and feel more vital.

Feelings of depression and anxiety were assessed using the Hospital Anxiety and Depression Scale (HADS).³⁵⁻³⁷ Seven items assess feelings of anxiety (HADS-A) and seven items assess feelings of depression (HADS-D), with a maximum of 21 points for the HADS-A and HADS-

D scales. A score of 8 or more on the HADS-A or HADS-D is an indication of anxiety or depressed mood.

Coping strategies were assessed using the short version of the Coping Inventory for Stressful Situations (CISS-21).^{38,39} The CISS-21 has 21 items divided into three categories: problem-targeted strategies (CISS-P), emotion-focused strategies (CISS-E) and avoidance strategies (CISS-A). Response options range from ‘not at all’(1) to ‘very applicable’(5). A higher score indicates a preference to use this particular coping strategy.

Statistical analyses

Descriptive statistics were used to describe characteristics of the study population and participation scores. Comparisons between all SMA types were not performed because of the small sample size. Since differences in quality of life were found between patients with a relatively early onset (i.e. SMA types 1-3a) and those with onset later in life (i.e. SMA types 3b-4),⁴⁰ we dichotomized the variable SMA types to early onset SMA (types 1-3a) versus late onset SMA (types 3-4) to compare outcomes between these two subgroups.

In addition to the total scores on the USER-P, individual items of the Restrictions and Satisfaction subscales were dichotomized to quantify the presence of restrictions in and dissatisfaction with specific aspects of participation. The options “with difficulty”, “with assistance” and “not possible” on the restrictions subscale were defined as “restrictions”. The option “without difficulty” was defined as “no restrictions”. The answer options “satisfied” and “very satisfied” on the satisfaction subscale were defined as “satisfaction”. The answer options “very dissatisfied”, “dissatisfied” and “neutral” were defined as “dissatisfaction”. The answer option ‘not applicable’ was defined as a missing variable.

To detect differences in determinants of participation between early and late-onset SMA patients, we used the independent samples Mann Whitney U test. Spearman correlations were

computed to determine relationships between potential determinants and participation of the total sample. Using Cohen's rule of thumb, a correlation of 0.10 was considered 'small', of 0.30 'medium' and of 0.50 'large'.⁴¹

Determinants that showed a p-value < 0.1 in the bivariate correlation analysis were entered into a hierarchical linear regression model (Multi-step enter method). Variables were always entered in the same order: step 1: disease severity variables and demographics; step 2: subjective complaints. Residual analyses were performed and multi-collinearity was tested to search for violations of necessary assumptions in multiple regression.⁴² SPSS version 24 for Windows was used for analysis.

RESULTS

Sixty-two of 80 (78%) invited patients participated. Descriptive statistics are displayed in Table 1. We included four patients with SMA type 1c and onset before 6 months of age who never learned to sit independently but survived into adulthood. Median HADS-A and HADS-D scores were low (Table 2). Patients with early onset SMA had significantly lower motor skills (HFMSE-scores) than those with late onset SMA. Significantly fewer patients with early onset SMA had a partner than patients with late onset SMA. Late onset patients reported more pain and more fatigue than early onset patients.

Frequency (level) of participation: Median participation scores are displayed in Table 3.

There were no significant differences between patients with early and late onset SMA in the frequency of participation (Table 3). SMA patients spent most hours on unpaid work and household activities (Table 4). Only 39% of them had paid work and only 16% had a full-time job (36 hours a week or more). A detailed description of all participation activities performed by all patients is given in Table 4.

Restrictions and satisfaction: Patients with early onset SMA experienced significantly more participation restrictions compared to patients with late onset SMA (Table 3). Table 5 shows that 16-97 % of patients felt restricted in daily activities. They felt most restricted in work/education, household chores, mobility outdoors (e.g. traveling by car, public transport or bike), going out (e.g. to a movie, pub, church, or shopping), physical exercise and visits to family and friends. There were no significant differences between patients with early and late onset SMA in satisfaction with participation (Table 3). Table 5 shows that 8-58 % of patients were dissatisfied about daily activities. They were most dissatisfied about performing household chores and physical exercise.

Correlates of participation (Table 6). Bi-variable analysis showed a medium correlation between lower frequency of participation and older age, reduced motor skills and pain. We found a medium correlation between lower frequency of participation and more feelings of depression. Participation restrictions were largely correlated with reduced motor skills. There was a medium correlation between satisfaction with participation and level of education, , fatigue and a large correlation with fewer feelings of depression.

Table 6 summarizes results of the multivariable analysis. 33% of the variance in frequency of participation was explained by motor skills in combination with feelings of feelings of depression; 26% of the variance in restrictions in participation was explained by motor skills and 50 % of the variance in satisfaction with participation was explained by fatigue together with feelings of feelings of depression.

DISCUSSION

Patients with early onset SMA (type 1,2,3a) experienced more participation restrictions than patients with later onset SMA, but reported similar levels of frequency of participation and resulting satisfaction. Motor skills were independently associated with the frequency of and

restrictions in participation. In addition, subjective complaints, namely fatigue, pain and particularly feelings of depression, were associated with participation scores in bivariate analyses. Coping styles were not associated with participation scores. In the multiple regression analyses, having feelings of depression proved to be the only subjective complaint independently related to participation (frequency and satisfaction).

This study identifies the participation activities that pose most problems for adult patients with SMA. There are a few studies on specific domains of participation that enrolled patients with a broader range of neuromuscular diseases. Employment rates of 40-57 % were, for example, found in a mixed sample of patients with neuromuscular disorders.^{12,13,14} This percentage is not only similar to our findings, but also comparable to that of patients with spinal cord injury.²⁸ Our data, therefore, seem to be in line with previous reports on participation in work activities of patients with moderate to severe motor impairments. Satisfaction scores of patients with SMA were also similar to those previously reported by patients with spinal cord injury and patients after stroke.^{28,29} It reflects the fact that patients in different situations are able to reorganize their social activities on average, in a satisfactory way, regardless of their restrictions and type of condition.

Age at onset of SMA may also be relevant for satisfaction with participation, since patients with SMA type 1c-3a experienced more restrictions but similar levels of satisfaction as patients with a later onset. A possible explanation for this finding is that patients with early onset SMA might have adapted more successfully to living with severe physical limitations from a very early age onwards.⁴³ Patients with late onset SMA initially have normal motor development. Relevant physical limitations due to disease progression occur in adulthood. Continuous adaptation attempts might lead to enduring distress as these patients have to redefine their goals and concepts about their daily functioning.

Although the incidence of feelings of depression among patients with SMA is not higher than in those with other neuromuscular diseases,^{18,20,21} the presence of feelings of depression was inversely associated with both frequency of participation activities and satisfaction with participation. This suggests that it is important to monitor whether feelings of depression are present in adult patients with SMA, and if so, to consider psychological interventions in order to reduce these feelings that might be the result of impaired emotional adaptation to disease progression.^{43,44}

Coping style was not associated with any aspect of participation. To the best of our knowledge, there are no studies on the relationship between coping style and social participation in patients with neuromuscular disorders. Studies have, however, been carried out on MS patients and spinal cord injury patients, and these also failed to find a relationship between coping styles and participation.^{45,46} The fact that healthy individuals exhibit higher levels of coping variability than patients with chronic disease may play a role in this.⁴⁷ We cannot, however, exclude the possibility that coping styles are relevant for participation (satisfaction) in a subgroup of patients, in particular among patients with SMA type 3-4 who may face challenges of adaptation in later life.

As in other studies, pain was reported frequently, in particular by late onset patients.^{18,48}

Causes of pain are multiple, and include spinal deformities, muscle cramps or neurogenic pain. Univariable analysis showed that pain is associated with the frequency of participation activities in patients with SMA, but when feelings of depression was entered into the model, pain did not add significantly to the model (see table 6).

The limited sample size is an important limitation of this study. It allowed the (pre)selection of a limited number of possible correlates. We made this preselection based on the existing literature and clinical experience. In our study we included adult patients with SMA over the whole spectrum (type 1c-4) and assessed participation in more detail than before. This gave

us the opportunity to assess the relevance of already known variables in a broader group of patients with SMA, thus expanding the clinical relevance of our study. Follow-up studies should aim at a larger sample size to address the importance of other factors including endurance and stamina for motor activities, social support, upper extremity function and personal factors such as self-efficacy or illness perceptions, in particular since the largest portion of the variance in participation remains unexplained. Larger sample sizes would also allow more detailed subgroup analysis, for example, of subgroups with early and later onset. Although almost 80% of invited patients participated in this population-based study, we cannot fully exclude the possibility of inclusion bias, i.e., the selection of patients in a relatively good condition or of patients who experienced increasing problems (e.g. patients with SMA type 3b), which may have influenced the results of this study.

In conclusion, this study showed that although less restricted, patients with late onset SMA do not feel greater satisfaction with their participation in daily life than patients with early onset SMA. Compared to other diagnoses (e.g. spinal cord injury), SMA patients appear to be as satisfied with their participation in daily activities. Late onset patients reported more fatigue and experienced more pain than patients with early onset SMA. Motor skills, fatigue and feelings of depression in particular are correlates of participation in daily life. Although these findings do not fully explain variation in participation, addressing these problems may be helpful in optimizing and personalising rehabilitation care for adult patients with SMA.

Abbreviations:

SMA	spinal muscular atrophy
SMN gene	survival motor neuron
USER-P	Utrecht Scale of Evaluation Rehabilitation-Participation
HFMSE	Expanded Hammersmith Functional Motor Scale
SF-36	Short Form 36-item Health Survey
HADS	Hospital Anxiety and depression Scale
CISS-21	Coping Inventory for Stressful Situations (Short version)

REFERENCES

1. Lefebvre S, Bürglen L, Reboullet S, Clermont O, Burlet P, Viollet L et al. Identification and characterization of a spinal muscular atrophy-determining gene. *Cell* 1995;80:155-65.
2. Finkel RS, Chiriboga CA, Vajsar J, et al. Treatment of infantile-onset spinal muscular atrophy with nusinersen: a phase 2, open-label, dose-escalation study. *Lancet* 2016; 388: 3017–3026.
3. Bharucha-Goebel, D, Kaufmann, P. Treatment Advances in Spinal Muscular Atrophy. *Curr Neurol Neurosci Rep* 2017; 17: 91.
4. Parente V, Corti S. Advances in spinal muscular atrophy therapeutics. *Ther Adv Neurol Disord* 2018, Vol. 11: 1–13
5. Wadman RI, Bosboom WM, van der Pol WL, van den Berg LH, Wokke JH, Iannaccone ST, Vrancken AF. Drug treatment for spinal muscular atrophy types II and III. *Cochrane Database Syst Rev* 2012 Apr 18;(4):CD006282.
6. Darras BT. Spinal muscular atrophies. *Pediatr Clin N Am* 2015;62:743–766.

7. Zerres K, Rudnik-Schöneborn S. Natural history in proximal spinal muscular atrophy. Clinical Analysis of 445 Patients and Suggestions for a Modification of Existing Classifications. *Arch Neurol* 1995;52:518 -523.
8. Kaufmann P, McDermott MP, Darras BT et al. Prospective cohort study of spinal muscular atrophy types 2 and 3. *Neurology* 2012; 79:1889-1897.
9. Deymeer F, Serdaroglu P, Parman Y, Poda M. Natural history of SMA IIIb: Muscle strength decreases in a predictable sequence and magnitude. *Neurology* 2008;71:644-649.
10. Wadman RI, Wijngaarde CA, Stam M, Bartels B, Otto LAM, Lemmink HH, Schoenmaker MAGC, Cuppen I, Berg van den LH, Pol van der WL. Muscle strength and motor function throughout life in a cross-sectional cohort of 180 patients with spinal muscular atrophy types 1c–4. *European journal of neurology* 2018; 25:512 -518.
11. Wang CH, Finkel RS, Bertini ES, Schroth M, Simonds A, Wong B, et al. Consensus Statement for Standard of Care in Spinal Muscular Atrophy. *J Child Neurol* 2007;22:1027-1049.
12. Andries F, Wevers CWJ, Wintzen AR, Busch HFM, Höweler CJ, de Jager AEJ et al. Vocational perspectives and neuromuscular disorders. *Int J Rehabil Res* 1997;20:255-273.
13. Fowler Jr WM, Abresch RT, Koch TR, Brewer ML, Bowden RK, Wanlass RL. Employment profiles in neuromuscular diseases. *Am J Phys Med Rehabil.* 1997;76:26-37
14. Minis M-A H, Kalkman JS, Akkermans RP, Engels JA, Huijbregts PA, Bleijenberg G, Oostendorp RA, Engelen van BGM. Employment status of patients with

- neuromuscular diseases in relation to personal factors, fatigue and health status: a secondary analysis. *J Rehabil Med* 2010;42: 60-65
15. Lund ML, Lexell J. Perceived participation in life situations in persons with late effects of polio. *J Rehabil Med* 2008; 40: 659-664.
16. Jeppesen J, Madsen A, Marquardt J, Rahbek J. Living and ageing with spinal muscular atrophy type 2: Observations among an unexplored patient population. *Dev Neurorehabil* 2010; 13:10-18.
17. Lamb C, Peden A. Understanding the Experience of Living with Spinal Muscular Atrophy: A qualitative description. *J Neurosci Nurs* 2008;40:250-256.
18. Abresch, R T. Assessment of pain and health-related quality of life in slowly progressive neuromuscular disease. *American journal of hospice & palliative medicine* 2002;19:39-48.
19. Cardol M, de Jong BA, van den Bos GA, Beelen, de Groot IJ, de Haan RJ. Beyond disability: perceived participation in people with chronic disabling condition. *Clin Rehabil* 2002; 16:27-35.
20. Graham CD, Weinman J, Sadjadi R et al. A multicentre postal survey investigating the contribution of illness perceptions, coping and optimism to quality of life and mood in adults with muscle disease. *Clin Rehabil*. 2014;28:508-19.
21. Van Groenestijn AC^{1,2}, Schröder CD¹, Kruitwagen-van Reenen ET^{1,3}, van den Berg LH⁴, Visser-Meily JM^{1,3}. Participation restrictions in ambulatory ALS patients: Physical and psychological factors. *Muscle Nerve*. 2017 Jan 10. doi: 10.1002/mus.25574.
22. www.treat-nmd.eu/resources/patient-registries/SMA-national-registries/.
23. Mercuri E, Bertini W, Iannaccone ST. Childhood spinal muscular atrophy: controversies and challenges. *Lancet Neurol* 2012; 11: 443-452.

24. Piepers S, van den Berg JH, Brugman F et al (2008) A natural history study of late onset spinal muscular atrophy types 3b and 4. *J Neurol* 255:1400-1404.
25. Zee CH van der, Priesterbach AR, Dussen L van der, Kap A, Schepers VPM, Visser-Meily JMA, Post MW. Reproducibility of three self-report participation measures: the ICF measure of participation and activities screener, the participation scale and the Utrecht Scale for Evaluation of Rehabilitation-Participation. *J Rehabil Med* 2010;42: 752–757.
26. Post MW, Van der Zee CH, Hennink J, Schafrat CG, Visser-Meily JM, Van Berlekom SB. Validity of the Utrecht scale for evaluation of rehabilitation-participation. *Disabil Rehabil* 2012; 34:478-485.
27. Zee CH van der, Baars-Elsinga A, Visser-Meily JMA, Post MWM. Responsiveness of two participation measures in an outpatient rehabilitation setting. *Scand J Occup Ther* 2013; 20:201-208.
28. Zee CH van der, Post MW, Brinkhof MW, Wagenaar RC. Comparison of the Utrecht Scale for Evaluation of Rehabilitation-Participation with the ICF Measure of Participation and Activities Screener and the WHO Disability Assessment Schedule II in persons with spinal cord injury. *Arch Phys Med Rehabil* 2014; 95: 87-93.
29. Graaf JA de, Mierlo ML van, Post MWM, Achterberg WP, Kappelle LJ, Visser-Meily JMA. Long-term restrictions in participation in stroke survivors under and over 70 years of age. *Disabil Rehabil* 2017 Jan 5:1-9.
30. O'Hagen JM, Glanzman AM, McDermott MP et al. An expanded version of the hammersmith functional motor scale for SMA II and III patients. *Neuromuscul Disord* 2007;17:693-697.

31. Krosschell KJ, Maczulski JA, Crawford TO et al. A modified hammersmith functional motor scale for use in multi-center research on spinal muscular atrophy. *Neuromuscul Disord* 2006;16:417-426.
32. Ware JE. SF-36 health survey. 1993. Manual & interpretation guide. Boston: The Health Institute, New England Medical Center.
33. Aaronson NK, Muller M, Cohen PDA et al. Translation, validation, and norming of the Dutch language version of the SF-36 health survey in community and chronic disease populations. *J Clin Epidemiol* 1998;51:1055-1068.
34. Brown LF, Kroenke K, Theobald DE, Wu J. Comparison of SF-36 vitality scale and Fatigue Symptom Inventory in assessing cancer-related fatigue. *Support care cancer* 2011 19:1255-9.
35. Zigmond AS, Snaith RP. The hospital anxiety and feelings of depression scale. The hospital anxiety and feelings of depression scale. *Acta Psychiatr Scand.* 1983;67:361-70.
36. Spinhoven PH, Ormel J, Sloekers PA et al. A validation study of the hospital anxiety and feelings of depression scale (HADS) in different groups of dutch subjects. *Psychol Med* 1997; 27:363-370.
37. Taylor L, Wicks P, Leigh PN et al. Prevalence of feelings of depression in amyotrophic lateral sclerosis and other motor disorders. *Eur J Neurol* 2010;17:1047–1053.
38. Endler NS, Parker JDA. Coping inventory for Stressful situations (CISS) manual 20 (2nd edd). Toronto: Multi Health Systems. 1999.
39. Ridder DTD de, Heck GL van. Coping inventory for stressful situations. CISS handleiding. 2004. Swets test Publishers, Lisse.

40. Kruitwagen-van Reenen ET, Wadman RI, Visser-Meily JMA, van den Berg LH, Schröder C, van der Pol WL. Correlates of health related quality of life in adult patients with Spinal Muscular Atrophy. *Muscle and Nerve* 2016;54:850-855.
41. Cohen J: statistical power analysis for the behavioural sciences. 1988. New York: Academy Press.
42. Field A. Discovering statistics using IBM SPSS statistics. Thousand Oaks, California: Sage Publications. 2013. Fourth edition. 916 p
43. Ridder de D, Geenen R, Kuijer R, Middendorp van H. Psychological adjustment to chronic disease. *Lancet* 2008; 372:246-255.
44. Graham CD, Simmons Z, Stuart SR, Rose MR. Issues & Opinions ? The potential of psychological interventions to improve quality of life and mood in muscle disorders. *Muscle & Nerve* 2015; 52:131-6.
45. Peter C, Muller R, Post MW, van Leeuwen, CM, Werner CS, Geyh S. Psychological resources, appraisals, and coping and their relationship to participation in spinal cord injury: A path analysis. *Arch Phys Med Rehabil* 2014; 95:1662–1671.
46. Lizanne Eva van den Akker LE, Beckerman H, Collette EH, Bleijenberg G, Dekker J, Knoop H, de Groot V. TREFAMS-ACE study group. The role of appraisal and coping style in relation with societal participation in fatigued patients with multiple sclerosis: a cross-sectional multiple mediator analysis. *J Behav Med* 2016; 39:855–865.
47. Schwartz CE, Daltroy LH. Learning from unreliability: the importance of inconsistency in coping dynamics. *Soc Sci Med* 1999; 48: 619-631.
48. Groot de IJM, Witte de LP. Physical complaints in ageing persons with spinal muscular atrophy. *J Rehabil Med* 2005;37(4):258 -262.

Table 1. Demographics in the total population and in 2 subgroups, early versus late onset SMA

	Total	Type 1	Type 2	Type 3a	Type 3b	Type 4	Early onset	Late onset
N	62	4	21	13	20	4	38	24
Sex, female	55%	75%	62%	62%	45%	60%	63%	50%
Age in years	43.0 (20-70)	40.5 (24-51)	31.0 (20-68)	51.0 (20-66)	48.0 (20-70)	53.0 (43-69)	37.0 (20-68)	48.0 (20-70)
Partner	61%	25%	38%	69%	80%	100%	47%*	83%
Paid job	39%	25%	29%	31%	55%	50%	30%	54%
Education								
- No /Secondary school	66%	50%	72%	54%	65%	100%	64%	71%
- Higher education	34%	50%	29%	46%	35%	0	37%	29%
Living independent	68%	75%	38%	85%	80%	100%	58%	83%

Note: values are median (range). *Results differ significantly between groups (SMA types or early versus late onset) with $p < 0.05$. NA, not applicable. Education, highest grade level completed; Partner: living separate or together = yes; Living independent: independent with/without self-coordinated care = yes; Paid employment: frequency scale USER-P, Yes/No;

Table 2. Disease severity and psychological factors in the total population and in 2 subgroups, early versus late onset SMA

	Total	Type 1	Type 2	Type 3a	Type 3b	Type 4	Early onset	Late onset
N	62	4	21	13	20	4	38	24
Disease severity								
Motor skills (HFMSE)	3.5 (0-66)	0	2.0 (0-20)	4.0 (0-35)	29.0 (0-66)	48.0 (43-53)	2.0* (0-35)	32.5 (0-66) n=22
Age at diagnosis	1.5 (0-44)	0.42 (0.33-0.5)	0.75 (0-2)	1.5 (0.5-3)	6.5 (3.5-25)	40.3 (31-44)	0.95 (0-3)*	9.5 (4-44)
Loss ambulation	18.5 (2-59)	NA	NA	17.6 (4-46)	35.1 (9-59)	NA	13.9* (1-45)	34.5 (9-59)
Subjective complaints								
Pain (SF-36)	84.0 (0-100)	73.0 (61-84)	84.0 (20-100)	84.0 (31-100)	57.0 (0-100)	73.0 (41-100)	84.0* (20-100)	62.0 (0-100)
Vitality (SF-36)	62.5 (5-100)	70.0 (45-75)	70.0 (30-100)	80.0 (5-100)	55.0 (15-70)	57.5 (45-95)	75.0* (5- 100)	55.0 (15-95)

HADS-A	3.5 (0-21)	4.5 (3-9)	4.0 (0-10)	3.0 (0-9)	3.0 (0-21)	2.5 (2-3)	4.0 (0-10)	3.0 (0-21)
HADS-D	2.0 (0-15)	2.0 (1-4)	1.0 (0-9)	2.0 (0-15)	3.5 (10-12)	2.0 (1-5)	1.5 (0-15)*	3.0 (1-12)
Psychological factor								
CISS:								
- Task-oriented	26.6 (12-33)	25.5 (19-32)	26.4 (12-33)	25.7 (12-33)	27.8 (20-33)	24.7 (19-28)	27.0 (12-33)	28.0 (19-33)
- Emotional	14.6 (7-32)	12.5 (8-15)	15.3 (7-25)	12.5 (7-19)	15.7 (7-32)	14.7 (11-18)	14.1 (7-25)	15.5 (7-32)
- Avoidance	19.3 (7-31)	22.0 (16-27)	18.9 (12-28)	19.1 (7-31)	18.7 (9-27)	21.3 (11-28)	19.5 (7-31)	19.1 (9-28)

Note: values are median (range). *Results differ significantly between groups (SMA types or early versus late onset) with $p < 0.05$. NA, not applicable.

Loss of ambulation: stop walking with aid, in years.

Abbreviations: SMA, Spinal muscular Atrophy; HFMSE, Expanded Hammersmith Functional Motor Scale; MRC, Medical Research Council; SF-36, Short Form 36-item Health Survey; HADS, Hospital Anxiety and Depression Scale; HADS-A, Hospital Anxiety and Depression Scale; anxiety subscale; HADS-D, Hospital Anxiety and Depression Scale; depression subscale; CISS-21, Short Form of the Coping Inventory for Stressful Situations

Table 3. USER-P scores for the total population, and early versus late onset SMA

	SMA type						Onset	
	Total	Type 1	Type 2	Type 3a	Type 3b	Type 4	Early	Late
USER-P	N=62	n=4	n=21	n=13	n=20	n=4	n=38	n=24
Frequencies	34.5 (7-66)	32.1(21-43)	36.7 (17-49)	30.9 (7-50)	33.5 (18-48)	42.2 (23-66)	35.4 (7-50)	35.4 (18-66)
total								
- Frequency:	20.0 (0-45)	17.5 (10-40)	25.0 (0-40)	10.0 (0-30)	22.5 (0-45)	15.0 (5-35)	20.0 (0-40)	20.0 (0-45)
work/education								
- Frequency:	48.6 (14-74)	42.9 (31-54)	54.3 (31-71)	48.6 (14-74)	44.3 (26-74)	55.7 (40-65)	50.0 (14-74)	47.1 (25-74)
leisure								
activities								
Restrictions	61.9	40.6	58.2	61.1	67.1	79.4	57.6	73.3
	(18-100)	(33-59)	(36-93)	(18-89)	(30-97)	(59-100)	(18-93)*	(30-100)
Satisfaction	72.5	71.9	72.5	77.8	71.0	74.3	75.0	71.0
	(32-100)	(47-86)	(50-100)	(32-100)	(33-89)	(58-98)	(32-100)	(33-98)

Note: values are median (range).

*Results differ significant between groups (SMA types or early versus late onset) with $p < 0.05$

SMA type, SMA types 1-4; Onset, early versus late onset SMA.

Abbreviations: SMA, Spinal Muscular atrophy; USER-P, Utrecht Scale for Evaluation Rehabilitation-Participation.

Frequency, work/education: hours/ week; Frequency, leisure activities: times/month.

Table 4. USER-P, Frequency scale: hours per week (work/education) and times per month (leisure activities) spent per item, for the total population, N = 62

Domains	Hours/week		
	Not at all	1-24 hrs	≥25 hrs
Paid work	61%	18%	21%
Unpaid work	44%	42%	15%
Education	71%	18%	12%
Household	34%	60%	7%
	Times/month		
	Not at all	1-10 times	>10 times
Sports and physical exercise	44%	42%	15%
Going out	10%	87%	2%
Daytrips	11%	81%	8%
Leisure activities at home	5%	39%	57%

Visiting family or friends	3%	86%	11%
Receiving visitors	3%	87%	10%
Contact phone, computer	0	16%	84%

Note: Education: only activities in the course of work, or in order to get work.

Abbreviations: USER-P, Utrecht Scale for Evaluation Rehabilitation-Participation

Table 5. Percentage of patients with perceived restrictions and dissatisfaction, for the total population, N=62

Restrictions	%	Dissatisfaction	%
Persisting problems			
Work/education	67 % (n=52)	Work/education	15 % (n=39)
Household chores	97 % (n=59)	Household chores	43 % (n=56)
Mobility outdoors	82%	Mobility outdoors	31 % (n=61)
Physical exercise	95 % (n=57)	Physical exercise	58 % (n=50)
Going out	71%	Going out	31 % (n=61)
Outdoor activities	79%	Outdoor activities	31 % (n=61)
Leisure indoors	41 % (n=61)	Leisure indoors	12 % (n=61)
Relationship partner	33 % (n=40)	Partner relationship	7 % (n=39)
Visits to family/friends	66%	Family relationships	10%
Visits from family/friends	20 % (n=61)	Friends and acquaintances	8%
Telephone/computer contact	16 %		

(n.): n per item NB: Not all items were scored by all patients.

Table 6. Bivariate and multivariable linear regression analyses for USER-P frequency, restrictions and satisfaction. Stepwise regression.

	Frequency of participation			Participation restrictions			Satisfaction with participation		
	Bivariable	Multivariable		Bivariable	Multivariable		Bivariable	Multivariable	
		Step 1	Step 2		Step 1	Step 2		Step 1	Step 2
Characteristics	ρ	β	β	ρ	β	β	ρ	β	β
Sex, female	0.08	N.E.	N.E.	0.05	N.E.	N.E.	-0.03	N.E.	N.E.
Age in years	-0.28*	-0.24	-0.05	-0.10	N.E.	N.E.	-0.01	N.E.	N.E.
Partner	0.03	N.E.	N.E.	0.11	N.E.	N.E.	0.06	N.E.	N.E.
Education, High	0.05	N.E.	N.E.	0.06	N.E.	N.E.	0.32*	-0.37	-0.20
Living independent	-0.13	N.E.	N.E.	-0.01	N.E.	N.E.	-0.13	N.E.	N.E.
Motor skills (HFMSE)	0.26*	0.27*	0.32*	0.59*	0.45*	0.47*	-0.04	N.E.	N.E.
Pain (SF-36)	0.28*	N.E.	0.06	0.04	N.E.	N.E.	0.21	N.E.	N.E.

Fatigue (SF-36)	0.19	N.E.	N.E.	-0.01	N.E.	N.E.	0.42*	N.E.	0.30*
HADS-A	0.01	N.E.	N.E.	-0.05	N.E.	N.E.	-0.29	N.E.	N.E.
HADS-D	-0.41*	N.E.	-0.44*	-0.24*	N.E.	-0.23	-0.49*	N.E.	-0.41*
CISS									
Task-oriented	-0.06	N.E.	N.E.	-0.14	N.E.	N.E.	-0.03	N.E.	N.E.
Emotional	0.21	N.E.	N.E.	-0.11	N.E.	N.E.	-0.00	N.E.	N.E.
Avoidance	-0.09	N.E.	N.E.	-0.05	N.E.	N.E.	0.02	N.E.	N.E.
ΔR^2		0.14*	0.19*		0.21*	0.05		0.14*	0.36*
R²		0.33			0.26			0.50	

Note: ΔR^2 , change in R^2 between two equations; R^2 , explained variance.

*, all included variables significant correlation with outcome measure $p < 0.1$.

Abbreviations: USER-P, Utrecht Scale for Evaluation Rehabilitation-Participation; HFMSE, Expanded Hammersmith Functional Motor Scale; SF-36, Short Form 36-item Health Survey; HADS-A, Hospital Anxiety and Depression Scale; anxiety subscale; HADS-D, Hospital Anxiety and Depression Scale; depression subscale; CISS-21, Short Form of the Coping Inventory for Stressful Situations.