University of Groningen

Altruism and infidelity among warblers

Richardson, David S.; Komdeur, Jan; Burke, Terry

Published in:
Nature

DOI:
[10.1038/422580a](https://doi.org/10.1038/422580a)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2003

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Richardson, D. S., Komdeur, J., & Burke, T. (2003). Altruism and infidelity among warblers. *Nature*, 422(6932), 580-580. <https://doi.org/10.1038/422580a>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Avian behaviour

Altruism and infidelity among warblers

In cooperatively breeding vertebrates, indirect fitness benefits^{1,2} are maximized by subordinates who choose to help their own closely related kin after accurately assessing their relatedness to the group's offspring³⁻⁵. Here we show that in the cooperatively breeding Seychelles warbler (*Acrocephalus sechellensis*), female subordinates help to raise new nestlings by providing them with food only when the offspring are being raised by parents who also fed the subordinates themselves when they were young³. These helper females use the continued presence of the primary female, rather than of the primary male, as their provisioning cue — presumably because female infidelity is rife in this species⁶, making their relatedness to the father less reliable.

In *A. sechellensis*, female offspring normally become subordinate within their natal group^{7,8} and often help to raise dependent young⁷. We studied the provisioning behaviour (feeds given per hour) of 21 colour-ringed female subordinates. Reproductive sharing and high levels of extra-pair paternity occur in this species⁹, we therefore assessed parentage and genetic relatedness by using microsatellite DNA markers⁸. Subordinate parents (that is, subordinates who had also produced a nestling in the helped nest) always provisioned and were excluded from subsequent analysis ($n = 8$).

In a general linear model analysis, the provisioning behaviour of non-parent subordinates was influenced positively by subordinate–nestling relatedness ($F_{1,11} = 4.82$, $P = 0.05$, $R^2 = 0.31$), whereas subordinate age, sex, number of nestlings, number of subordinates (or feeding adults) and territory quality had no significant effect. A comparison of relatedness⁹ to the nestlings of subordinates who helped with those that did not (Fig. 1a) indicates that helpers were significantly related to nestling(s), whereas non-helpers were not (mean relatedness, 0.27 versus -0.05 , respectively).

The continued presence of the primary female, but not of the primary male, was a reliable cue of subordinate–nestling relatedness; subordinate–primary female relatedness predicted subordinate–nestling relatedness ($F_{1,38} = 27.37$, $P < 0.001$), but subordinate–primary male relatedness did not ($F_{2,37} = 2.11$, $P = 0.16$). Subordinates provisioned significantly more often when their putative mother was still present, but provisioning was unaffected by the presence or absence of the putative father (Fig. 1b).

Subordinates may determine when to provision by directly assessing their relatedness to either the nestling(s) or the primary

Figure 1 Provisioning by subordinate female, non-parent Seychelles warblers (*Acrocephalus sechellensis*). **a**, Provisioning in relation to the pairwise relatedness between subordinates and nestlings (subordinate–helper–nestling relatedness versus non-related: 0.27 ± 0.16 versus 0 , one-sample t -test, $t_7 = 4.64$, $P < 0.002$; helpers versus non-helpers: 0.27 ± 0.16 , $n = 8$ versus -0.05 ± 0.20 , $n = 5$; $t_{11} = 3.13$, $P = 0.01$). **b**, Provisioning in relation to the presence (purple bars) or absence (green bars) of the subordinate's putative parents (presence versus absence of putative mother: 7.57 ± 2.44 , $n = 6$ versus 1.43 ± 3.60 , $n = 7$; t -test, $t_{11} = 3.71$, $P = 0.003$; presence versus absence of putative father: 4.30 ± 4.89 , $n = 4$ versus 4.24 ± 4.29 , $n = 8$, $t_{11} = 0.02$, $P = 0.98$). Error bars, means $\pm$ s.e.; asterisks indicate statistical significance.

female. However, the continued presence of the primary female explained provisioning more reliably ($F_{1,11} = 13.72$, $P = 0.003$, $R^2 = 0.55$) than genetic estimates of subordinate–nestling or subordinate–primary female relatedness ($F_{1,11} = 2.14$, $P = 0.17$; $F_{1,11} = 0.06$, $P = 0.81$, respectively). These results confirm that the presence of the primary female who raised the subordinate is likely to be used as a cue to determine when to provision. Previous findings that helpers can distinguish between siblings and half-siblings³ may have been an artefact caused by helpers assisting only their mothers.

Our results show that female subordinates can use an indirect but reliable cue to assess their relatedness to nestlings, and that this assessment of kinship determines provisioning rates. In *A. sechellensis*, the amount of food brought to the nestling determines fledging success and first-year survival, and the presence of a helper increases the number of young who are fledged on a territory¹⁰. The preferential provisioning of related nestlings by female subordinates will therefore increase the

number of related offspring produced. In our study, the number of fledglings produced on a territory increased significantly (by 17%) for each subordinate present (after exclusion of direct parentage)⁸.

Both direct⁸ and indirect benefits may be important within this cooperative breeding system as, by using effective discrimination, subordinates are able to maximize the indirect benefit gained within a system that is driven primarily by direct benefits⁸. Our findings show that, in the presence of a high frequency of female infidelity, an associative learning mechanism has evolved to focus on the mother at the nest — the only sex to which the subordinates are reliably related.

David S. Richardson^{*†}, Jan Komdeur[‡], Terry Burke^{*}

^{*}Department of Animal and Plant Sciences, University of Sheffield, Sheffield S10 2TN, UK

[†]Department of Animal Ecology, University of Lund, 223 62 Lund, Sweden

e-mail: david.richardson@zoekol.lu.se

[‡]Zoological Laboratory, University of Groningen, PO Box 14, 9750 AA Haren, The Netherlands, and Department of Zoology, University of Melbourne, Victoria 3010, Australia

- Hamilton, W. D. *J. Theor. Biol.* **7**, 1–52 (1964).
- Emlen, S. T. *Proc. Natl Acad. Sci. USA* **92**, 8092–8099 (1995).
- Komdeur, J. *Proc. R. Soc. Lond. B* **256**, 47–52 (1994).
- Russell, A. F. & Hatchwell, B. J. *Proc. R. Soc. Lond. B* **268**, 2169–2174 (2001).
- Hatchwell, B. J., Ross, D. J., Fowles, M. K. & McGowan, A. *Proc. R. Soc. Lond. B* **268**, 885–890 (2001).
- Richardson, D., Jury, F., Blaakmeer, K., Komdeur, J. & Burke, T. *Mol. Ecol.* **10**, 2263–2273 (2001).
- Komdeur, J. *Proc. R. Soc. Lond. B* **263**, 661–666 (1996).
- Richardson, D. S., Burke, T. & Komdeur, J. *Evolution* **56**, 2313–2321 (2002).
- Goodnight, K. F. & Queller, D. C. *Mol. Ecol.* **8**, 1231–1234 (1999).
- Komdeur, J. *Behav. Ecol. Sociobiol.* **34**, 175–186 (1994).

Competing financial interests: declared none.

COMMUNICATIONS ARISING

Plant ecology

Tree-species competition and coexistence

How apparently similar plant species coexist is a puzzle. Kelly and Bowler¹ claim that environmental fluctuation promotes the coexistence of tree species by alternately favouring recruitment of common and rare congeners in a dry tropical forest. Here I argue that current knowledge of tropical-forest ecology does not support the authors' focus on congeneric competition, and show that their use of diameter distributions to date recruitment fluctuations may be misleading. It is therefore doubtful, at this stage, that recruitment patterns of the authors' congeneric pairs can be linked to the sort of competition dynamic that they envisage.

Kelly and Bowler's two-component lottery model assumes that seedlings of