University of Groningen

Phosphatases modulate the bistable sporulation gene expression pattern in *Bacillus subtilis*

Veening, JW; Hamoen, LW; Kuipers, OP

Published in:
Molecular Microbiology

DOI:
[10.1111/j.1365-2958.2005.04659.x](https://doi.org/10.1111/j.1365-2958.2005.04659.x)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2005

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Veening, JW., Hamoen, LW., & Kuipers, OP. (2005). Phosphatases modulate the bistable sporulation gene expression pattern in *Bacillus subtilis*. *Molecular Microbiology*, 56(6), 1481-1494.
<https://doi.org/10.1111/j.1365-2958.2005.04659.x>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Phosphatases modulate the bistable sporulation gene expression pattern in *Bacillus subtilis*

Jan-Willem Veening, Leendert W. Hamoen,[†] and Oscar P. Kuipers*

Department of Genetics, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, Kerklaan 30, 9751 NN Haren, the Netherlands.

Summary

Spore formation in the Gram-positive bacterium *Bacillus subtilis* is a last resort adaptive response to starvation. To initiate sporulation, the key regulator in this process, Spo0A, needs to be activated by the so-called phosphorelay. Within a sporulating culture of *B. subtilis*, some cells initiate this developmental program, while other cells do not. Therefore, initiation of sporulation appears to be a regulatory process with a bistable outcome. Using a single cell analytical approach, we show that the autostimulatory loop of *spo0A* is responsible for generating a bistable response resulting in phenotypic variation within the sporulating culture. It is demonstrated that the main function of RapA, a phosphorelay phosphatase, is to maintain the bistable sporulation gene expression. As *rapA* expression is quorum regulated, it follows that quorum sensing influences sporulation bistability. Deletion of *spo0E*, a phosphatase directly acting on Spo0A~P, resulted in abolishment of the bistable expression pattern. Artificial induction of a heterologous Rap phosphatase restored heterogeneity in a *rapA* or *spo0E* mutant. These results demonstrate that with external phosphatases, *B. subtilis* can use the phosphorelay as a tuner to modulate the bistable outcome of the sporulating culture. This shows that *B. subtilis* employs multiple pathways to maintain the bistable nature of a sporulating culture, stressing the physiological importance of this phenomenon.

Introduction

Spore formation in *Bacillus subtilis* is a complex adaptive response to starvation (Sonenshein, 2000). The process

of sporulation is governed by a multicomponent phosphorelay which consists of five histidine kinases (KinA, KinB, KinC, KinD and KinE) and two phosphorelay proteins (Spo0F and Spo0B) (Perego and Hoch, 2002) (see Fig. 1). Multiple environmental and physiological signals are fed into this system, and under appropriate conditions this leads to phosphorylation of Spo0A, the key sporulation transcription factor (for reviews see Grossman, 1995; Sonenshein, 2000). Within an isogenic culture of sporulating *B. subtilis*, some cells initiate the developmental program of sporulation, whereas others do not (Chung *et al.*, 1994). Therefore, initiation of sporulation appears to be a regulatory process with a bistable outcome. Phenotypic variation could benefit the fitness of the species, because the heterogeneous population is able to quickly react to changing environments (Balaban *et al.*, 2004). As sporulation is an energy intensive process, and irreversible after its earliest stage (Parker *et al.*, 1996), cells that are delayed to commit to sporulate could have an advantage over sporulating cells if food resources were to become plentiful again. Theoretical modelling and experiments in both prokaryotic and eukaryotic model systems have demonstrated that positive feedback of a transcriptional regulator, together with a non-linear response to an activator, can lead to a bimodal probability distribution in expression (Hasty *et al.*, 2000; Becskei *et al.*, 2001; Isaacs *et al.*, 2003). It is believed that stochastic fluctuations causes some cells to reach the threshold level to activate the feedforward loop and these cells will end up in the 'high expressing' population, whereas others do not reach this threshold and remain in the 'low expressing' state (Ferrell, 2002; Rao *et al.*, 2002). In the present study, we examined whether the positive feedback architecture of *spo0A* regulation is responsible for bistability in sporulation gene expression and studied the influence of various phosphatases on the maintenance of bistability.

Activity of Spo0A is subject to several auto-stimulatory loops (Strauch *et al.*, 1992; Strauch *et al.*, 1993; Fujita and Sadaie, 1998). These loops involve transcription of *spo0A* and phosphorylation of Spo0A. Transcription of *spo0A* is directly activated by Spo0A~P and indirectly activated by induced expression of *sigH*. The *sigH* gene encodes an RNA polymerase sigma factor (σ^H) that recognizes an alternative promoter located upstream of *spo0A* (Predich *et al.*, 1992) and activates transcription of genes involved in the phosphorylation of Spo0A such as *kinA* and *spo0F*.

Accepted 18 March, 2005 *For correspondence. E-mail o.p.kuipers@rug.nl; Tel. (+31) 50 3632 093; Fax (+31) 50 3632 348.

[†]Present address: Sir William Dunn School of Pathology, University of Oxford, Oxford OX1 3RE, UK.

Fig. 1. Simplified schematic representation of the regulatory network that governs initiation of sporulation. Perpendiculars and arrows represent the negative and positive actions respectively.

Furthermore, sigma-H activates transcription of the *spoIIA* operon, which contains the sporulation specific sigma factor, sigma-F (Hoch, 1991). A major role of phosphorylated Spo0A is to repress the expression of *abrB*, a gene encoding a transcriptional regulator that represses various stationary phase processes (Robertson *et al.*, 1989). During exponential growth, AbrB represses expression of *sigH*, *kinA* and *abrB* itself (Strauch, 1995). Thus, alleviation of AbrB repression by Spo0A~P at the beginning of the stationary growth phase, stimulates *sigH* and *kinA* expression and therefore *spo0A* transcription and indirectly phosphorylation of Spo0A (see Fig. 1). In conclusion, the complex autostimulation of *spo0A* could be the basis of the bistable sporulation gene expression.

KinA is the primary kinase in the phosphorelay and is necessary for the phosphorylation of Spo0A (Burbulys *et al.*, 1991). It has been demonstrated that the fraction of cells that initiate sporulation is decreased in a *kinA* mutant background (Chung *et al.*, 1994). This result suggests that a certain threshold concentration of Spo0A~P is necessary to initiate sporulation and that the activity of the phosphorelay determines the threshold level for autostimulation of Spo0A. This implies that influences on the phosphorelay by external phosphatases could alter the heterogeneous sporulation gene expression. An interesting candidate is RapA. RapA is an aspartyl-phosphatase that dephosphorylates one of the components of the phosphorelay, Spo0F~P, and consequently reduces accumulation of Spo0A~P (Perego and Hoch, 2002) (see Fig. 1). In an earlier study it was shown that *rapA*, by that time known as *gsiAA* (glucose starvation induced protein

A), was activated under glucose limiting conditions and high cell densities, signals which also trigger initiation of sporulation (Mueller *et al.*, 1992). Apparently, RapA increases in concentration coincidentally with the activation of Spo0A~P. This seeming contradiction suggests that RapA might play a role in the heterogeneous induction of sporulation. This possibility is especially interesting because RapA activity is subject to a complex quorum sensing regulation. First of all, expression of *rapA* is activated by the ComA-ComP two-component signal transduction system, which is activated by cell density signals via the quorum sensor molecule, ComX (Mueller *et al.*, 1992; Lazazzera *et al.*, 1999). This regulatory system is essential for another stationary growth phase process; the development of genetic competence. Secondly, the phosphatase activity of RapA is negatively modulated by a small protein, PhrA, encoded by an adjacent gene on the same transcript as *rapA*. PhrA is exported by the SecA-dependent system and subsequently processed to the carboxyl-terminal five-amino acid active inhibitor (ARNQT). The pentapeptide presumably functions as a quorum sensor signal. PhrA is re-imported by the oligopeptide permease transport system (Opp) and inhibits the dephosphorylating action of RapA (Perego and Hoch, 2002). Deletion of the *phrA* gene causes uncontrolled RapA phosphatase activity leading to poor sporulation (Perego and Hoch, 1996).

To study the role of the RapA-PhrA system and the autostimulation of Spo0A in sporulation heterogeneity, a single cell approach was used. By flow cytometric analyses and fluorescence microscopy, we demonstrate that the autostimulatory activation of *spo0A* is responsible for the bistable expression pattern in sporulating cultures. In addition, we show that external phosphatases, such as RapA, Spo0E and Rap60, can modulate sporulation bistability.

Results

Initiation of sporulation is heterogeneous

When *B. subtilis* cells reach the end of exponential growth, and various environmental signals promote the activation of the sporulation response regulator Spo0A, this protein in turn activates more than 40 genes directly, including the sigma-H dependent *spoIIA* operon comprising the early sporulation genes *spoIIAA*, *spoIIAB* and *sigF* (Molle *et al.*, 2003). To determine how the initiation of sporulation is distributed among cells within an isogenic population, we constructed a strain in which the *gfp* gene (encoding Green Fluorescent Protein) is under control of the *spoIIA* promoter (integrated at the *spoIIA* promoter region, leaving the original operon intact). The resulting strain, IIA-*gfp* (*P_{spoIIA}-gfp*), was grown in sporulation medium (SM), and

Fig. 2. Heterogeneous expression of $P_{spoIIA}\text{-}gfp$. Cells were grown in sporulation medium and collected for flow cytometric analysis. Time is given in hours relative to the transition point between the exponential and stationary growth phase (T0). As a control, the parental *B. subtilis* 168 strain was analysed as well (indicated with the asterisk). Arrows indicate cells in a low and high *spoIIA* expression state.

cells were collected at different time points for analysis by flow cytometry (Fig. 2). As a control for GFP specific fluorescence, the parental *B. subtilis* 168 strain was also analysed. Growth curves related to these experiments can be found in the *Supplementary material* (Appendix S1 and Fig. S1). During exponential growth (T-1), production of GFP was not detected. At the beginning of the stationary growth phase (T0), expression of GFP from the *spoIIA* promoter could be observed. This expression was not equal in all cells, confirming previous results (Chung *et al.*, 1994). After one hour from the transition between exponential and stationary growth (T1), cells are in a low (Fig. 2, left peak) or high expressing state (Fig. 2, right peak). The flow cytometric graphs show relative numbers of cells, since part of the culture lysis during sporulation. It is assumed that cells that are unable to activate Spo0A (in time) are vulnerable to lysis (Gonzalez-Pastor *et al.*, 2003; see *Discussion*). These analyses show that two distinct subpopulations are present early after entry into the stationary growth phase: cells in a low *spoIIA* expression state and cells in a high *spoIIA* expression state.

Activation of the *spoIIA* promoter is not subject to transcriptional noise

Previous studies have shown that substantial stochastic fluctuations in promoter activities may account for non-genetic variability within isogenic populations (McAdams and Arkin, 1997; Elowitz *et al.*, 2002; Ozbudak *et al.*, 2002; Blake *et al.*, 2003; Raser and O'Shea, 2004). Theoretically, such stochastic effects could contribute to the large cell-to-cell variation in *spoIIA* expression. Intrinsic noise of a given gene can be defined as the extent to which the activities of two identical copies of that gene, in the same intracellular environment, fail to correlate. To

verify that the observed heterogeneity is not a consequence of intrinsic transcriptional noise of the *spoIIA* promoter itself, a double-labelled strain that enables the detection of stochasticity in gene expression was constructed. Therefore, the *spoIIA* promoter was fused to improved variants of the cyan- and yellow fluorescent proteins (ICFP and IYFP, respectively, Veening *et al.*, 2004). In the resulting strain (IIA/double), the $P_{spoIIA}\text{-}icfp$ construct was integrated at the *amyE* locus and the $P_{spoIIA}\text{-}iyfp$ construct was integrated at the *spoIIA* promoter region (a map showing the chromosomal locations of the constructs is indicated in Fig. S2 of the *Supplementary material*). The experimental set-up and fluorescence microscopy measurements were performed in a similar fashion as described by Elowitz *et al.* (2002). They showed that in the absence of intrinsic noise, the CFP and YFP signals fluctuate in a correlated fashion over time in a single cell. However, the amount of fluorescent proteins produced will differ between cells because of extrinsic noise. The correlated YFP and CFP signals in the individual cells in Fig. 3, suggest that the levels of intrinsic noise are negligible for the *spoIIA* promoter (cells that exhibit ICFP production, expressed from the *spoIIA* promoter at the *amyE* locus, show a similar level of IYFP expressed from the *spoIIA* promoter at the *spoIIA* promoter region). The results of a more quantitative analysis are presented in Table 1. To quantify noise, a statistical analysis was performed in a similar fashion as described previously (Elowitz *et al.*, 2002). Strain IIA/double was grown in SM and cells were collected for fluorescent microscopy two hours after entry into stationary growth phase. Images were analysed and noise data were calculated as described in the *Experimental procedures*. Table 1 shows the calculated noise levels for strain IIA/double. The value for intrinsic noise (η_{int}) in strain IIA/double was approximately 0.09, indicating that the two colours, on average, differ by about 0.09% within individual cells. In the work of Elowitz *et al.* (2002), similar intrinsic noise values were obtained with *Escherichia coli* strains that displayed low levels of intrinsic noise (using strong constitutive promoters). The extrinsic noise levels were somewhat higher compared with the levels of intrinsic noise ($\eta_{ext} \sim 0.13$) but still quite low. This indicates that once a cell is activated, the expression level reaches similar heights among cells. Because the measured extrinsic noise is significantly

Table 1. Noise of the *spoIIA* promoter.

	Intrinsic noise, $\eta_{int} (\times 10^{-2})$	Extrinsic noise, $\eta_{ext} (\times 10^{-2})$	Total noise, $\eta_{tot} (\times 10^{-2})$
IIA/double ^a	9.0 ± 0.8	13.1 ± 2.6	16.1 ± 2.4

a. Measurements are from four independent experiments in which at least 100 cells per experiment were counted; $\pm$ indicates the standard error in these data.

Fig. 3. Transcriptional noise of the *spoIIA* promoter. Strain IIA/double was grown in sporulation medium and cells were collected for fluorescence microscopy two hours after entry into the stationary growth phase. IYFP and ICFP images were combined in the red and green channels respectively. The left panel shows IYFP produced from activation of the *spoIIA* promoter construct integrated at the *spoIIA* locus (red cells) and the middle panel indicates ICFP produced from the P_{spoIIA} -*icfp* construct integrated at the *amyE* locus (green cells). An overlay of both channels is depicted in the right panel. Yellow cells express both transcriptional units.

higher than the intrinsic noise, it is the major source for the total observed variation (η_{tot}) because the three values are related as $\eta_{int}^2 + \eta_{ext}^2 = \eta_{tot}^2$ (Elowitz *et al.*, 2002). Overall, these results indicate that intrinsic noise does not play a role in the observed heterogeneity of *spoIIA* gene expression, but is a consequence of specific regulation.

Spo0A autoactivated cells initiate sporulation

To confirm that the *spoIIA* promoter activity is a good indicator for cells that activate Spo0A, we constructed a strain that enables the visualization of activation of both *spoIIA* and *spo0A* transcriptional units. The double-labelled strain *spo0A*/IIA (P_{spo0A} -*iyfp*, $amyE::P_{spoIIA}$ -*icfp*) was grown in SM and collected at hourly intervals for fluorescence microscopic analyses. After entry into the stationary growth phase, production of IYFP from the *spo0A* promoter could be observed in most cells (Fig. 4).

However, some cells clearly show stronger *spo0A* expression. These cells also show high activation of the *spoIIA* promoter, indicating that activation of the *spoIIA* promoter can be used to monitor the Spo0A activation state.

The Spo0A autostimulatory loop is essential for sporulation bistability

As described in the introduction, positive feedback can be the basis for bistable gene expression. To examine whether Spo0A autoactivation is the mechanism responsible for bistable induction of sporulation, we bypassed the transcriptional autostimulation by replacing the wild-type *spo0A* promoter with an IPTG inducible promoter. In addition, a constitutively active form of Spo0A (Spo0A-Sad67) was used in order to bypass the autostimulatory effects on the phosphorelay (Ireton *et al.*, 1993). All strains contained the P_{spoIIA} -*gfp* reporter. Strains IIA/*spo0A* + (P_{spoIIA} -

Fig. 4. Activation of *spo0A* and *spoIIA* within a sporulating culture. Strain *spo0A*/IIA was grown in sporulation medium and collected for fluorescence microscopy one hour after entry into the stationary growth phase. IYFP and ICFP images were combined in the red and green channels respectively. The left panel shows IYFP produced from activation of the *spo0A* promoter (red cells) and the middle panel indicates ICFP produced from the *spoIIA* promoter (green cells). An overlay of both channels is depicted in the right panel. Yellow cells express both transcriptional units.

gfp, P_{spac} -*spo0A*), *IIA/sad67* (P_{spolIA} -*gfp*, P_{spac} -*spo0A*-*sad67*) and *IIA/spo0A/sad67* (P_{spolIA} -*gfp*, $\Delta spo0A$, P_{spac} -*spo0A*-*sad67*) were grown at varying IPTG concentrations, and cells were collected for flow cytometric measurements and immunoblot analyses (Fig. 5). To reduce possible effects of the phosphorelay, cells were grown in rich medium and samples were taken at the logarithmic growth phase. As depicted in panel A, the *spolIA* promoter could not be activated by induction of wild-type Spo0A. In accordance with this, the intracellular Spo0A levels in strain *IIA/spo0A* + accumulate with increasing IPTG concentrations (Fig. 5D, upper panel). This demonstrates that unphosphorylated Spo0A (Spo0A is not activated under these growth conditions) is not able to activate the *spo0A* autostimulatory loop. This is in agreement with previous results (Ireton *et al.*, 1993). Figure 5B shows the expression patterns of *spolIA* expression upon the induction of the constitutively active form of Spo0A (Sad67) in a wild-type background in which the autostimulatory loop is present. At low IPTG levels, all cells are in the low expression state, whereas at high induction levels all cells are in the high expression state. At intermediate induction levels (with a threshold concentration of approximately 50 μ M of IPTG), a clear bistable expression pattern can be observed. In agreement with this, immunoblot analysis of strain *IIA/sad67* (Fig. 5D, middle panel) shows that the *spo0A* autostimulatory loop is activated when the constitutively active Spo0A-Sad67 is induced at levels higher than 50 μ M of IPTG, indicated by an increase of wild-type Spo0A. It should be noted that the Sad67 protein migrates faster than wild-type Spo0A because of the N-terminal deletion. This demonstrates that Spo0A-Sad67 is not only able to activate transcription of *spo0A*, but also results in increased Spo0A phosphorylation. This is most likely caused by increased expression of phosphorelay genes like *kinA* and *spo0F* (Hoch, 1991). In contrast to this, when wild-type *spo0A* is deleted and only the IPTG inducible *spo0A-sad67* is present, a graded response in *spolIA* gene expression can be observed (Fig. 5C). At low IPTG concentrations, little activation of the *spolIA* promoter could be observed. Upon increasing IPTG levels, the population fluorescence is shifted to higher intensities in a Normal-Distribution fashion without the occurrence of a dual population. Also, at maximal induction, the average

Fig. 5. Bistability requires a positive feedback loop. Strains containing the P_{spolIA} -*gfp* reporter fusion were grown for 4 h in TY medium containing varying concentrations of IPTG and samples were analysed on a flow cytometer. Fluorescence distributions of strain *IIA/spo0A* + (P_{spolIA} -*gfp*, P_{spac} -*spo0A*) (A), strain *IIA/sad67* (P_{spolIA} -*gfp*, P_{spac} -*sad67*) (B) and (P_{spolIA} -*gfp*, P_{spac} -*sad67*, $\Delta spo0A$) (C) grown in the presence of the indicated IPTG concentration. D shows an immunoblot analysis using polyclonal anti-Spo0A antibodies and polyclonal anti-GroEL antibodies on whole-cell extracts which were subjected to SDS-PAGE (12%) and immunoblotting. The anti-GroEL antibody served as a control for loading.

fluorescence intensity of strain IIA/*spo0A*/*sad67* does not reach the level of strain IIA/*sad67*, in which the autoloop is present. This trend is also seen in the corresponding immunoblot analysis (Fig. 5D, lower panel) in which upon increasing IPTG levels a linear increase in *Sad67* protein levels is shown, and wild-type *Spo0A* is not present because of the *spo0A* mutation. In summary, these results demonstrate that the *spo0A* autostimulatory loops are responsible for the bistable expression of *spoIIA*.

RapA modulates sporulation bistability

RapA lowers the phosphorelay activity and could influence the bistable sporulation gene expression as described in the introduction. In one of the first studies on *rapA* it was shown that the expression of this gene was strongly repressed by *Spo0A*-P (Mueller and Sonenshein, 1992). Because expression of *rapA* is activated by factors outside the *Spo0A*-P regulatory network (ComA/P/X) but is repressed by *Spo0A*-P, it is conceivable that cells which do not reach the threshold level of *Spo0A*-P in time, will

accumulate high levels of RapA, thus delaying sporulation initiation. To examine whether there is a correlation between activation of *rapA* and initiation of sporulation, the *rapA* promoter was fused to the *iyfp* gene, and this construct was combined with a strain carrying the *spoIIA* promoter fused to the *icfp* gene. The resulting strain, *rapA*/IIA (*P_{rapA}-iyfp*, *amyE::P_{spoIIA}-icfp*), was grown in SM and cells were collected for fluorescence microscopy. As shown in Fig. 6, at the end of logarithmic growth cells begin to express *rapA*, albeit at low levels. At this time-point (*T* = -1), activation of the *spoIIA* operon cannot be observed. Later in growth (*T* = 2), some cells have activated the *spo0A* autostimulatory loop and highly express *spoIIA*. Cells that do not show clear activation of *spoIIA*, show a strong induction of *rapA* expression. In the *spoIIA* active cells, this *rapA* induction is clearly absent. Thus, expression of the two genes is reciprocally distributed within the culture (Fig. 6C–E). To further substantiate these observations, strain *rapA-iyfp* (*P_{rapA}-iyfp*) was subjected to flow cytometric analyses (Fig. 6F). Again, it is shown that at the end of logarithmic growth, cells express

Fig. 6. Reciprocal expression of *rapA* and *spoIIA*. Strain *rapA*/IIA was grown in sporulation medium and cells were collected one hour before (*T* = -1) and two hours after (*T* = 2) entry into the stationary growth phase for analysis by fluorescence microscopy. IYFP and ICFP images were combined in the red and green channels respectively. Red cells produce IYFP of which expression is driven by activity of the *rapA* promoter; green cells represent production of ICFP, of which expression is driven by activity of the *spoIIA* promoter.

A. Light phase contrast picture of cells at *T* = -1.

B. IYFP image of cells from panel A.

C–D. Fluorescent microscopic graphs of cells from *T* = 2.

E. An overlay of panels C and D.

F. Strain *rapA-iyfp* was grown in sporulation medium and cells were collected one hour before (*T* = -1) and two hours after (*T* = 2) entry into the stationary growth phase for analysis by flow cytometry. Because YFP has different spectral properties compared with GFP, settings of the flow cytometer were set more sensitive. Therefore, the parental 168 strain was also included in the fluorescence histogram (*T* = 2).

YFP from the *rapA* promoter at low levels but in a mono-modal distribution. In the stationary phase, a fraction of cells express *rapA* at high levels and the other fraction at a low level.

Because induction of *rapA* seems to prevent activation of *spoIIA*, a mutation in *rapA* could have an effect on sporulation bistability. To test this hypothesis, we introduced a *rapA* mutation in our reporter strain. The resulting strain, IIA-gfp/ $\Delta rapA$ (P_{spoIIA} -gfp, $\Delta rapA$), was grown in SM and cells analysed by flow cytometry. As shown in Fig. 7A, the final bistable expression of *spoIIA* almost completely disappeared in a *rapA* mutant background, and the vast majority of cells express high levels of *spoIIA*. Figure 7B shows a quantitative analysis of the effect of a *rapA* muta-

Fig. 7. The effect of $\Delta rapA$ on P_{spoIIA} -gfp expression. Strains IIA-gfp and IIA-gfp/ $\Delta rapA$ were grown in sporulation medium and collected for flow cytometric analysis. Time is given in hours relative to the transition point between the exponential and stationary growth phase (T0).

A. Fluorescence distributions of cells collected at T2.

B. The fraction of cells in a high *spoIIA* state, relative to the total measurable population, is indicated. Three independent transformants of each strain were analysed and error bars depict the standard error in these experiments.

Fig. 8. The effect of $\Delta spo0E$ on P_{spoIIA} -gfp expression. Strains IIA-gfp and IIA-gfp/ $\Delta spo0E$ were grown in sporulation medium and collected for flow cytometric analysis. The fluorescence distributions of cells collected at two hours after entry into the stationary growth phase (T = 2) is depicted.

tion. When we consider cells with signals higher than fluorescent channel 670 to be in a high expressing state, approximately 70% of the wild-type population reaches the *spoIIA* high expression state, compared with more than 90% in the *rapA* mutant. Initially, induction of sporulation is still heterogeneous and a similar fraction of cells highly express *spoIIA* in the *rapA* mutant when compared with the wild type (Fig. 7B; T0). This indicates that sporulation bistability is not caused by RapA, but stabilized by RapA.

Phosphorelay phosphatases modulate the bistable outcome

The dephosphorylating action of RapA reduces the activity of the phosphorelay and consequently will raise the threshold level of Spo0A-P to fully initiate autostimulation. This suggests that other phosphatases that act on the sporulation phosphorelay, such as Spo0E, have the potential to alter sporulation bistability. The Spo0E phosphatase can dephosphorylate Spo0A-P (Ohlsen *et al.*, 1994). Overproduction of Spo0E was shown to inhibit sporulation, whereas deletion of this locus resulted in premature sporulation (Perego and Hoch, 1991). A *spo0E* mutation was introduced in our reporter strain. The resulting strain, IIA-gfp/ $\Delta spo0E$ (P_{spoIIA} -gfp, $\Delta spo0E$), was grown in SM and cells analysed by flow cytometry. As shown in Fig. 8, the final bistable expression of *spoIIA* is completely abolished in a *spo0E* mutant background, and the vast majority of cells express high levels of *spoIIA*. Although activation of the *spoIIA* operon is premature in the *spo0E* mutant, the initial expression distribution is still bistable (data not shown). This result suggests that by modulating phosphorelay activity through external phosphatases, the kinetics of the autostimulatory Spo0A acti-

vation can be adjusted, resulting in different bistable distributions.

To substantiate this supposition, we tested the effect of the artificial induction of a heterologous RapA homologue, Rap60. The Rap60 phosphatase is encoded on pTA1060, a rolling-circle plasmid of *Bacillus amyloliquefaciens* (Meijer *et al.*, 1998). Previously, it was shown that Rap60, which is highly homologous to RapA, could dephosphorylate a component of the phosphorelay of *B. subtilis* (Koetje *et al.*, 2003). The *rap60* gene was placed under a xylose-inducible promoter and introduced into the *amyE* locus of a strain in which either *rapA* or *spo0E* was deleted. Strains IIA/0E/Xrap ($P_{spoIIA}\text{-gfp}$, $\Delta spo0E$, *amyE::P_{xyI}-rap60*) and IIA/rapA/Xrap ($P_{spoIIA}\text{-gfp}$, $\Delta rapA$, *amyE::P_{xyI}-rap60*) were grown in SM and xylose was added 1 h before entry into the stationary growth phase. Cultures were analysed using flow cytometry two hours after entry into the stationary growth phase (Fig. 9). As shown in Fig. 9, xylose induction of Rap60 complements the abolished bistable expression pattern of a *rapA* and *spo0E* mutant. These results show that the phosphorelay can be used as a tuner to modulate the bistable outcome of the sporulating culture.

Discussion

The *spoIIA* operon, encoding the sporulation sigma factor σ^F , is one of the earliest targets that is activated by Spo0A~P at the onset of sporulation (Sonenshein, 2000). Using GFP as a reporter, activation of this operon was shown to be heterogeneously distributed within the sporulating culture (Fig. 2). By following an approach pioneered by Elowitz and coworkers, we showed that the heterogeneity in activation of the *spoIIA* promoter is not a consequence of transcriptional noise of this promoter (Fig. 3). Elowitz *et al.* have shown that low intrinsic noise of target promoters can be observed when intracellular concentrations of activator proteins are high (Elowitz *et al.*, 2002). This suggests that the *spoIIA* promoter requires relatively high amounts of regulator protein(s), such as Spo0A, to activate its transcription. In support of this, we demonstrate that cells that have a high *spo0A* expression state are the ones that also activate expression of the *spoIIA* operon (Fig. 4). These results agree with previous findings (Bird *et al.*, 1992; Chung *et al.*, 1994). The *spoIIA* promoter is therefore a useful reporter to study the heterogeneous activation of Spo0A in a sporulating culture.

Modelling of autostimulatory gene regulation networks has shown that stochastic fluctuations of regulatory components in these systems can lead to a bistable response (Hasty *et al.*, 2000). This was biologically supported using relative simple well-defined, positive feedback circuits as models, such as that derived from bacteriophage lambda in *E. coli* (Hasty *et al.*, 2000; Isaacs *et al.*, 2003), and a

Fig. 9. A RapA homologue can modulate sporulation bistability. Strains were grown in sporulation medium and cultures were split in equal volumes and xylose was added to the indicated concentration 1 h before entry into the stationary growth phase. Cultures were analysed by flow cytometry two hours after entry into the stationary growth phase. As a control for a possible xylose effect on sporulation heterogeneity, strains IIA-gfp and IIA-gfp/ $\Delta spo0E$ were also analysed and showed no decrease in expression of *spoIIA*, rather a slight increase (data not shown). Fluorescence distributions of strain IIA/rapA/Xrap ($P_{spoIIA}\text{-gfp}$, $\Delta rapA$, *amyE::P_{xyI}-rap60*) (A) and IIA/0E/Xrap ($P_{spoIIA}\text{-gfp}$, $\Delta spo0E$, *amyE::P_{xyI}-rap60*) (B) grown in the presence of the indicated xylose concentration.

tetracycline-responsive transactivator system in yeast (Becskei *et al.*, 2001). Although sporulation is initiated by the autostimulatory expression of *spo0A*, we cannot speak of a simple positive feedback circuit. As indicated in Fig. 1, and described in the introduction, there are several positive feedback loops working on the *spo0A* promoter. The activation of Spo0A by the phosphorelay is also subject to autostimulatory influences (Strauch *et al.*, 1992; Strauch *et al.*, 1993; Fujita and Sadaie, 1998). By using an inducible and constitutively active variant of Spo0A, we demonstrated that sporulation bistability is abolished when the Spo0A-autostimulation is replaced by a graded induction (Fig. 5). This corresponds with previously described regulatory systems in which autostimula-

tion of a regulator can convert a graded response to a bistable response (Gardner *et al.*, 2000; Hofer *et al.*, 2002). In conclusion, these results indicate that the complex autostimulatory Spo0A regulation cascade can be considered as a classical autostimulatory loop with a bistable output. The mechanism that decides which cell reaches the Spo0A threshold first, is likely based on a random process (Dawes and Thornley, 1970; Hasty *et al.*, 2000; Maughan and Nicholson, 2004).

Notably, during the revision of this manuscript, Fujita and coworkers have also shown, using different methodologies and independent of our work, that initiation of sporulation requires a threshold level of Spo0A and can be regarded as a bistable switch, results consistent with the work presented here (Fujita *et al.*, 2005).

Heterogeneity in gene expression in *B. subtilis* cultures is not only limited to the process of sporulation but also takes place during competence development (Cahn and Fox, 1968; Hadden and Nester, 1968). Because this process is also influenced by Spo0A~P (Hahn *et al.*, 1995), it raises the question whether heterogeneity observed in this process is the consequence of *spo0A* autoactivation. However, in our group we have recently obtained evidence that the autostimulatory transcription loop of *comK* (encoding the Competence Transcription Factor), is responsible for bistability in competence development (Smits *et al.*, 2005).

lacZ reporter studies has shown that *rapA* expression coincides with the initiation of sporulation, which is peculiar considering the negative effect of RapA on this process. We showed that this discrepancy is resolved when looking at expression levels in individual cells. As a result of the bistable nature of sporulation initiation, some cells initiate sporulation and repress *rapA* gene expression while the non-sporulating cells continue to accumulate RapA (Fig. 6). When *rapA* was deleted, the final bistable distribution of sporulation gene expression was almost abolished (Fig. 7). Based on these findings, we propose that the main function of RapA is to maintain the bistable gene expression that originates from the autostimulatory *spo0A* activation.

The fascinating aspect of *rapA* regulation is that the induction is subjected to multiple quorum sensing pathways. The initial *rapA* expression is not heterogeneous, which suggests that induction of all ComA-regulated genes is homogeneous (Fig. 6). This corresponds with the observation that *srfA* induction, directly activated by the ComA/P/X system, also showed homogeneous expression (Hahn *et al.*, 1994). One could imagine that within shaken liquid cultures, quorum sensors, such as ComX, are evenly distributed. Quorum sensing might be a mechanism to reduce noise in signal transduction pathways: although cells might differ in the expression of regulatory systems, they will sense the same level of pheromones.

Of course this situation will be different in dense biofilm populations.

The phosphatase activity of RapA is modulated by PhrA, a small-secreted peptide encoded by an adjacent gene on the same transcript as *rapA*. Previously, it was shown that sporulation is poor in a *phrA* mutant strain (Perego and Hoch, 1996). Eventually, the accumulation of PhrA seems to antagonize the dephosphorylating action of RapA, and thus allows delayed cells to initiate sporulation. Perego suggested that the RapA-PhrA system is not a cell-density control mechanism but that processed PhrA primarily accumulates in the periplasmic space (Perego, 1999). This was based on the observation that: (i) free PhrA peptide could only be detected in medium of a *opp* mutant strain (owing to a lack of re-import) and (ii) the peptide is susceptible to the high amounts of extracellular proteases produced by *B. subtilis*. Because of the temporal sequence of export, processing and import, the PhrA pheromone might provide a proscribed and finite time frame for information processing and decision-making meaning that the RapA-PhrA system functions as a sporulation timing device (Perego, 1999). In this work, we show that only a limited number of cells express high levels of *rapA* and are delayed in sporulation. As a result of cotranscription, these cells are also likely to express high levels of *phrA*. It is envisioned that in time, RapA inhibition can be counteracted by the continuous accumulation of PhrA. Whether this would occur *in cis* (PhrA in periplasm) or *in trans* (PhrA in medium) remains to be established. However, Perego's idea that the RapA-PhrA functions as a sporulation-timing device corresponds with our postulated role as stabilizer of bistable gene expression in sporulating cultures.

Surprisingly, deletion of *comA*, the activator of *rapA*, did not promote sporulation under the conditions tested (J.-W. Veening, L. W. Hamoen and O. P. Kuipers, unpubl. results), whereas overexpression of ComA was previously shown to inhibit sporulation (Weinrauch *et al.*, 1989). This suggests that the ComA-ComP system, besides activating expression of negative regulators like *rapA*, also regulates genes that promote sporulation initiation. A possible candidate is the *srfA* operon, which is activated by ComA~P (Nakano *et al.*, 1991a). The induction of *srfA* is also important for the development of genetic competence. In the 5' end of this operon a small gene, *comS*, is embedded which is essential for the activation of ComK. Interestingly, the 3' part of this operon was found to be required for sporulation (Nakano *et al.*, 1991b). Whether there is another (small) gene present in this operon that is required for sporulation remains to be investigated.

Another mechanism that could act as a sporulation delay system was previously described by Gonzalez-Passtor *et al.* (2003). They showed that, within a sporulating culture, sporulating cells are able to kill their siblings that

have not yet initiated this process. This phenomenon is responsible for the reduction in cell density of a sporulating culture. The operon responsible for producing the sporulation killing factor (*skf*) is activated by Spo0A~P. The Spo0A-inactive subpopulation is sensitive to the sporulation-killing factor secreted by the Spo0A-active cells, and will subsequently lyse and release nutrients available for the remainder of the population. Early in the stationary growth phase, the low Spo0A-active population (Fig. 2 left peak) seems to be resistant to the killing factor. This is most likely caused by the high efficiency of the coexpressed export pump proteins SkfE and SkfF, which confer resistance against the killing factor (J.E. Gonzalez-Pastor, pers. comm.). The Skf system could provide the sporulating culture with an alternative mechanism to maintain sporulation bistability by impeding cell growth via the Skf antibiotic. However, a disruption of the *skf* operon did not change the bistable expression of *spoIIA* (data not shown). Using a dye that specifically stains cells with damaged membranes (propidium iodide), no dead cells among the high RapA-active (and low Spo0A-active) subpopulation were detected (data not shown). Thus, the RapA-active subpopulation is not targeted for lysis, but is able to commit to other differentiation pathways than sporulation.

In conclusion, the RapA results show that phosphorelay phosphatases have a modulating action on the bistable expression pattern of a sporulating culture by altering the threshold-level of Spo0A~P. By monitoring *spoIIA* expression in a *spo0E* mutant, and in strains containing artificially inducible Rap60, we substantiated that the phosphorelay can be used as a tuner to modulate the

bistable outcome of the sporulating culture (Figs 8 and 9). The exact function of Spo0E in sporulation regulation is unclear. AbrB represses *spo0E* gene expression (Strauch *et al.*, 1989; Perego and Hoch, 1991). Apparently, cells that initiate *spo0A* autoactivation also induce Spo0E levels. This coexpression suggests that Spo0E serves as some kind of a 'safety lid' to prevent over-stimulation of *spo0A* autoactivation. Rap phosphatases are common among *Bacillus* species and many of them are encoded on plasmids isolated from industrially relevant strains. These strains are used for high production of enzymes, and generally show a reduction in sporulation (Meijer *et al.*, 1998). Apparently, natural selection chose to alter the sporulation outcome by using external phosphatases to modulate the phosphorelay.

Experimental procedures

Plasmids, bacterial strains and media

Table 2 lists the plasmids and bacterial strains used. TY medium (tryptone/yeast extract) contained Bacto-Tryptone (1%), Bacto-yeast extract (0.5%) and NaCl (1%). SM contained dehydrated nutrient broth (0.8%), NaOH (0.5 mM), MgSO₄ (1 mM), KCl (1 g l⁻¹), Ca(NO₃)₂ 1 mM, MnCl₂ (0.01 mM) and FeSO₄ (0.001 mM). Minimal medium for *B. subtilis* was prepared as described before (Veening *et al.*, 2004). When required, medium for *E. coli* was supplemented with ampicillin (Ap; 100 µg ml⁻¹); media for *B. subtilis* were supplemented with chloramphenicol (Cm; 5 µg ml⁻¹), tetracycline (Tc; 6 µg ml⁻¹), spectinomycin (Sp; 100 µg ml⁻¹) or kanamycin (Km; 10 µg ml⁻¹). When indicated, IPTG (isopropyl-β-D-thiogalactosidase) or xylose was added to the medium.

Table 2. Bacterial strains and plasmids.

	Genotype	Reference
Strains		
<i>E. coli</i>		
MC1061	F ⁻ , <i>araD</i> 139, Δ(<i>ara-leu</i>)7696, Δ(<i>lac</i>)X74, <i>galU</i> , <i>galK</i> , <i>hsdR2</i> , <i>mcrA</i> , <i>mcrB</i> 1, <i>rspL</i>	(Wertman <i>et al.</i> , 1986)
<i>B. subtilis</i>		
168	<i>trpC</i> 2	(Kunst <i>et al.</i> , 1997)
IIA-gfp	168, P _{<i>spoIIA</i>} -gfp, Cm ^r	This study
IIA/double	168, P _{<i>spoIIA</i>} -lyfp, Cm ^r , amyE::P _{<i>spoIIA</i>} -icfp, Km ^r	This study
spo0A/IIA	168, P _{<i>spo0A</i>} -lyfp, Cm ^r , amyE::P _{<i>spoIIA</i>} -icfp, Km ^r	This study
rapA/IIA	168, P _{<i>rapA</i>} -lyfp, Cm ^r , amyE::P _{<i>spoIIA</i>} -icfp, Km ^r	This study
rapA-lyfp	168, P _{<i>rapA</i>} -lyfp, Cm ^r	This study
IIA-gfp/ΔrapA	168, P _{<i>spoIIA</i>} -gfp, Cm ^r , rapA::Sp, Sp ^r	This study
IIA-gfp/Δspo0E	168, P _{<i>spoIIA</i>} -gfp, Cm ^r , spo0E::Sp, Sp ^r	This study
IIA/spo0A +	168, P _{<i>spoIIA</i>} -gfp, Sp ^r , amyE::P _{<i>spac</i>} -spo0A, Cm ^r	This study
IIA/sad67	168, P _{<i>spoIIA</i>} -gfp, Cm ^r , amyE::P _{<i>spac</i>} -sad67, Tc ^r	This study
IIA/spo0A/sad67	168, P _{<i>spoIIA</i>} -gfp, Cm ^r , spo0A::Km, Km ^r , amyE::P _{<i>spac</i>} -sad67, Tc ^r	This study
IIA/0E/Xrap	168, P _{<i>spoIIA</i>} -gfp, Cm ^r , spo0E::Sp, Sp ^r , amyE::P _{<i>xyIA</i>} -rap60, Km ^r	This study
IIA/rapA/Xrap	168, P _{<i>spoIIA</i>} -gfp, Cm ^r , rapA::Sp, Sp ^r , amyE::P _{<i>xyIA</i>} -rap60, Km ^r	This study
Plasmids		
pGFP-IIA	<i>bla</i> , cat, P _{<i>spoIIA</i>} -gfp	This study
pLYFP-spo0A	<i>bla</i> , cat, P _{<i>spo0A</i>} -lyfp	This study
pLYFP-rapA	<i>bla</i> , cat, P _{<i>rapA</i>} -lyfp	This study

Recombinant DNA techniques and oligonucleotides

Procedures for DNA purification, restriction, ligation, agarose gel electrophoresis and transformation of *E. coli* were carried out as described before (Sambrook *et al.*, 1989). Enzymes were obtained from Roche (Mannheim, GER). Oligonucleotides were purchased from Biolegio BV (Malden, NL). *B. subtilis* was transformed as described before (Veening *et al.*, 2004).

Construction of plasmids

To construct plasmid pGFP-IIA, carrying the *B. subtilis* *spolIIA* promoter region fused with the *gfpmut1* gene, a polymerase chain reaction (PCR) with the primers *spolIIA*-F (5'-CCCAAGCTTAGGCCAAGAGCTTGGCACT-3') and *spolIIA*-R (5'-TTCTGCAGGCTCATGCTCATTCCTCCTTG-3') was performed, using chromosomal DNA of *B. subtilis* 168 as a template. The amplified fragment was subsequently cleaved with *HindIII* and *PstI*, and ligated into the corresponding sites of pSG1151 (Lewis and Marston, 1999), resulting in plasmid pGFP-IIA.

To construct plasmid pIYFP-spo0A, carrying the *B. subtilis* *spo0A* promoter region fused with the *iyfp* gene, a PCR with the primers *spo0A*-F (5'-CCCAAGCTTCGTACAGGGGATGAGCGGAAGCCC-3') and *spo0A*-R (5'-CGGAATTCCTCACGTTTCTTCTCCCAATG-3') was performed, using chromosomal DNA of *B. subtilis* 168 as a template. The amplified fragment was subsequently cleaved with *HindIII* and *EcoRI*, and ligated into the corresponding sites of pIYFP (Veening *et al.*, 2004), resulting in plasmid pIYFP-spo0A.

To construct plasmid pIYFP-rapA, carrying the *B. subtilis* *rapA* promoter region fused with the *iyfp* gene, a PCR with the primers *rapA*-F (5'-CCCAAGCTTCGCTCCTGCAGTCTGGTCATCGAGG-3') and *rapA*-R (5'-CGGAATTCCTGCTTCATCCTCAATTAATCCCC-3') was performed, using chromosomal DNA of *B. subtilis* 168 as a template. The amplified fragment was subsequently cleaved with *HindIII* and *EcoRI*, and ligated into the corresponding sites of pIYFP (Veening *et al.*, 2004), resulting in plasmid pIYFP-rapA.

Construction of bacterial strains

Bacillus subtilis strain IIA-gfp was obtained by a Campbell-type integration (single crossover) of plasmid pGFP-IIA into the chromosomal *spolIIA* promoter region of *B. subtilis* 168. *B. subtilis* strain rapA-iyfp was obtained by a Campbell-type integration of plasmid pIYFP-rapA into the chromosomal *rapA* promoter region of *B. subtilis* 168. Transformants were selected on TY agar plates containing Cm, after overnight incubation at 37°C. Correct integration was verified by PCR (data not shown).

Bacillus subtilis strain IIA/double was obtained by transformation of strain iyfp-IIA (Veening *et al.*, 2004) with chromosomal DNA of strain icfp-IIA-amyE (Veening *et al.*, 2004). Transformants were selected on TY agar plates containing Cm and Km, after overnight incubation at 37°C. Correct integration into the *amyE* gene was tested and confirmed by lack of amylase activity upon growth on plates containing 1% starch.

Bacillus subtilis strains spo0A/IIA and rapA/IIA were obtained by a Campbell-type integration of plasmids pIYFP-spo0A and pIYFP-rapA, respectively, into the chromosome of *B. subtilis* icfp-IIA-amyE (Veening *et al.*, 2004). Transformants were selected on TY agar plates containing Cm and Km, after overnight incubation at 37°C. Correct integration was verified by PCR (data not shown).

Bacillus subtilis strain IIA-gfp/ Δ spo0E ($P_{spolIIA}$ -gfp, Δ spo0E) was obtained by transformation of strain IIA-gfp with chromosomal DNA of strain 0ES (Hosoya *et al.*, 2002). Transformants were selected on TY agar plates containing Cm and Sp, after overnight incubation at 37°C. *B. subtilis* strain Δ rapA-Sp was obtained by transformation of strain Δ rapA-Cm (Tjalsma *et al.*, 2004) with plasmid pCm::Sp (Steinmetz and Richter, 1994). Transformants were selected on TY agar plates containing Sp, after overnight incubation at 37°C. To confirm correct switching of the Cm resistance marker to the Sp resistance marker, transformants were scored on Cm sensitivity and Sp resistance. *B. subtilis* strain IIA-gfp/ Δ rapA ($P_{spolIIA}$ -gfp, Δ rapA) was obtained by transformation of strain IIA-gfp with chromosomal DNA of strain Δ rapA-Sp. Transformants were selected on TY agar plates containing Cm and Sp, after overnight incubation at 37°C.

Bacillus subtilis strain IIA-gfp::Sp was obtained by transformation of strain IIA-gfp with plasmid pCm::Sp (Steinmetz and Richter, 1994). Transformants were selected on TY agar plates containing Sp, after overnight incubation at 37°C. To confirm correct switching of the Cm resistance marker to the Sp resistance marker, transformants were scored on Cm sensitivity and Sp resistance. *B. subtilis* strain IIA/spo0A + ($P_{spolIIA}$ -gfp, P_{spac} -spo0A) was obtained by transformation of strain IIA-gfp::Sp with chromosomal DNA of strain Sik243 (Ireton *et al.*, 1993). Transformants were selected on TY agar plates containing Sp and Cm, after overnight incubation at 37°C.

Bacillus subtilis strain Xrap::Neo was obtained by transformation of strain R (Koetje *et al.*, 2003) with plasmid pCm::Nm (Steinmetz and Richter, 1994). Transformants were selected on TY agar plates containing Km, after overnight incubation at 37°C. To confirm correct switching of the Cm resistance marker to the Km resistance marker, transformants were scored on Cm sensitivity and Km resistance. *B. subtilis* strain IIA-gfp, Xrap was obtained by transformation of strain IIA-gfp with chromosomal DNA of strain Xrap::Neo. Transformants were selected on TY agar plates containing Cm and Km, after overnight incubation at 37°C. Strains IIA/0E/Xrap ($P_{spolIIA}$ -gfp, Δ spo0E, *amyE*:: P_{xyIA} -rap60) and IIA/rapA/Xrap ($P_{spolIIA}$ -gfp, Δ rapA, *amyE*:: P_{xyIA} -rap60) were obtained by transformation of strain IIA-gfp, Xrap with chromosomal DNA of strains 0ES (Hosoya *et al.*, 2002) and Δ rapA-Sp respectively. Transformants were selected on TY agar plates containing Cm, Km and Sp, after overnight incubation at 37°C.

Bacillus subtilis strain sad67::Cm (P_{spac} -spo0A-sad67) was obtained by transformation of strain 168 with chromosomal DNA of strain Sik31 (Ireton *et al.*, 1993). Transformants were selected on TY agar plates containing Cm, after overnight incubation at 37°C. To obtain *B. subtilis* strain sad67::Tc, in which the Cm resistance marker of strain sad67::Cm is switched to a Tc resistance marker, strain sad67::Cm was transformed with linearized plasmid pCm::Tc (Steinmetz and Richter, 1994). Correct transformants were scored on Cm

sensitivity and Tc resistance. To obtain *B. subtilis* strain IIA-gfp/sad67, strain sad67::Tc was transformed with chromosomal DNA of strain IIA-gfp. Transformants were selected on TY agar plates containing Tc and Cm, after overnight incubation at 37°C. Finally, *B. subtilis* strain IIA/spo0A/sad67 was obtained by transforming strain IIA-gfp/sad67 with chromosomal DNA of strain SWV215 (*spo0A*::Km) (Xu and Strauch, 1996). Transformants were selected on TY agar plates containing Tc, Cm and Km, after overnight incubation at 37°C.

Microscopy

Cells were prepared for microscopy and applied to agarose slides as described before (Veening *et al.*, 2004), and images were acquired using an Axiophot microscope equipped with an AxioVision camera (Zeiss, Oberkochen, GER). Fluorescence filter sets used to visualize the green, cyan and yellow fluorescent proteins were obtained from Zeiss. Fluorescent signals of GFP were visualized using set 09 (excitation, 450–490 nm; emission, > 520 nm), fluorescent ICFP signals were visualized using set 47 (excitation, 426–446 nm; emission, 460–500 nm) and fluorescent signals of IYFP using set 46 (excitation, 490–510 nm; emission, 520–550 nm). AxioVs20 software (Zeiss) was used for image capturing and figures were prepared for publication using Corel Graphics Suite 11 (Corel Corporation).

Flow cytometry

Cells were 100× diluted in 0.2 µM filtered minimal medium and directly measured on a Coulter Epics XL-MCL flow cytometer (Beckman Coulter, Mijdrecht, NL) operating an argon laser (488 nm). For each sample, at least 20 000 cells were analysed. Data containing the green fluorescent signals were collected by a FITC filter and the photomultiplier voltage was set between 700 and 800 V. Data were captured using EXPO32 software (Beckman Coulter) and further analysed using WinMDI 2.8 software (<http://facs.scripps.edu/software.html>). Figures were prepared for publication using WinMDI 2.8 and Corel Graphics Suite 11. To distinguish background fluorescence from GFP specific fluorescence, parental strain *B. subtilis* 168 was also analysed with each flow cytometric experiment.

Calculation of noise

Strain IIA/double was grown in SM and cells were collected for fluorescent microscopy two hours after entry into stationary growth phase. Images were analysed using Imagemaster 2D elite software v3.1 (Amersham Pharmacia biotech) and Microsoft Excel (as described by Smits *et al.*, 2005). Data were calculated essentially as described before (Elowitz *et al.*, 2002). In brief, intrinsic noise (η_{int}), extrinsic noise (η_{ext}) and total variation (η_{tot}) were defined as follows:

$$\eta_{\text{int}}^2 = \frac{\langle (c - \bar{c})^2 \rangle}{2\langle c \rangle \langle y \rangle}; \quad \eta_{\text{ext}}^2 = \frac{\langle cy \rangle - \langle c \rangle \langle y \rangle}{\langle c \rangle \langle y \rangle};$$

$$\eta_{\text{tot}}^2 = \frac{\langle c^2 + y^2 \rangle - 2\langle c \rangle \langle y \rangle}{2\langle c \rangle \langle y \rangle}$$

Here the i^{th} element of vectors c and y contain the average CFP or YFP intensity, respectively, of the i^{th} cell in the sample. Angled brackets denote means over the cell population.

Western blot analysis and immunodetection

Before harvesting of the cells, the optical density at 600 nm was measured and was used to correct for the total amount of protein loaded on the gel. Cells were separated from the growth medium by centrifugation (14 000 r.p.m., 1 min, room temperature). Pelleted cells were resuspended in protoplast buffer (20 mM potassium phosphate pH 7.5, 15 mM MgCl₂, 20% sucrose and 1 mg ml⁻¹ of lysozyme) and incubated at 37°C for 30 min. The resulting protoplasts were diluted with 2× SDS sample buffer, incubated at 95°C for 5 min and separated by SDS-polyacrylamide gel electrophoresis (SDS-PAGE) as described (Sambrook *et al.*, 1989). Next, proteins were transferred to a polyvinylidene difluoride (PVDF) membrane (Roche) as described (Sambrook *et al.*, 1989). Spo0A and Spo0A-Sad67 were detected with polyclonal anti-Spo0A antibodies (kindly provided by M. Fujita), GroEL was detected with polyclonal anti-GroEL antibodies (our laboratory collection) and horseradish peroxidase-antirabbit-IgG conjugate (Amersham Biosciences, Little Chalfont, UK). Anti-Spo0A and anti-GroEL was added in a 1:5000 dilution and anti-rabbit at a 1:7500 dilution. Anti-Spo0A antibodies can be used to detect both wild-type Spo0A and the N-terminal deletion mutant Spo0A-Sad67 (Fujita and Losick, 2003).

Acknowledgements

We thank W.H. van de Poll for assistance with the flowcytometer. We thank M. Fujita and I. Barak for kindly providing us with antibodies against Spo0A. We thank the members of our group and two anonymous referees for helpful advice and we thank I. Barak, D. Dubnau, R. Losick, M. Fujita and J.E. Gonzalez-Pastor for personal communications. J.W.V. was supported by Grant ABC-5587 from NWO-STW and L.W.H. was supported by a BBSRC Grant awarded to J. Errington.

Supplementary material

The following supplementary material is available for this article online:

Appendix S1. Specific growth conditions used in this study.

Fig. S1. Typical growth of *B. subtilis* IIA-gfp grown in sporulation medium.

Fig. S2. Chromosome map of strain IIA/double.

References

- Balaban, N.Q., Merrin, J., Chait, R., Kowalik, L., and Leibler, S. (2004) Bacterial persistence as a phenotypic switch. *Science* **305**: 1622–1625.
- Becskei, A., Seraphin, B., and Serrano, L. (2001) Positive feedback in eukaryotic gene networks: cell differentiation

- by graded to binary response conversion. *EMBO J* **20**: 2528–2535.
- Bird, T., Burbulys, D., Wu, J.J., Strauch, M.A., Hoch, J.A., and Spiegelman, G.B. (1992) The effect of supercoiling on the in vitro transcription of the *spoIIA* operon from *Bacillus subtilis*. *Biochimie* **74**: 627–634.
- Blake, W.J., Kaern, M., Cantor, C.R., and Collins, J.J. (2003) Noise in eukaryotic gene expression. *Nature* **422**: 633–637.
- Burbulys, D., Trach, K.A., and Hoch, A. (1991) Initiation of sporulation in *B. subtilis* is controlled by a multicomponent phosphorelay. *Cell* **64**: 545–552.
- Cahn, F.H., and Fox, M.S. (1968) Fractionation of transformable bacteria from competent cultures of *Bacillus subtilis* on renografin gradients. *J Bacteriol* **95**: 867–875.
- Chung, J.D., Stephanopoulos, G., Ireton, K., and Grossman, A.D. (1994) Gene expression in single cells of *Bacillus subtilis*: evidence that a threshold mechanism controls the initiation of sporulation. *J Bacteriol* **176**: 1977–1984.
- Dawes, I.W., and Thornley, J.H. (1970) Sporulation in *Bacillus subtilis*. Theoretical and experimental studies in continuous culture systems. *J Gen Microbiol* **62**: 49–66.
- Elowitz, M.B., Levine, A.J., Siggia, E.D., and Swain, P.S. (2002) Stochastic gene expression in a single cell. *Science* **297**: 1183–1186.
- Ferrell, J.E., Jr. (2002) Self-perpetuating states in signal transduction: positive feedback, double-negative feedback and bistability. *Curr Opin Cell Biol* **14**: 140–148.
- Fujita, M., and Losick, R. (2003) The master regulator for entry into sporulation in *Bacillus subtilis* becomes a cell-specific transcription factor after asymmetric division. *Genes Dev* **17**: 1166–1174.
- Fujita, M., and Sadaie, Y. (1998) Feedback loops involving Spo0A and AbrB in *in vitro* transcription of the genes involved in the initiation of sporulation in *Bacillus subtilis*. *J Biochem (Tokyo)* **124**: 98–104.
- Fujita, M., Gonzalez-Pastor, J.E., and Losick, R. (2005) High- and low-threshold genes in the Spo0A regulon of *Bacillus subtilis*. *J Bacteriol* **187**: 1357–1368.
- Gardner, T.S., Cantor, C.R., and Collins, J.J. (2000) Construction of a genetic toggle switch in *Escherichia coli*. *Nature* **403**: 339–342.
- Gonzalez-Pastor, J.E., Hobbs, E.C., and Losick, R. (2003) Cannibalism by sporulating bacteria. *Science* **301**: 510–513.
- Grossman, A.D. (1995) Genetic networks controlling the initiation of sporulation and the development of genetic competence in *Bacillus subtilis*. *Annu Rev Genet* **29**: 477–508.
- Hadden, C., and Nester, E.W. (1968) Purification of competent cells in the *Bacillus subtilis* transformation system. *J Bacteriol* **95**: 876–885.
- Hahn, J., Kong, L., and Dubnau, D. (1994) The regulation of competence transcription factor synthesis constitutes a critical control point in the regulation of competence in *Bacillus subtilis*. *J Bacteriol* **176**: 5753–5761.
- Hahn, J., Roggiani, M., and Dubnau, D. (1995) The major role of Spo0A in genetic competence is to downregulate *abrB*, an essential competence gene. *J Bacteriol* **177**: 3601–3605.
- Hasty, J., Pradines, J., Dolnik, M., and Collins, J.J. (2000) Noise-based switches and amplifiers for gene expression. *Proc Natl Acad Sci USA* **97**: 2075–2080.
- Hoch, J.A. (1991) *Spo0* genes, the phosphorelay, and the initiation of sporulation. In *Bacillus Subtilis and Other Gram-Positive Bacteria: Biochemistry, Physiology, and Molecular Genetics*. Sonenshein, A.L., Hoch, J.A., and Losick, R. (eds). Washington, DC: American Society for Microbiology, pp. 747–755.
- Hofer, T., Nathansen, H., Lohning, M., Radbruch, A., and Heinrich, R. (2002) GATA-3 transcriptional imprinting in Th2 lymphocytes: a mathematical model. *Proc Natl Acad Sci USA* **99**: 9364–9368.
- Hosoya, S., Asai, K., Ogasawara, N., Takeuchi, M., and Sato, T. (2002) Mutation in *yaaT* leads to significant inhibition of phosphorelay during sporulation in *Bacillus subtilis*. *J Bacteriol* **184**: 5545–5553.
- Ireton, K., Rudner, D.Z., Siranosian, K.J., and Grossman, A.D. (1993) Integration of multiple developmental signals in *Bacillus subtilis* through the Spo0A transcription factor. *Genes Dev* **7**: 283–294.
- Isaacs, F.J., Hasty, J., Cantor, C.R., and Collins, J.J. (2003) Prediction and measurement of an autoregulatory genetic module. *Proc Natl Acad Sci USA* **100**: 7714–7719.
- Koetje, E.J., Hajdo-Milasnovic, A., Kiewiet, R., Bron, S., and Tjalsma, H. (2003) A plasmid-borne Rap-Phr system of *Bacillus subtilis* can mediate cell-density controlled production of extracellular proteases. *Microbiology* **149**: 19–28.
- Kunst, F., Ogasawara, N., Moszer, I., Albertini, A.M., Alloni, G., Azevedo, V., *et al.* (1997) The complete genome sequence of the Gram-positive bacterium *Bacillus subtilis*. *Nature* **390**: 249–256.
- Lazazzera, B.A., Palmer, T., Quisel, J., and Grossman, A.D. (1999) Cell density control of gene expression and development in *Bacillus subtilis*. In *Cell-cell Signaling in Bacteria*. Dunny, G.M., and Winans, S.C. (eds). Washington, DC: American Society for Microbiology, pp. 27–46.
- Lewis, P.J., and Marston, A.L. (1999) GFP vectors for controlled expression and dual labelling of protein fusions in *Bacillus subtilis*. *Gene* **227**: 101–110.
- McAdams, H.H., and Arkin, A. (1997) Stochastic mechanisms in gene expression. *Proc Natl Acad Sci USA* **94**: 814–819.
- Maughan, H., and Nicholson, W.L. (2004) Stochastic processes influence stationary-phase decisions in *Bacillus subtilis*. *J Bacteriol* **186**: 2212–2214.
- Meijer, W.J., Wisman, G.B., Terpstra, P., Thorsted, P.B., Thomas, C.M., Holsappel, S., *et al.* (1998) Rolling-circle plasmids from *Bacillus subtilis*: complete nucleotide sequences and analyses of genes of pTA1015, pTA1040, pTA1050 and pTA1060, and comparisons with related plasmids from Gram-positive bacteria. *FEMS Microbiol Rev* **21**: 337–368.
- Molle, V., Fujita, M., Jensen, S.T., Eichenberger, P., Gonzalez-Pastor, J.E., Liu, J.S., and Losick, R. (2003) The Spo0A regulon of *Bacillus subtilis*. *Mol Microbiol* **50**: 1683–1701.
- Mueller, J.P., and Sonenshein, A.L. (1992) Role of the *Bacillus subtilis* *gsiA* gene in regulation of early sporulation gene expression. *J Bacteriol* **174**: 4374–4383.
- Mueller, J.P., Bukusoglu, G., and Sonenshein, A.L. (1992) Transcriptional regulation of *Bacillus subtilis* glucose starvation-inducible genes: control of *gsiA* by the ComP-

- ComA signal transduction system. *J Bacteriol* **174**: 4361–4373.
- Nakano, M.M., Xia, L.A., and Zuber, P. (1991a) Transcription initiation region of the *srfA* operon, which is controlled by the comP-comA signal transduction system in *Bacillus subtilis*. *J Bacteriol* **173**: 5487–5493.
- Nakano, M.M., Magnuson, R., Myers, A., Curry, J., Grossman, A.D., and Zuber, P. (1991b) *srfA* is an operon required for surfactin production, competence development, and efficient sporulation in *Bacillus subtilis*. *J Bacteriol* **173**: 1770–1778.
- Ohlsen, K.L., Grimsley, J.K., and Hoch, J.A. (1994) Deactivation of the sporulation transcription factor Spo0A by the Spo0E protein phosphatase. *Proc Natl Acad Sci USA* **91**: 1756–1760.
- Ozbudak, E.M., Thattai, M., Kurtser, I., Grossman, A.D., and van Oudenaarden, A. (2002) Regulation of noise in the expression of a single gene. *Nat Genet* **31**: 69–73.
- Parker, G.F., Daniel, R.A., and Errington, J. (1996) Timing and genetic regulation of commitment to sporulation in *Bacillus subtilis*. *Microbiology* **142**: 3445–3452.
- Perego, M. (1999) Self-signaling by Phr peptides modulates *Bacillus subtilis* development. In *Cell-cell Signaling in Bacteria*. Dunny, G.M., and Winans, S.C. (eds). Washington, DC: American Society for Microbiology, pp. 243–258.
- Perego, M., and Hoch, J.A. (1991) Negative regulation of *Bacillus subtilis* sporulation by the spo0E gene product. *J Bacteriol* **173**: 2514–2520.
- Perego, M., and Hoch, J.A. (1996) Cell-cell communication regulates the effects of protein aspartate phosphatases on the phosphorelay controlling development in *Bacillus subtilis*. *Proc Natl Acad Sci USA* **93**: 1549–1553.
- Perego, M., and Hoch, J.A. (2002) Two-component systems, phosphorelays, and regulation of their activities by phosphatases. In *Bacillus subtilis and its Closest Relatives: from Genes to Cells*. Sonenshein, A.L., Hoch, J.A., and Losick, R. (eds). Washington, DC: American Society for Microbiology Press, pp. 473–481.
- Predich, M., Nair, G., and Smith, I. (1992) *Bacillus subtilis* early sporulation genes *kinA*, *spo0F*, and *spo0A* are transcribed by the RNA polymerase containing sigma H. *J Bacteriol* **174**: 2771–2778.
- Rao, C.V., Wolf, D.M., and Arkin, A.P. (2002) Control, exploitation and tolerance of intracellular noise. *Nature* **420**: 231–237.
- Raser, J.M., and O'Shea, E.K. (2004) Control of stochasticity in eukaryotic gene expression. *Science* **304**: 1811–1814.
- Robertson, J.B., Gocht, M., Marahiel, M.A., and Zuber, P. (1989) AbrB, a regulator of gene expression in *Bacillus*, interacts with the transcription initiation regions of a sporulation gene and an antibiotic biosynthesis gene. *Proc Natl Acad Sci USA* **86**: 8457–8461.
- Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989) *Molecular Cloning: a Laboratory Manual* Cold Spring Harbor. New York, NY: Cold Spring Harbor Laboratory Press.
- Smits, W.K., Eschevins, C.C., Susanna, K.A., Bron, S., Kuipers, O.P., and Hamoen, L.W. (2005) Stripping *Bacillus*: ComK auto-stimulation is responsible for the bistable response in competence development. *Mol Microbiol* **56**: 604–614.
- Sonenshein, A.L. (2000) Control of sporulation initiation in *Bacillus subtilis*. *Curr Opin Microbiol* **3**: 561–566.
- Steinmetz, M., and Richter, R. (1994) Plasmids designed to alter the antibiotic resistance expressed by insertion mutations in *Bacillus subtilis*, through *in vivo* recombination. *Gene* **142**: 79–83.
- Strauch, M.A. (1995) Delineation of AbrB-binding sites on the *Bacillus subtilis* *spo0H*, *kinB*, *ftsAZ*, and *pbpE* promoters and use of a derived homology to identify a previously unsuspected binding site in the *bsuB1* methylase promoter. *J Bacteriol* **177**: 6999–7002.
- Strauch, M.A., Spiegelman, G.B., Perego, M., Johnson, W.C., Burbulys, D., and Hoch, J.A. (1989) The transition state transcription regulator *abrB* of *Bacillus subtilis* is a DNA binding protein. *EMBO J* **8**: 1615–1621.
- Strauch, M.A., Trach, K.A., Day, J., and Hoch, J.A. (1992) Spo0A activates and represses its own synthesis by binding at its dual promoters. *Biochimie* **74**: 619–626.
- Strauch, M.A., Wu, J.J., Jonas, R.H., and Hoch, J.A. (1993) A positive feedback loop controls transcription of the *spo0F* gene, a component of the sporulation phosphorelay in *Bacillus subtilis*. *Mol Microbiol* **7**: 967–974.
- Tjalsma, H., Koetje, E.J., Kiewiet, R., Kuipers, O.P., Kolkman, M., van der Laan, J., et al. (2004) Engineering of quorum-sensing systems for improved production of alkaline protease by *Bacillus subtilis*. *J Appl Microbiol* **96**: 569–578.
- Veening, J.W., Smits, W.K., Hamoen, L.W., Jongbloed, J.D., and Kuipers, O.P. (2004) Visualization of Differential Gene Expression by Improved Cyan Fluorescent Protein and Yellow Fluorescent Protein Production in *Bacillus subtilis*. *Appl Environ Microbiol* **70**: 6809–6815.
- Weinrauch, Y., Guillen, N., and Dubnau, D.A. (1989) Sequence and transcription mapping of *Bacillus subtilis* competence genes *comB* and *comA*, one of which is related to a family of bacterial regulatory determinants. *J Bacteriol* **171**: 5362–5375.
- Wertman, K.F., Wyman, A.R., and Botstein, D. (1986) Host/vector interactions which affect the viability of recombinant phage lambda clones. *Gene* **49**: 253–262.
- Xu, K., and Strauch, M.A. (1996) Identification, sequence, and expression of the gene encoding gamma-glutamyl-transpeptidase in *Bacillus subtilis*. *J Bacteriol* **178**: 4319–4322.