

University of Groningen

## ABC transporter architecture and regulatory roles of accessory domains

Biemans-Oldehinkel, E; Doeven, MK; Poolman, B

*Published in:*  
 FEBS Letters

*DOI:*  
 [10.1016/j.febslet.2005.11.079](https://doi.org/10.1016/j.febslet.2005.11.079)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
 Publisher's PDF, also known as Version of record

*Publication date:*  
 2006

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*  
 Biemans-Oldehinkel, E., Doeven, MK., & Poolman, B. (2006). ABC transporter architecture and regulatory roles of accessory domains. *FEBS Letters*, 580(4), 1023-1035. <https://doi.org/10.1016/j.febslet.2005.11.079>

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

## Minireview

## ABC transporter architecture and regulatory roles of accessory domains

Esther Biemans-Oldehinkel<sup>1</sup>, Mark K. Doeven<sup>1</sup>, Bert Poolman\**Department of Biochemistry, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands*

Received 4 November 2005; revised 30 November 2005; accepted 30 November 2005

Available online 12 December 2005

Edited by Gerrit van Meer

**Abstract** We present an overview of the architecture of ATP-binding cassette (ABC) transporters and dissect the systems in core and accessory domains. The ABC transporter core is formed by the transmembrane domains (TMDs) and the nucleotide binding domains (NBDs) that constitute the actual translocator. The accessory domains include the substrate-binding proteins, that function as high affinity receptors in ABC type uptake systems, and regulatory or catalytic domains that can be fused to either the TMDs or NBDs. The regulatory domains add unique functions to the transporters allowing the systems to act as channel conductance regulators, osmosensors/regulators, and assemble into macromolecular complexes with specific properties.

© 2005 Federation of European Biochemical Societies. Published by Elsevier B.V. All rights reserved.

**Keywords:** ABC transporter; Accessory domain; Architecture; ATP-binding cassette; Regulation of transport

## 1. Introduction

ATP-binding cassette (ABC) transporters are vital to any living system and are involved in the translocation of a wide variety of substrates ranging from ions, sugars, amino acids, vitamins, lipids, antibiotics and drugs to larger molecules such as oligosaccharides, oligopeptides and even high molecular weight proteins [1]. Owing to their function ABC transporters are involved in many crucial processes, including nutrient uptake, lipid trafficking, drug and antibiotic excretion, secretion of macromolecules, antigen presentation to cytotoxic T cells, and cell volume regulation, and they are often major players in complex pathways affecting gene expression (e.g., sporulation, competence and virulence development). In humans a still growing number of diseases is related to (dys)function of ABC transporters, the most common known examples being cystic fibrosis caused by defects in the Cystic Fibrosis Transmembrane conductance Regulator (CFTR/ABCC7), an ABC-type chloride channel, and multidrug-resistance conferred to tumor cells by P-glycoprotein (Pgp/MDR1/ABCB1) activity.

Irrespective of the direction of transport (uptake or excretion), the nature of the transported substrate (low or high

molecular weight molecule), or the physiological function, ABC transporters are composed of two hydrophobic transmembrane domains (TMDs) and two water soluble nucleotide-binding domains (NBDs) bound to the cytosolic face of the TMDs (Fig. 1). In addition to these core domains, hereafter referred to as the translocator or translocation unit, accessory domains or proteins can be part of the ABC transporter. In this paper, we review the architecture of ABC transporters, with emphasis on the types of accessory domains associated with the translocator. These accessory domains provide additional, often regulatory, functionalities to the transporters. The accessory domains can be recognized as additional transmembrane segments or internal or external domains linked to the TMDs, or extensions added to the NBDs. In many cases, it has been shown that the accessory domains can be deleted without affecting the translocation function, confirming that they do not form part of the core of the translocator.

## 2. Architecture of ABC uptake systems

The first identified components of ABC transporters were the substrate-binding proteins (SBPs) of ABC uptake systems that reside in the periplasmic space of Gram-negative bacteria, from which they could be released by cold osmotic shock [2–5]. These water soluble proteins bind the substrate in a cleft between two lobes and deliver it to the translocator in the cytoplasmic membrane. In Gram-positive bacteria and archaea, i.e., microorganisms lacking an outer membrane, SBPs are exposed on the cell surface and attached to the cytoplasmic membrane via a lipid-anchor or a transmembrane-peptide (to date, the latter has been only observed in archaea), or they can be fused to the TMDs resulting in two substrate-binding domains (SBDs) per functional complex (Fig. 1B; OpuA). In some cases, two SBDs fused in tandem are linked to the TMDs and these systems have a total of four extracytoplasmic substrate-binding sites (e.g., GlnPQ). Systems with SBDs fused to the TMDs can be also found in Gram-negative bacteria but less frequently than in Gram-positives [6].

In terms of number of different subunits, the most complex ABC type uptake systems known are the nickel- and (oligo)peptide transporters, in which each domain is present as a separate polypeptide, giving a total of (at least) five unique proteins required for transport (Fig. 1A and B; Nik and Opp). In many cases, two copies of the same NBD plus two different TMDs (e.g., the maltose, histidine, an glucose transporters) or, less frequently, two copies of one NBD plus one TMD are present (the vitamin B12 transporter Btu). Occasionally, the

\*Corresponding author. Fax: +31 50 3634165.  
E-mail address: b.poolman@rug.nl (B. Poolman).

<sup>1</sup> These authors contributed equally to this work.




Fig. 1. Domain architecture of ABC transporters. Schematically indicated are: SBPs (pac-man shaped), TMDs (rectangles), and NBDs (ovals). (A) Gram-negative bacteria (all the examples are from *Escherichia coli*): Nik, nickel transporter; Mal, maltose/maltodextrin transporter; Btu, vitamin B12 transporter; Fhu, siderophore/haem/vitamin B12 transporter; Rbs, ribose transporter. (B) Gram-positive bacteria and Archaea: Opp, oligopeptide transporter from *Lactococcus lactis*; Glc, glucose transporter from *Sulfolobus solfataricus*; OpuA, glycine betaine transporter from *Lactococcus lactis*; Gln, glutamine/glutamic acid transporter from *Lactococcus lactis*. (C,D) Functional and structural homologues are present in all three kingdoms of life. (C) Msb, lipid flippase from *Escherichia coli*; Cyd, cysteine exporter from *Escherichia coli*. (D) Mdl, mitochondrial peptide transporter from *Saccharomyces cerevisiae*; TAP1/2 (ABC2/3), human peptide transporter; PDR5, yeast pleiotropic drug resistance transporter; P-gp (MDR1/ABC1), human multidrug transporter.

NBDs or TMDs are fused (e.g., siderophore/haem and ribose transporters) such that in total only three unique proteins are used (Fig. 1A). The smallest number of unique subunits of which ABC uptake systems are composed, known to date, is two (Fig. 1B). This is the case for the glycine betaine and glutamine/glutamic acid transporters in which the SBPs are fused to either the N- or C-terminus of their cognate TMDs, and the second unique subunit is formed by the NBD [6].

### 3. Architecture of ABC efflux systems

In contrast to ABC uptake systems, which are present in the cytoplasmic membrane of prokaryotic species (bacteria and archaea), ABC efflux systems are found in every life-form ranging from microorganisms to man. ABC efflux pumps can be found in the cytoplasmic membrane of prokaryotes and eukaryotes, as well as in the organellar membranes of the higher organisms, that are, the endoplasmic reticulum, inner mitochondrial membrane, peroxisomal and vacuolar membranes. Although the common four-domain structure of the translocator (two TMDs plus two NBDs) is conserved, ABC efflux pumps are somewhat less complex compared to ABC uptake systems in that they do not require a SBP for function (Fig. 1C and D). The direction of transport can be deduced from the presence or not of this latter component. An excep-

tion to this rule has yet to be found. Even more frequently than in ABC uptake systems, the four core-domains of the translocator are fused, yielding only one or two unique polypeptides forming the functional unit (Fig. 1C and D). The most common organization is the “half-size transporter” in which a single TMD is fused to the N- or C-terminus of a NBD. The functional complex is either a homo- or heterodimer of two half-transporters (e.g., the lipid flippases, multi-drug transporters, cysteine exporter, and peptide transporters; Fig. 1C and D). Finally, all four core-domains (two half-transporters) can also be fused as is the case for the multi-drug transporters (e.g., P-gp, MRP1 and 2, and PDR5), and the ABC type chloride channel CFTR (Fig. 1D), and these are termed “full-length transporters”.

### 4. Transmembrane domains

The TMDs are two highly hydrophobic proteins or protein domains that create the channel through which the substrate passes during translocation (Fig. 1). In ABC uptake systems, they form the docking site for the SBP and transduce a signal to the NBDs to bind and hydrolyze ATP. The TMDs are bundles of  $\alpha$ -helices that traverse the membrane multiple times in a zig-zag fashion. Many typical ABC transporters conform to the paradigmatic “two-times-six”  $\alpha$ -helices structure, yielding

a total of 12 transmembrane segments (TMSs) per functional unit (Table 1). However, the number of TMSs may vary from anywhere between 5 and 11 for each individual TMD.

Whereas the NBDs show considerable sequence identity (see below), the TMDs (as well as the SBPs) have been less well conserved during evolution, reflecting the large diversity of substrates transported. TMDs of ABC uptake systems contain the so-called EAA motif (L-loop), which usually is found in the cytoplasmic loop between TMS 5 and 6 [7]. This motif is absent in ABC exporters.

In ABC efflux systems, the TMDs harbor the substrate-binding site(s). Evidence for the existence of a substrate-binding site in the TMDs of ABC uptake systems came from the analysis of SBP-independent mutants [8–11]. Mutations in the NBD of the histidine transporter (HisP) or the TMDs of the maltose transporter (MalF and G) resulted in an uncoupled phenotype, in which the SBP no longer interacted with the TMDs and the NBDs hydrolyzed ATP constitutively. In addition, a SBP-independent maltose transporter with a truncated

form of MalF had its substrate specificity changed from maltose to lactose, albeit with very high affinity constants [12]. Finally, the crystal structure of the BtuCD translocator revealed a cleft in the periplasmic half of the TMDs that could accommodate a vitamin B12 molecule and represent a genuine substrate-binding site [13].

## 5. Nucleotide-binding domains

The engines of ABC transporters are the NBDs as these power substrate translocation by ATP-binding and hydrolysis. Whereas the SBPs and TMDs are poorly conserved, the NBDs are more closely related and group the different transporter families in the ABC superfamily. They contain the ATP-binding cassettes, hence the name ABC transporters. Many crystal structures of isolated NBDs are available, yielding detailed information about the overall fold and the nucleotide-binding site(s) [14]. A NBD consists of two domains, a RecA-like and a

Table 1  
Overview of TMD topology in well characterized ABC transporters

TMD	Transport substrate	Number of TMSs			Total number of TMSs	Reference
		N-terminal	Core-domain	C-terminal		
<b>ABC uptake systems</b>						
HisQ	Histidine		5		10 (2 × 5)	[133]
HisM			5			
GlnP	Glutamine/glutamic acid		5 <sup>a</sup>		10 (2 × 5)	[52]
ProW	Glycine betaine/Proline betaine	2	5		14 (2 × 7)	[26]
OpuABC	Glycine betaine	2	5	1 <sup>b</sup>	16 (2 × 8)	[6]
OppB	Oligopeptides		6		12 (2 × 6)	[134]
OppC			6			
MalF	Maltose/ maltodextrins	2	6		14 (8 + 6)	[54,135]
MalG			6			
RbsC	Ribose		10 <sup>c</sup>		20 (2 × 10)	[136]
BtuC	Vitamin B12		10		20 (2 × 10)	[13]
FhuB	Siderophores/haem/vitamin B12		10 × 2		20	[137]
<b>ABC efflux systems</b>						
<b>Half-transporters</b>						
LmrA	Drugs		6 <sup>d</sup>		12 (2 × 6)	[138]
BmrA	Drugs		6 <sup>d</sup>		12 (2 × 6)	[139]
MsbA	Lipids		6		12 (2 × 6)	[16]
CydC	Cysteine/GSH		6		12 (2 × 6)	[140]
CydD			6			
HlyB	Haemolysin	(2) <sup>e</sup>	6		12–16 <sup>e</sup> (2 × 6–8)	[141,142]
TAP1 (ABCB2)	Antigenic	4	6		19 (10 + 9)	[58]
TAP2 (ABCB3)	peptides	3	6			
<b>Full-length transporters</b>						
P-gp (MDR1/ABCB1)	Drugs		6 × 2		12	[143]
BCRP (ABCG2)	Drugs		6 × 2		12	[144]
MRP1 (ABCG2)	Drugs		6 × 2		17 (5+12)	[144]
MRP2 (ABCC1)	Drugs	5	6 × 2		17 (5 + 12)	[144]
SUR1 (ABCC8)	None/unknown <sup>f</sup>	5	6 × 2		17 (5 + 12)	[145]
CFTR (ABCC7) <sup>g</sup>	Chloride		6 × 2		12	[146]
Ste6	Mating pheromone		6 × 2 <sup>h</sup>		12	[147]
ABCR (ABCA4)	Retinal/ retinal phosphatidylethanolamine		6 × 2 <sup>i</sup>		12	[24]

<sup>a</sup>Two SBDs are fused in tandem to the first TMS. The N-terminal signal sequence, preceding the SBDs, is cleaved off.

<sup>b</sup>The SBD is fused to a C-terminal signal anchor sequence (the 8th TMS).

<sup>c</sup>Initially claimed to have 6 TMSs [148]; later shown to possess 10 TMSs [136].

<sup>d</sup>No experimental evidence; topology based on homology with other multidrug-efflux systems.

<sup>e</sup>HlyB is thought to contain 6–8 TMSs; the presence of two N-terminal TMSs has not been unambiguously determined.

<sup>f</sup>Sulfonylurea receptors (SURx) associate with K<sub>IR6.x</sub> to form ATP-sensitive potassium channels.

<sup>g</sup>CFTR is a chloride channel rather than transporter although it has the classical two TMDs plus two NBDs structure observed for genuine ABC transporters.

<sup>h</sup>Experimental data only for the N-terminal TMD.

<sup>i</sup>Two large extracytoplasmic glycosylated domains are inserted between the first and second TMS of both the N- and C-terminal TMDs.

helical domain, the latter being specific for ABC transporters. In the NBD monomer, the ATP-binding site is formed by two conserved sequences, the Walker A (P-loop) and Walker B motifs, present in many ATP-binding proteins and not restricted to ABC proteins [15]. These motifs bind the phosphates of ATP and ADP, and coordinate a  $Mg^{2+}$ -ion via  $H_2O$  in the nucleotide-binding site, respectively. A glutamate directly following the Walker B motif might be the catalytic base for hydrolysis, since it binds to the attacking water and  $Mg^{2+}$ -ion, and mutation of this residue leads to loss of ATPase activity [14]. The Q-loop also forms hydrogen bonds with  $Mg^{2+}$  and the attacking water. A conserved histidine in the H motif hydrogen bonds with the  $\gamma$ -phosphate of ATP. In the NBD dimer, the signature sequence, LSGGQ, (the motif that distinguishes ABC proteins from other nucleotide-binding proteins) of one NBD binds to the  $\gamma$ -phosphate of ATP bound to the other NBD, thereby completing the nucleotide-binding sites [14]. Thus, two ATPs are sandwiched in head-to-tail orientation between two NBDs in the catalytically active dimer.

## 6. ABC transporter mechanism

Given the conserved architecture of two TMDs plus two NBDs across the entire ABC transporter superfamily, one could expect a common translocation mechanism for ABC uptake and efflux systems. Indeed, from the culmination of vast amounts of functional data, in combination with the recent X-ray structures of complete ABC transporters (for both efflux and uptake systems; [13,14,16–21]), a consensus mechanism is emerging. The simplest model of SBP-dependent uptake encompasses four steps (Fig. 2): (i) substrate binds to the SBP; (ii) the closed, liganded SBP docks onto the TMDs and signals the NBDs to cooperatively bind ATP; (iii) upon ATP-binding, a binding site in the TMDs is made available to the outside, the SBP opens and substrate is transferred to




Fig. 2. Model of the translocation mechanism of a SBP-dependent ABC transporter. For explanation, see text.

the binding site in the TMDs; and (iv) ATP hydrolysis dissociates the NBD dimer and results in reorientation of the binding site from an outward- to an inward-facing conformation. The substrate arrives in the cytoplasm and the SBP is released from the TMDs. Alternatively, it cannot be ruled out that upon ATP binding the substrate is transferred directly to the cytoplasm (step iii) and ATP hydrolysis in combination with dissociation of ADP and inorganic phosphate is solely used to reset the system for another translocation cycle. Whereas step (i) is unique to SBP-dependent uptake systems, steps (ii–iv) could be envisaged for ABC efflux systems as well. In step (ii), a substrate bound to the inward-facing binding site would signal the NBDs to bind ATP. Subsequently, step (iii), ATP-binding reorients the binding site and substrate is released to the outside. (iv) ATP-hydrolysis resets the system by converting the TMDs back to the inward-facing conformation.

Although ABC uptake and efflux systems could share a common translocation mechanism, there are some indications that things might be more complicated. First of all, although in nature TMDs and NBDs appear as separate modules or fused together in almost any possible combination (Fig. 1), to the best of our knowledge, no ABC uptake systems are known that have a TMD fused to a NBD as is the case with half- or full-length efflux systems. For ABC efflux systems, this latter architecture seems to be a universal feature. Second, the conserved EAA motif (see above) is found exclusively in ABC uptake systems. This motif is thought to be important for the interaction of the TMDs with the NBDs. Third, the structures of the ABC efflux system MsbA showed an additional intracellular domain between the TMD and NBD that was not present in the structure of the SBP-dependent vitamin B12 uptake system BtuCD [13,16–18]. Fourth, NBD2 of CFTR was shown, in addition to hydrolyzing ATP ( $Mg\text{-ATP} + H_2O \rightarrow Mg\text{-ADP} + Pi$ ), to display adenylate kinase activity ( $Mg\text{-ATP} + AMP \leftrightarrow Mg\text{-ADP} + ADP$ ), which influenced channel activity [22]. It remains to be established whether the adenylate kinase activity is unique to CFTR or ABC efflux systems, but some important structural differences between ABC uptake and efflux systems seem to exist. The significance of these observations in terms of a general ABC transport mechanism awaits further experimentation.

## 7. Accessory domains in ABC transporters

In addition to the ubiquitous four core-domains, that are two TMDs plus two NBDs, accessory domains can be found in several ABC transporters (Fig. 3). These extra domains often perform different functions and can be found separate from, attached to, or integrated into different parts of the core ABC transporter. On the basis of their location (extrinsic or intrinsic to the membrane) and function (regulatory or catalytic) within the cognate transporters, we divide the well-characterized accessory domains of ABC transporters into four different groups: (i) extracytoplasmic domains; (ii) membrane-embedded domains; (iii) cytosolic regulatory domains and (iv) cytosolic catalytic domains.

### 7.1. Extracytoplasmic domains

One of the characteristics of accessory domains is that they are not part of the general ABC transporter core complex. There are three main examples of known extracytoplasmic






Fig. 3. Well-characterized accessory domains in ABC transporters. TMD core domains are boxed (grey lines). (A) Structure of the oligopeptide-binding protein AppA from *Bacillus subtilis* (PDB Accession No. 1XOC). Domain I and III (blue and red, respectively) are common to SBPs. The extra domain (II, orange) contributes with a few residues to the binding site by interacting with the N-terminal part of the bound nonameric peptide (yellow). (B) Topology of OpuABC, TMD with SBP fused to C-terminus, of the glycine betaine transporter OpuA from *Lactococcus lactis*. The NBD is a separate protein (Fig. 1B). (C) Topology of members of the ABCA family. Extracellular loops (ECD-1 and ECD-2) and cytosolic regulatory domains (R1 and R2) are shown. (D) TMD topology of TAP1 and 2 half-transporters. (E) Topology of the SUR/MRP (ABCC) family. The extra TMD (TMD0) and L0-loop are indicated. (F) Structure of dimeric nucleotide-bound MalK from *Escherichia coli* (PDB Accession No. 1Q12), the NBD of the maltose/maltodextrin transporter. The common helical and RecA-like domains are shown in red and orange for the first monomer and dark purple and blue for the second, respectively. Two ATPs (yellow) are sandwiched between the NBDs. The additional C-terminal domain is shown in yellow for the first and bright purple for the second dimer subunit. (G) Structure of a tandem CBS domain from *Thermotoga maritima* (PDB Accession No. 1O50) as present in the C-terminal part of the NBD of OpuA (OpuAA). The N- and C-terminal halves are colored red and blue, respectively. (H) Topology of CFTR. The N-tail and regulatory (R) domain are indicated. (I) Putative TMD topology of LcnC, a bacteriocin exporter with a N-terminal peptidase-like domain.

domains or proteins that can be considered accessory units (i) the large extracytoplasmic domains (ECDs) found in members of the ABCA family; (ii) the extracellular N-terminal domain

of ProW (and closely related homologues), the membrane component of the ProU system from *Escherichia coli*; and (iii) the SBDs or proteins found in prokaryotes.

Members of the ABCA family are full-length transporters consisting of two TMDs of 6 TMSs each and two NBDs, each located C-terminal of one of the TMDs (Table 1; [23]). The two ECDs are predicted after the first TMS of the N- and C-terminal half of the transporter (Fig. 3C) and are 600 and 275 amino acids long in ABCA4 (ABCR) [24]. Eight *N*-glycosylation sites were mapped by mutagenesis onto the ECDs of ABCA4, four in the N-terminal half and four in the C-terminal half of the protein [24]. Removal of these *N*-glycosylation sites resulted in reduced expression levels, although the mutants retained the capacity to bind ATP. It has been postulated that the two ECDs are linked by at least one disulfide bridge. Intra- and intermolecular crosslinking of cysteine residues in an ECD was shown for ABCG2 (BCRP) [25], a half-transporter with the ECD located between TMS 5 and TMS 6. Two ABCG2 subunits were linked by a disulfide bridge, and two remaining cysteine residues formed intramolecular disulfide bonds within the ECD. Although mutations in the ECDs have been shown to influence the substrate specificity of the transporters, a distinct function is not known for these relatively large ECDs.

The second example of an ECD is found in the proline and glycine betaine transport system (ProU) of *E. coli*. Here the membrane component, ProW, is composed of an extended N-terminal region of about 100 amino acid residues followed by 7 TMS (Table 1). The function of the N-terminal region is unknown, but a *phoA* and *LacZ* fusion study showed that its location is periplasmic [26].

The best-studied ECDs or proteins are the SBDs or proteins of ABC-type uptake systems. These domains or proteins are considered to be accessory, because these components are not intrinsic to the translocation process as has been demonstrated for ABC transporter mutants that function in the absence of SBPs (see Section 4). SBPs are the main determinants of substrate specificity of ABC uptake systems and confer high-affinity on the transport process with dissociation constants ( $K_d$ ) in the (sub)micromolar range. They have been a gold-mine for structural studies due to their well behavior during overexpression, purification and crystallization trials. Although some early experiments suggested that SBPs might self-associate to form dimers or higher order oligomers [27–30], crystal structures of numerous SBPs have led to the generally accepted view that they are monomers with one substrate-binding site per molecule [31]. Even though systems are known that use multiple SBPs to broaden the substrate specificity of the transporter (several peptide transporters; [32–34]), only a single SBP is required for transport function. The high resolution structure of the vitamin B12 importer BtuCD confirmed that there is space for only a single SBP to dock onto the TMDs [35]. The ABC transporters with one or two SBDs fused to the TMD are thought to function similarly, except for that the presence of a second, third or fourth SBD might affect the kinetics of the translocation process [36] or perhaps broaden the substrate specificity.

The archetype SBP consists of two domains or lobes connected by a flexible linker. The two lobes close and engulf the ligand upon substrate-binding (Venus's Flytrap model; [37]). Structures are available of SBPs in open-unliganded, closed-unliganded, open-liganded, and closed-liganded forms [38–42], of which the latter conformation is thought to productively interact with the membrane complex [43–45]. In some cases, unliganded SBPs also have access to the TMDs, thereby inhibiting ABC transporter function (e.g., in case of the malt-

ose- and histidine-binding proteins; [46,47]). These observations, however, could not be confirmed in similar experiments performed with the oligopeptide ABC transporter [45]. Mutational and suppressor analyses suggested that each lobe of the SBP binds to one of the TMDs [8,48]. Site-directed mutagenesis and pre-steady-state kinetic analysis indicated that donation of the substrate by the SBP to the TMDs is the rate-limiting step during the ABC uptake catalytic cycle of the oligopeptide transporter Opp [49,50].

Although a typical SBP consists of two lobes (domains) that are connected by a flexible hinge [31], exceptions to this rule are the SBPs of the nickel/oligopeptide transporter family (Fig. 3A). OppA contains an extra, third domain [38]. The function of this additional domain is unknown, although in AppA, an OppA homologue from *Bacillus subtilis*, it has been shown to form part of the binding site for nonameric peptides [51].

Two ABC transporter families (OTCN and PAO) are known in which one or two SBPs are fused to either the N- or C-terminus of one TMD, yielding a total of two or four SBDs per functional complex (Fig. 1B, OpuA and GlnPQ; [6]). Two SBDs are present in OpuA, the osmoregulatory ABC transporter from *Lactococcus lactis*, although for activity only one SBD is needed. The SBDs of OpuA interact in a cooperative manner by stimulating either the docking of one domain onto the TMDs or substrate donation to the TMDs (Fig. 2, steps (ii) and (iii), respectively; [36]). However, this cooperative interaction has not been established for other SBP dependent ABC transporters, e.g., the glutamate/glutamine transporter GlnPQ from *L. lactis*, containing four SBDs [52]. In GlnPQ, the SBD in the primary sequence nearest to the translocator domain (SBD2) is used as high affinity receptor that delivers the substrate to the TMDs. SBD1, on the contrary, is not required for high affinity transport and is not sufficient for transport but does bind glutamine with a low affinity (dissociation constant in the (sub)millimolar instead of micromolar range; Schuurman–Wolters, unpublished).

## 7.2. Membrane-embedded domains

The majority of ABC transporters are predicted to have six TMSs per TMD core (Table 1). Thus, in a typical transporter a total of 12 TMSs are present. Exceptions are known for ABC uptake systems that contain 5 or 10 TMSs as a single TMD core that is both essential and sufficient for transport. In addition to this somewhat variable number of TMSs in the TMDs, several ABC transporters contain extra TMSs that are not part of the core TMD and can be considered as accessory domains. Although this phenomenon frequently occurs, the function of these extra domains is in many cases unknown. For example, within the OTCN family of ABC transporters [53], sequence alignments reveal a core TMD of 5 TMSs but homologues can have in total either 5, 6 or 8 TMSs. The core TMSs are generally much better conserved than the accessory TMSs ([6]; unpublished). Within the OTCN family, some polypeptides (f.i., the OpuABC subunit of OpuA, Fig. 3B, [6]; and the ProW subunit of ProU, [26]) have two extra TMSs N-terminal of the TMD core but their function is unknown. In addition, some members have an 8th TMS (or 6th TMS in case the first two are missing), C-terminal of the TMD core, that serves as signal anchor sequence for the translocation of the extracytoplasmic SBD. Also, members of the OSP family to which the maltose transporter belongs can have two extra N-terminal

TMSs (MalF; [54]). For the maltose transporter, the extra TMSs of MalF have been proposed to be an export signal, but most of the two TMSs could be removed without any effect on maltose transport activity [54–56].

A system in which a clear function could be assigned to the accessory TMSs is the transporter associated with antigen processing (TAP), present in the endoplasmic reticulum membrane. Based on hydrophobicity analysis and sequence alignments 10 (4 accessory plus 6 core) and 9 (3 accessory plus 6 core) TMSs were predicted for human TAP1 and TAP2, respectively (Table 1 and Fig. 3D; [57,58]). Although the core-TAP complex (the C-terminal 6 TMSs from TAP1 and TAP2 combined) represented the minimal functional unit for transport, the N-terminal domains of TAP1 and TAP2 were essential for recruitment of tapasin [58,59], a glycoprotein known to mediate association between TAP and MHC class I molecules [60]. Consequently, the extra domains were proposed to mediate the assembly of the macromolecular peptide-loading complex. Subsequently, it was shown that in TAP2 the tapasin-binding site is exclusively located in the N-terminal domain. Upon deletion of the N-terminal domain of TAP1, tapasin could still interact with the TAP1 TMD core [61]. TAP variants lacking the N-terminal domain of TAP2, but not of TAP1, were disturbed in the quality control of MHC I loading, and formed peptide loading complexes that were disturbed in their physical interaction with calreticulin, calnexin, and ER60.

Instead of a few extra TMSs, some ABC systems, including the sulfonyleurea receptors SUR1 and SUR2, and the multi-drug-resistance proteins MRP1, 2, 3, 6 and 7, have 5 accessory TMSs that may form a domain by itself (Table 1 and Fig. 3E; [62–64]). These transporters belong to the ABCC family of the ABC transporter superfamily. The core complex in this subfamily is a full-length transporter composed of two TMDs (TMD1 and TMD2, each containing 6 TMSs) and two NBDs (NBD1 and NBD2) that follow TMD1 and TMD2, respectively. The extra TMSs (indicated as TMD0) are linked via a cytoplasmic loop (L0) to the N-terminus of TMD1 (Fig. 3E). In MRP1, the entire TMD0 could be deleted without having any effect on transport activity or on the trafficking and insertion of the protein into the basolateral membrane [65,66]. This contrasted results obtained with SUR1 and MRP2, where deletion of TMD0 affected channel activity, processing and trafficking [67–71]. Recent studies revealed that TMD0 is needed for trafficking if the C-terminal region of MRP1 is mutated or deleted [72]. It seems that in MRP1 TMD0 and the C-terminal region contain redundant trafficking signals that become essential when one or the other region is missing or mutated.

In SUR, TMD0 is required for activation of the inwardly rectifying potassium channel ( $K_{IR}$ ) subunits and anchoring of the  $K_{IR}$  subunits to the ABC transporter core of SUR, thereby generating functional  $K_{ATP}$  channels [67,68]. Thus, similar to the TAP system, the accessory membrane-embedded domains of SUR provide additional protein interaction sites and serve a regulatory purpose.

### 7.3. Cytosolic regulatory domains

Regulation of many ABC transporters takes place at the level of gene expression. Most ABC transporter operons encode a transcriptional regulator that connects the level of protein to environmental conditions. Expression of the osmoregulatory ABC transporter OpuA from *L. lactis* is, for example, regu-

lated in response to osmotic stress via the transcription factor OpuR [73]. However, the activity of ABC transporters, once expressed, may also be regulated. For this purpose, additional (autoregulatory) domains in the transporter itself are required. Examples are the L0 linker in MRP1 and SUR, the C-terminal domain of MalK of the maltose transporter from *E. coli* and *S. typhimurium*, the tandem cystathionine- $\beta$ -synthase (CBS) domains of OpuA from *L. lactis* and other members of the OTCN family, and the R-domain of CFTR (Fig. 3E, F, G and H, respectively). Although these domains all regulate ABC transporter activity, they use different mechanisms to do so.

The maltose/maltodextrin transporter from *E. coli* and *S. typhimurium* is directly involved in transcriptional regulation of, amongst others, itself. The regulation takes place at the C-terminal domain of MalK, the NBD of the transporter (Fig. 1A (Mal) and 3F; [74]). Crystal structures are available of MalK [75], and GlcV, the NBD of the glucose transporter from *Sulfolobus solfataricus* [76]. These NBDs contain an accessory C-terminal domain with a common tertiary fold. The C-terminal domain of MalK is involved in two regulatory processes, at least in enteric bacteria [77]. First, via its C-terminal domain, MalK is subject to regulation by  $IIA^{glc}$ , a component of the bacterial phosphoenolpyruvate-dependent sugar:phosphotransferase system (PTS) [75,78,79].  $IIA^{glc}$  can be in the unphosphorylated and phosphorylated state, depending on the availability of PTS sugars. In the unphosphorylated form, i.e., when a PTS sugar such as glucose is present,  $IIA^{glc}$  binds to MalK and thereby prevents maltose uptake. The inhibition is relieved when glucose is exhausted and  $IIA^{glc}$  becomes phosphorylated. This regulatory process determines the hierarchy of sugar utilization and is known as inducer exclusion mechanism. GlcV is unlikely to be involved in this type of regulation, since archaea lack a PTS. Second, MalK inhibits the activity of MalT, the positive regulator of the maltose regulon, by competing with MalT for maltotriose binding [75,80–82]. Binding of maltotriose to MalT is required for oligomerization of the transcription factor, a prerequisite for the interaction of MalT with its DNA binding sites [83,84]. In addition to MalK, two other proteins, MalY and Aes, can also bind MalT [85,86], creating an intricate regulatory network that will not be detailed further in this paper. MalK is thought to possess two distinct protein binding sites, one for MalT and one for  $IIA^{glc}$ , because mutations in MalK affecting inducer exclusion do not interfere with MalT inactivation [78,87]. By mapping known mutations onto the structure of MalK (the one from *Thermococcus litoralis*), the MalT interaction site has been proposed to be located on an exposed surface of the C-terminal regulatory domain. Point mutations that confer an inducer exclusion insensitive phenotype form a patch adjacent to and oriented perpendicularly to the MalT interaction site [75].

The osmoregulatory ABC transporter OpuA protects the cell against hyperosmotic stress, ultimately plasmolysis, through uptake of the compatible solute glycine betaine, and thereby reverses the osmotic shrinkage of the cell [88]. When reconstituted in proteoliposomes, OpuA is activated by increasing internal ionic strength [89]. The threshold for activation by ions is dependent on the ionic lipid content (charge density) of the membrane, indicating that the ionic signal is transduced to the transporter via critical interactions of protein domains with membrane lipids [90]. The ABC transporter



OpuA, as do other members of the OTCN family, has two CBS domains (Fig. 3G) in tandem at the C-terminal end of the ATPase subunit OpuAA. With two ATPase subunits per translocation unit, a total of four CBS domains are present in OpuA. This so-called CBS module of OpuA constitutes the sensor that switches the transporter between an inactive, electrostatically locked and an active, thermally relaxed state, by interacting with the negatively charged membrane surface in response to the ionic strength [89]. The switching mechanism is an effective means for cells to respond to osmotic shifts, as an increase in medium osmolality will result in a decrease in cell volume, and the accompanying increase in cytoplasmic ionic strength will activate the transporter. Glycine betaine accumulation, followed by water influx, will increase the cell volume and decrease the ionic strength in due time, and the transporter will ultimately be switched off. This inherent feedback mechanism prevents over-accumulation of glycine betaine and couples transporter activity with ionic strength in the cell.

CFTR is an ABC-type chloride channel that is regulated by protein kinase A-mediated phosphorylation of its cytoplasmic regulatory (R) domain [91]. The R-domain is located between NBD1 and TMD2 of the full-length transporter (Fig. 3G), and contains a number of charged residues and multiple consensus phosphorylation sites [92,93]. The unphosphorylated R-domain is thought to inhibit channel activity by interaction with the CFTR core transporter. Release of the R-domain by phosphorylation would activate the channel [94–96]. Thus far, phosphorylation dependent release of the R-domain has not been demonstrated unequivocally. Correspondingly, a recent study strongly suggested that phosphorylation results in enhanced association of the R-domain with the transporter rather than in its release [97]. The mechanism of regulation suggested by Chappe et al. was that binding of the R-domain to the transporter induces intramolecular rearrangements that allow nucleotide-induced conformational changes to be transmitted from the NBDs to the TMDs. A second system that was shown to be regulated by phosphorylation of a cytoplasmic domain is ABCA1 [98]. This half transporter is a member of the ABCA family in which the R-domains (R1 and R2) are located C-terminal of the NBDs of the C- and N-terminal half of the transporter, respectively (Fig. 3C). Phosphorylation of the R1 domain of ABCA1 diminished the phospholipid flip-flop, apolipoprotein binding, and cholesterol/lipid extrusion activities.

The N-terminal tail of CFTR (N-tail; Fig. 3H), next to its R-domain, also plays a role in regulating CFTR channel activity. It modulates the rate of channel openings, probably by interacting with components of CFTR (R-domain or NBD), or with the pore itself [99]. Next to a suggested physical interaction with a portion of the R-domain [100], the N-tail and the C-terminal tails of CFTR have been shown to bind several inhibitors and stimulators (physical modulators) of CFTR activity [101]. These interaction partners either bind directly to CFTR or mediated through various PDZ domain-containing proteins. One of the PDZ containing proteins is the  $\text{Na}^+/\text{H}^+$  exchanger regulatory factor (NHERF) that can form homo- and hetero-oligomers between CFTR and other membrane proteins, e.g., the  $\beta_2$ -adrenergic receptor [102,103]. It has been recently shown that the assembly of the C-terminal domains of CFTR is regulated by ezrin, an actin binding protein [104]. The di-

meric CFTR-NHERF-ezrin interaction would provide a pathway to regulate CFTR activity via the cytoskeleton actin network. Moreover, there is evidence that when two subunits of CFTR are brought in close proximity by NHERF, the open probability of the channels is increased [105]. Finally, the increasing number of proteins interacting with CFTR suggests that the chloride conductor is not only regulated itself, but that CFTR might also regulate the activities of other ion channels, receptors, or transporters.

The N-tail of CFTR contains a cluster of acidic residues which are located onto one face of a predicted  $\alpha$ -helix [100,106]. A similar amphipatic  $\alpha$ -helix has been predicted for the L0 linker in the sulfonyleurea receptor, SUR [107]. This linker is located between TMD0 and TMD1 (Fig. 3E), and consequently links TMD1 and TMD2 of SUR to the TMD0/ $\text{K}_{\text{IR}}$  complex. Interaction of the N-terminus of  $\text{K}_{\text{IR}}$  with L0 has a bidirectional effect on the open probability of the  $\text{K}_{\text{ATP}}$  channel, i.e. structurally distinct segments of the linker can saturate or attenuate the open probability of the channel [68,108,109]. Additionally, the L0 linker has been reported to be involved in the binding of the sulfonyleurea glibenclamide which inhibits  $\text{K}_{\text{ATP}}$  channels [110]. Similar to CFTR, there is an increasing number of proteins known to interact with  $\text{K}_{\text{ATP}}$  channels [111]. Although these proteins do not bind directly to the ABC component, the sulfonyleurea receptor, they form part of the same supramolecular complex. A variety of glycolytic enzymes, adenylate kinase and creatine kinase have been shown to be part of the  $\text{K}_{\text{ATP}}$  channel complex and regulate its function.

The requirement of the L0 linker for transport function has also been demonstrated for MRP1 [107]. In MRP1, two  $\alpha$ -helices have been predicted in L0, although the first one with low probability [66]. L0, and especially the second helix in the loop, are absolutely essential for MRP1 function [65,112–114], and might be required for correct folding and trafficking of the protein [66]. L0 has been proposed to interact both with the core domains of MRP1 and with the hydrophobic membrane regions of other proteins, or with the membrane itself [66,107]. In addition, it has been suggested that at least part of the glutathione binding site is located in L0, although there is no direct evidence for this notion [115].

#### 7.4. Accessory catalytic domains

The type I secretion pathway is used to export competence-stimulating peptides, bacteriocins, haemolysins, and non-proteinaceous material, independent of the Sec system [116–119]. In this pathway, ABC transporters have dual functions, that are, export of polypeptides and removal of leader peptides. Similar to the Sec pathway, the (poly)peptides are synthesized ribosomally and targeted for export by a N-terminal sequence of between 15 and 30 amino acids [120], that is removed upon secretion across the membrane. Removal of this signal sequence is taken care of by a peptidase domain associated with dedicated ABC transporters. The signal sequences with the consensus LSXXELXXIXGG are usually cleaved after the two conserved glycine residues [121] and are therefore called double-glycine-type leader peptides. The polypeptides that are released from the cell, after processing of the leader peptide, range in length between 17 and 80 amino acids [120] and often require post-translational modification prior to secretion to become biologically active. The secreted peptides rarely share significant homology, which is reflective of their

functions as molecules that act in a strain- or species-specific manner, typically as competence-stimulating peptides (ComA in *Streptococcus pneumoniae*) or as class I (lantibiotics) or class-II (non-lantibiotics) bacteriocins.

The secretion of ComA and bacteriocins is mediated by dedicated ABC half transporters that possess an accessory N-terminal domain of approximately 160 amino acids to remove the leader peptide [122]. The peptidase domain has characteristics of a cysteine protease, including conserved cysteine and histidine residues in the active site that are essential for proteolytic activity [123]. How the peptidase domain interacts with the other components of the transporter to promote secretion is unknown. For LcnC, the ABC transporter responsible for export of the bacteriocin LcnA, the peptidase domain was located at the cytoplasmic face of the membrane on the basis of PhoA and LacZ fusion studies (Fig. 3I; [124]).

In Gram-negative bacteria, the type I secretion systems are composed of a protein complex that spans the cytoplasmic membrane (ABC transporter), periplasm (membrane fusion protein) and outer membrane (porin-like structure). The best-characterized system is the one for secretion of HlyA, a 110 kDa haemolytic toxin, in *E. coli*. This system is composed of the ABC transporter HlyB, the membrane fusion protein HlyD, and the outer membrane factor TolC. Contrary to the type I secretions systems for peptides and bacteriocins, HlyB does not contain a peptidase like domain and HlyA does not contain a leader peptide. Instead, type I proteins like haemolysin contain a poorly conserved C-terminal secretion signal [125]. The secretion signal domain of HlyA has been shown to interact with the NBD in vitro [126], but the possibility cannot be excluded that the signal region of HlyA may interact with the TMD of HlyB. Also for the interaction between HlyB and HlyD there is not a clearly defined region or domain in the ABC transporter.

## 8. ABC transporter domains in other systems

Whereas accessory domains in ABC transporters carry out important regulatory and catalytic functions, (core) components of ABC transporters can have dual functions or be integrated into completely different systems where they are vital for function. The ABC Walker A and B motifs are found in many proteins [15], the best known examples probably being myosin, and the DNA repair enzymes Rad50 and MutS. Moreover, Rad50 and MutS not only contain the Walker A and B motifs but also the ABC signature sequence LSGGQ and are genuine ABC proteins, albeit without transport function. In fact, a subset of ABC proteins is involved in DNA maintenance and protein synthesis, e.g., recombination, DNA repair, chromosome condensation and segregation, and translation elongation. These latter proteins exert their functions in the cytoplasm and/or nucleus rather than in the membrane [127].

SBPs of bacterial uptake systems have been shown to act as chaperones [128], and can be used as chemotaxis receptors [129]. In addition, SBP-dependent secondary transporters have been discovered [130], but an evolutionary relatedness of these SBPs to those of ABC transporters is not evident. Finally, domains homologous to prokaryotic SBPs function as receptors in ionotropic glutamate receptors present in the central nervous system of higher eukaryotes [131,132].

## 9. Conclusions

The core complex of an ABC transporter is composed of two NBDs and two TMDs, each in most cases with five or six transmembrane  $\alpha$ -helical segments. A notable exception to this rule is the vitamin B12 transporter BtuCD with two times ten TMSs in the TMDs, that, on the basis of the crystal structure, all seem critical for the formation of the translocation path. The RbsC and FhuB transporters may also conform to this exception. Additional TMSs associated with the core ABC complex seem to play regulatory roles as is best documented for the transporter involved in antigen presentation and the TMD0 domain of the sulfonyleurea receptors.

The SBPs are accessory components of the ABC uptake systems that have evolved to signal substrate availability to (the activation of) the ATPase, following the binding of liganded SBPs to the TMDs. Moreover, the SBPs increase the affinity of the transporters for their substrate(s) by several orders of magnitude. For a number of SBPs, functions are known that are not related to transport, including chaperoning and signal transduction. Several ABC transporters have evolved to play crucial roles in processes such as cell volume regulation, control of channel activity and antigen presentation and therefore acquired accessory regulatory functions. The CBS domains, protein modules linked to a wide variety of proteins (channels, transporters, enzymes and transcription factors), in OpuA and ProU add osmosensing and osmoregulatory functions to the transporters. Finally, ABC transporters (type I secretion systems) can have a peptidase domain that cleaves off the signal sequence prior to translocating the mature peptide (bacteriocin) or protein.

*Acknowledgments:* This work was supported by a “Top-subsidie” grant (NWO-CW, Grant No. 700-50-302). We thank Barry Holland for valuable discussion on the type I secretion systems.

## References

- [1] Higgins, C.F. (1992) ABC transporters: from microorganisms to man. *Annu. Rev. Cell. Biol.* 8, 67–113.
- [2] Neu, H.C. and Heppel, L.A. (1965) The release of enzymes from *Escherichia coli* by osmotic shock during the formation of spheroplasts. *J. Biol. Chem.* 240, 3685–3692.
- [3] Pardee, A.B. (1968) Membrane transport proteins. Proteins that appear to be parts of membrane transport systems are being isolated and characterized. *Science* 162, 632–637.
- [4] Berger, E.A. (1973) Different mechanisms of energy coupling for the active transport of proline and glutamine in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 70, 1514–1518.
- [5] Berger, E.A. and Heppel, L.A. (1974) Different mechanisms of energy coupling for the shock-sensitive and shock-resistant amino acid permeases of *Escherichia coli*. *J. Biol. Chem.* 249, 7747–7755.
- [6] Van der Heide, T. and Poolman, B. (2002) ABC transporters: one, two or four extracytoplasmic substrate-binding sites? *EMBO Rep.* 3, 938–943.
- [7] Saurin, W. and Dassa, E. (1994) Sequence relationships between integral inner membrane proteins of binding protein-dependent transport systems: evolution by recurrent gene duplications. *Prot. Sci.* 3, 325–344.
- [8] Treptow, N.A. and Shuman, H.A. (1985) Genetic evidence for substrate and periplasmic-binding-protein recognition by the MalF and MalG proteins, cytoplasmic membrane components of the *Escherichia coli* maltose transport system. *J. Bacteriol.* 163, 654–660.
- [9] Speiser, D.M. and Ames, G.F.-L. (1991) *Salmonella typhimurium* histidine periplasmic permease mutations that allow transport in

- the absence of histidine-binding proteins. *J. Bacteriol.* 173, 1444–1451.
- [10] Petronilli, V. and Ames, G.F.-L. (1991) Binding protein-independent histidine permease mutants. Uncoupling of ATP hydrolysis from transmembrane signaling. *J. Biol. Chem.* 266, 16293–16296.
- [11] Covitz, K.M., Panagiotidis, C.H., Hor, L.I., Reyes, M., Trepow, N.A. and Shuman, H.A. (1994) Mutations that alter the transmembrane signalling pathway in an ATP binding cassette (ABC) transporter. *EMBO J.* 13, 1752–1759.
- [12] Merino, G. and Shuman, H.A. (1998) Truncation of MalF results in lactose transport via the maltose transport system of *Escherichia coli*. *J. Biol. Chem.* 273, 2435–2444.
- [13] Locher, K.P., Lee, A.T. and Rees, D.C. (2002) The *E. coli* BtuCD structure: a framework for ABC transporter architecture and mechanism. *Science* 296, 1091–1098.
- [14] Davidson, A.L. and Chen, J. (2004) ATP-binding cassette transporters in bacteria. *Annu. Rev. Biochem.* 73, 241–268.
- [15] Walker, J.E., Saraste, M., Runswick, M.J. and Gay, N.J. (1982) Distantly related sequences in the  $\alpha$ - and  $\beta$ -subunits of ATP synthase, myosin, kinases and other ATP-requiring enzymes and a common nucleotide binding fold. *EMBO J.* 1, 945–951.
- [16] Chang, G. and Roth, C.B. (2001) Structure of MsbA from *E. coli*: a homolog of the multidrug resistance ATP binding cassette (ABC) transporters. *Science* 293, 1793–1800.
- [17] Chang, G. (2003) Structure of MsbA from *Vibrio cholerae*: a multidrug resistance ABC transporter homolog in a closed conformation. *J. Mol. Biol.* 330, 419–430.
- [18] Reyes, C.L. and Chang, G. (2005) Structure of the ABC transporter MsbA in complex with ADP·vanadate and lipopolysaccharide. *Science* 308, 1028–1031.
- [19] Higgins, C.F. and Linton, K.J. (2004) The ATP switch model for ABC transporters. *Nat. Struct. Mol. Biol.* 11, 918–926.
- [20] Locher, K.P. (2004) Structure and mechanism of ABC transporters. *Curr. Opin. Struct. Biol.* 14, 426–431.
- [21] Van der Does, C. and Tampé, R. (2004) How do ABC transporters drive transport? *Biol. Chem.* 385, 927–933.
- [22] Randak, C.O. and Welsh, M.J. (2005) Adenylate kinase activity in ABC transporters. *J. Biol. Chem.* 280, 34385–34388.
- [23] Peelman, F., Labeur, C., Vanloo, B., Roosbeek, S., Devaud, C., Duverger, N., Deneffe, P., Rosier, M., Vandekerckhove, J. and Rosseneu, M. (2003) Characterization of the ABCA transporter subfamily: identification of prokaryotic and eukaryotic members, phylogeny and topology. *J. Mol. Biol.* 325, 259–274.
- [24] Bungert, S., Molday, L.L. and Molday, R.S. (2001) Membrane topology of the ATP binding cassette transporter ABCR and its relationship to ABC1 and related ABCA transporters. Identification of N-linked glycosylation sites. *J. Biol. Chem.* 276, 23539–23546.
- [25] Henriksen, U., Fog, J.U., Litman, T. and Gether, U. (2005) Identification of intra- and intermolecular disulfide bridges in the multidrug resistance transporter ABCG2. *J. Biol. Chem.* 280, 36926–36934.
- [26] Haardt, M. and Bremer, E. (1996) Use of *phoA* and *lacZ* fusions to study the membrane topology of ProW, a component of the osmoregulated ProU transport system of *Escherichia coli*. *J. Bacteriol.* 178, 5370–5381.
- [27] Antonov, V.K., Alexandrov, S.L. and Vorotyntseva, T.I. (1976) Reversible association as a possible regulatory mechanism for controlling the activity of the nonspecific leucine-binding protein from *Escherichia coli*. *Adv. Enzyme Regul.* 14, 269–278.
- [28] Rashed, I., Shuman, H. and Boos, W. (1976) The dimer of the *Escherichia coli* galactose-binding protein. *Eur. J. Biochem.* 69, 545–550.
- [29] Richarme, G. (1982) Associative properties of the *Escherichia coli* galactose binding protein and maltose binding protein. *Biochem. Biophys. Res. Commun.* 105, 476–481.
- [30] Richarme, G. (1983) Associative properties of the *Escherichia coli* galactose-binding protein and maltose-binding protein. *Biochim. Biophys. Acta* 748, 99–108.
- [31] Quijcho, F.A. and Ledvina, P.S. (1996) Atomic structure and specificity of bacterial periplasmic receptors for active transport and chemotaxis: variation of common themes. *Mol. Microbiol.* 20, 17–25.
- [32] Alloing, G., De Philip, P. and Claverys, J.-P. (1994) Three highly homologous membrane-bound lipoproteins participate in oligopeptide transport by the Ami system of the Gram-positive *Streptococcus pneumoniae*. *J. Mol. Biol.* 241, 44–58.
- [33] Lin, B., Short, S.A., Eskildsen, M., Klemperer, M.S. and Hu, L.T. (2001) Functional testing of putative oligopeptide permease (Opp) proteins of *Borrelia burgdorferi*: a complementation model in *opp* *Escherichia coli*. *Biochim. Biophys. Acta* 1499, 222–231.
- [34] Doeven, M.K., Kok, J. and Poolman, B. (2005) Specificity and selectivity determinants of peptide transport in *Lactococcus lactis* and other microorganisms. *Mol. Microbiol.* 57, 640–649.
- [35] Borths, E.L., Locher, K.P., Lee, A.T. and Rees, D.C. (2002) The structure of *Escherichia coli* BtuF and binding to its cognate ATP binding cassette transporter. *Proc. Nat. Acad. Sci. USA* 99, 16642–16647.
- [36] Biemans-Oldehinkel, E. and Poolman, B. (2003) On the role of the two extracytoplasmic substrate-binding domains in the ABC transporter OpuA. *EMBO J.* 22, 5983–5993.
- [37] Mao, B., Pear, M.R. and McCammon, J.A. (1982) Hinge-bending in L-arabinose-binding protein. The “Venus’s-flytrap” model. *J. Biol. Chem.* 257, 1131–1133.
- [38] Tame, J.R.H., Murshudov, G.N., Dodson, E.J., Neil, T.K., Dodson, G.G., Higgins, C.F. and Wilkinson, A.J. (1994) The structural basis of sequence-independent peptide binding by OpuA protein. *Science* 264, 1578–1581.
- [39] Nickitenko, A.V., Trakhanov, S. and Quijcho, F.A. (1995) 2 Å resolution structure of DppA, a periplasmic dipeptide transport/chemosensory receptor. *Biochemistry* 34, 16585–16595.
- [40] Hedde, J., Scott, D.J., Unzai, S., Park, S.-Y. and Tame, J.R.H. (2003) Crystal structures of the liganded and unliganded nickel-binding protein NikA from *Escherichia coli*. *J. Biol. Chem.* 278, 50322–50329.
- [41] Karpowich, N.K., Huang, H.H., Smith, P.C. and Hunt, J.F. (2003) Crystal structures of the BtuF periplasmic-binding protein for vitamin B12 suggest a functionally important reduction in protein mobility upon ligand binding. *J. Biol. Chem.* 278, 8429–8434.
- [42] Trakhanov, S., Vyas, N.K., Luecke, H., Kristensen, D.M., Ma, J. and Quijcho, F.A. (2005) Ligand-free and -bound structures of the binding protein (LivJ) of the *Escherichia coli* ABC leucine/isoleucine/valine transport system: trajectory and dynamics of the interdomain rotation and ligand specificity. *Biochemistry* 44, 6597–6608.
- [43] Prossnitz, E., Gee, A. and Ames, G.F.-L. (1989) Reconstitution of the histidine periplasmic transport system in membrane vesicles. Energy coupling and interaction between the binding protein and the membrane complex. *J. Biol. Chem.* 264, 5006–5014.
- [44] Dean, D.A., Hor, L.I., Shuman, H.A. and Nikaido, H. (1992) Interaction between maltose-binding protein and the membrane-associated maltose transporter complex in *Escherichia coli*. *Mol. Microbiol.* 6, 2033–2040.
- [45] Doeven, M.K., Abele, R., Tampé, R. and Poolman, B. (2004) The binding specificity of OpuA determines the selectivity of the oligopeptide ATP-binding cassette transporter. *J. Biol. Chem.* 279, 32301–32307.
- [46] Merino, G., Boos, W., Shuman, H.A. and Bohl, E. (1995) The inhibition of maltose transport by the unliganded form of the maltose-binding protein of *Escherichia coli*: experimental findings and mathematical treatment. *J. Theor. Biol.* 177, 171–179.
- [47] Ames, G.F.-L., Liu, C.E., Joshi, A.K. and Nikaido, K. (1996) Liganded and unliganded receptors interact with equal affinity with the membrane complex of periplasmic permeases, a subfamily of traffic ATPases. *J. Biol. Chem.* 271, 14264–14270.
- [48] Park, Y., Cho, Y.-J., Ahn, T. and Park, C. (1999) Molecular interactions in ribose transport: the binding protein module symmetrically associates with the homodimeric membrane transporter. *EMBO J.* 18, 4149–4156.
- [49] Lanfermeijer, F.C., Picon, A., Konings, W.N. and Poolman, B. (1999) Kinetics and consequences of binding of nona- and dodecapeptides to the oligopeptide binding protein (OpuA) of *Lactococcus lactis*. *Biochemistry* 38, 14440–14450.
- [50] Picon, A., Kunji, E.R.S., Lanfermeijer, F.C., Konings, W.N. and Poolman, B. (2000) Specificity mutants of the binding protein of

- the oligopeptide transport system of *Lactococcus lactis*. J. Bacteriol. 182, 1600–1608.
- [51] Levdikov, V.M., Blagova, E.V., Brannigan, J.A., Wright, L., Vagin, A.A. and Wilkinson, A.J. (2005) The structure of the oligopeptide-binding protein, AppA, from *Bacillus subtilis* in complex with a nonapeptide. J. Mol. Biol. 345, 879–892.
- [52] Schuurman-Wolters, G.K. and Poolman, B. (2005) Substrate specificity and ionic regulation of GlnPQ from *Lactococcus lactis*. J. Biol. Chem. 280, 23785–23790.
- [53] Dassa, E. and Bouige, P. (2001) The ABC of ABCs: a phylogenetic and functional classification of ABC systems in living organisms. Res. Microbiol. 152, 211–229.
- [54] Boyd, D., Manoil, C. and Beckwith, J. (1987) Determinants of membrane protein topology. Proc. Natl. Acad. Sci. USA 84, 8525–8529.
- [55] Ehrmann, M., Boyd, D. and Beckwith, J. (1990) Genetic analysis of membrane protein topology by a sandwich gene fusion approach. Proc. Natl. Acad. Sci. USA 87, 7574–7578.
- [56] Ehrmann, M. and Beckwith, J. (1991) Proper insertion of a complex membrane protein in the absence of its amino-terminal export signal. J. Biol. Chem. 266, 16530–16533.
- [57] Abele, R. and Tampé, R. (1999) Function of the transport complex TAP in cellular immune recognition. Biochim. Biophys. Acta 1461, 405–419.
- [58] Koch, J., Guntrum, R., Heintke, S., Kyritsis, C. and Tampé, R. (2004) Functional dissection of the transmembrane domains of the transporter associated with antigen processing (TAP). J. Biol. Chem. 279, 10142–10147.
- [59] Procko, E., Raghuraman, G., Wiley, D.C., Raghavan, M. and Gaudet, R. (2005) Identification of domain boundaries within the N-termini of TAP1 and TAP2 and their importance in tapasin binding and tapasin-mediated increase in peptide loading of MHC class I. Immunol. Cell Biol. 83, 475–482.
- [60] Bangia, N., Lehner, P.J., Hughes, E.A., Surman, M. and Cresswell, P. (1999) The N-terminal region of tapasin is required to stabilize the MHC class I loading complex. Eur. J. Immunol. 29, 1858–1870.
- [61] Leonhardt, R.M., Keusekotten, K., Bekpen, C. and Knittler, M.R. (2005) Critical role for the tapasin-docking site of TAP2 in the functional integrity of the MHC class I-peptide-loading complex. J. Immunol. 175, 5104–5114.
- [62] Bakos, E., Hegedus, T., Hollo, Z., Welker, E., Tusnády, G.E., Zaman, G.J., Flens, M.J., Varadi, A. and Sarkadi, B. (1996) Membrane topology and glycosylation of the human multidrug resistance-associated protein. J. Biol. Chem. 271, 12322–12326.
- [63] Stride, B.D., Valdimarsson, G., Gerlach, J.H., Wilson, G.M., Cole, S.P. and Deeley, R.G. (1996) Structure and expression of the messenger RNA encoding the murine multidrug resistance protein, an ATP-binding cassette transporter. Mol. Pharmacol. 49, 962–971.
- [64] Tusnády, G.E., Bakos, E., Varadi, A. and Sarkadi, B. (1997) Membrane topology distinguishes a subfamily of the ATP-binding cassette (ABC) transporters. FEBS Lett. 402, 1–3.
- [65] Bakos, E., Evers, R., Szakacs, G., Tusnády, G.E., Welker, E., Szabo, K., de Haas, M., van Deemter, L., Borst, P., Varadi, A. and Sarkadi, B. (1998) Functional multidrug resistance protein (MRP1) lacking the N-terminal transmembrane domain. J. Biol. Chem. 273, 32167–32175.
- [66] Westlake, C.J., Qian, Y.M., Gao, M., Vasa, M., Cole, S.P. and Deeley, R.G. (2003) Identification of the structural and functional boundaries of the multidrug resistance protein 1 cytoplasmic loop 3. Biochemistry 42, 14099–14113.
- [67] Chan, K.W., Zhang, H. and Logothetis, D.E. (2003) N-terminal transmembrane domain of the SUR controls trafficking and gating of  $K_{ATP}$  channel subunits. EMBO J. 22, 3833–3843.
- [68] Babenko, A.P. and Bryan, J. (2003) Sur domains that associate with and gate  $K_{ATP}$  pores define a novel gatekeeper. J. Biol. Chem. 278, 41577–41580.
- [69] Sharma, N., Crane, A., Clement, J.P.t., Gonzalez, G., Babenko, A.P., Bryan, J. and Aguilar-Bryan, L. (1999) The C terminus of SUR1 is required for trafficking of  $K_{ATP}$  channels. J. Biol. Chem. 274, 20628–20632.
- [70] Nies, A.T., Konig, J., Cui, Y., Brom, M., Spring, H. and Keppeler, D. (2002) Structural requirements for the apical sorting of human multidrug resistance protein 2 (ABCC2). Eur. J. Biochem. 269, 1866–1876.
- [71] Fernandez, S.B., Hollo, Z., Kern, A., Bakos, E., Fischer, P.A. and Borst, P. (2002) Role of the N-terminal transmembrane region of the multidrug resistance protein MRP2 in routing to the apical membrane in MDCKII cells. J. Biol. Chem. 277, 31048–31055.
- [72] Westlake, C.J., Cole, S.P. and Deeley, R.G. (2005) Role of the NH<sub>2</sub>-terminal membrane spanning domain of multidrug resistance protein 1/ABCC1 in protein processing and trafficking. Mol. Biol. Cell 16, 2483–2492.
- [73] Romeo, Y., Obis, D., Bouvier, J., Guillot, A., Fourcans, A., Bouvier, I., Gutierrez, C. and Mistou, M.Y. (2003) Osmoregulation in *Lactococcus lactis*: BusR, a transcriptional repressor of the glycine betaine uptake system BusA. Mol. Microbiol. 47, 1135–1147.
- [74] Chen, J., Lu, G., Lin, J., Davidson, A.L. and Quijcho, F.A. (2003) A tweezers-like motions of the ATP-binding cassette dimer in an ABC transport cycle. Mol. Cell 12, 651–661.
- [75] Böhm, A., Diez, J., Diederichs, K., Welte, W. and Boos, W. (2002) Structural model of MalK, the ABC subunit of the maltose transporter of *Escherichia coli*: implications for mal gene regulation, inducer exclusion, and subunit assembly. J. Biol. Chem. 277, 3708–3717.
- [76] Verdon, G., Albers, S.V., Dijkstra, B.W., Driessen, A.J. and Thunnissen, A.M. (2003) Crystal structures of the ATPase subunit of the glucose ABC transporter from *Sulfolobus solfataricus*: nucleotide-free and nucleotide-bound conformations. J. Mol. Biol. 330, 343–358.
- [77] Böhm, A. and Boos, W. (2004) Gene regulation in prokaryotes by subcellular relocalization of transcription factors. Curr. Opin. Microbiol. 7, 51–56.
- [78] Nelson, S.O. and Postma, P.W. (1984) Interactions in vivo between IIIGlc of the phosphoenolpyruvate:sugar phosphotransferase system and the glycerol and maltose uptake systems of *Salmonella typhimurium*. Eur. J. Biochem. 139, 29–34.
- [79] Dean, D.A., Reizer, J., Nikaido, H. and Saier Jr., M.H. (1991) Regulation of the maltose transport system of *Escherichia coli* by the glucose-specific enzyme III of the phosphoenolpyruvate-sugar phosphotransferase system. Characterization of inducer exclusion-resistant mutants and reconstitution of inducer exclusion in proteoliposomes. J. Biol. Chem. 265, 21005–21010.
- [80] Reyes, M. and Shuman, H.A. (1988) Overproduction of MalK protein prevents expression of the *Escherichia coli* mal regulon. J. Bacteriol. 170, 4598–4602.
- [81] Panagiotidis, C.H., Boos, W. and Shuman, H.A. (1998) The ATP-binding cassette subunit of the maltose transporter MalK antagonizes MalT, the activator of the *Escherichia coli* mal regulon. Mol. Microbiol. 30, 535–546.
- [82] Boos, W. and Böhm, A. (2000) Learning new tricks from an old dog: MalT of the *Escherichia coli* maltose system is part of a complex regulatory network. Trends Genet. 16, 404–409.
- [83] Schreiber, V. and Richet, E. (1999) Self-association of the *Escherichia coli* transcription activator MalT in the presence of maltotriose and ATP. J. Biol. Chem. 274, 33220–33226.
- [84] Danot, O. (2001) A complex signaling module governs the activity of MalT, the prototype of an emerging transactivator family. Proc. Natl. Acad. Sci. USA 98, 435–440.
- [85] Schreiber, V., Steegborn, C., Clausen, T., Boos, W. and Richet, E. (2000) A new mechanism for the control of a prokaryotic transcriptional regulator: antagonistic binding of positive and negative effectors. Mol. Microbiol. 35, 65–76.
- [86] Joly, N., Danot, O., Schlegel, A., Boos, W. and Richet, E. (2002) The Aes protein directly controls the activity of MalT, the central transcriptional activator of the *Escherichia coli* maltose regulon. J. Biol. Chem. 277, 16606–16613.
- [87] Kühnau, S., Reyes, M., Sievertsen, A., Shuman, H.A. and Boos, W. (1991) The activities of the *Escherichia coli* MalK protein in maltose transport, regulation, and inducer exclusion can be separated by mutations. J. Bacteriol. 173, 2180–2186.
- [88] van der Heide, T. and Poolman, B. (2000) Osmoregulated ABC-transport system of *Lactococcus lactis* senses water stress via changes in the physical state of the membrane. Proc. Natl. Acad. Sci. USA 97, 7102–7106.



- [89] Biemans-Oldehinkel, E., Poolman, B. (2005) A sensor for intracellular ionic strength. Submitted for publication.
- [90] van der Heide, T., Stuart, M.C. and Poolman, B. (2001) On the osmotic signal and osmosensing mechanism of an ABC transport system for glycine betaine. *EMBO J.* 20, 7022–7032.
- [91] Rich, D.P., Gregory, R.J., Anderson, M.P., Manavalan, P., Smith, A.E. and Welsh, M.J. (1991) Effect of deleting the R domain on CFTR-generated chloride channels. *Science* 253, 205–207.
- [92] Gadsby, D.C. and Nairn, A.C. (1999) Control of CFTR channel gating by phosphorylation and nucleotide hydrolysis. *Physiol. Rev.* 79, S77–S107.
- [93] Dahan, D., Evagelidis, A., Hanrahan, J.W., Hinkson, D.A., Jia, Y., Luo, J. and Zhu, T. (2001) Regulation of the CFTR channel by phosphorylation. *Pflügers Arch.* 443 (Suppl. 1), S92–S96.
- [94] Rich, D.P., Berger, H.A., Cheng, S.H., Travis, S.M., Saxena, M., Smith, A.E. and Welsh, M.J. (1993) Regulation of the cystic fibrosis transmembrane conductance regulator Cl-channel by negative charge in the R domain. *J. Biol. Chem.* 268, 20259–20267.
- [95] Csanády, L., Chan, K.W., Seto-Young, D., Kopsco, D.C., Nairn, A.C. and Gadsby, D.C. (2000) Severed channels probe regulation of gating of cystic fibrosis transmembrane conductance regulator by its cytoplasmic domains. *J. Gen. Physiol.* 116, 477–500.
- [96] Balduresson, O., Ostedgaard, L.S., Rokhlina, T., Cotten, J.F. and Welsh, M.J. (2001) Cystic fibrosis transmembrane conductance regulator Cl-channels with R domain deletions and translocations show phosphorylation-dependent and -independent activity. *J. Biol. Chem.* 276, 1904–1910.
- [97] Chappe, V., Irvine, T., Liao, J., Evagelidis, A. and Hanrahan, J.W. (2005) Phosphorylation of CFTR by PKA promotes binding of the regulatory domain. *EMBO J.* 24, 2730–2740.
- [98] Roosbeek, S., Peelman, F., Verhee, A., Labeur, C., Caster, H., Lensink, M.F., Cirulli, C., Grooten, J., Cochet, C., Vandekerckhove, J., Amoresano, A., Chimini, G., Tavernier, J. and Rosseneu, M. (2004) Phosphorylation by protein kinase CK2 modulates the activity of the ATP binding cassette A1 transporter. *J. Biol. Chem.* 279, 37779–37788.
- [99] Fu, J. and Kirk, K.L. (2001) Cysteine substitutions reveal dual functions of the amino-terminal tail in cystic fibrosis transmembrane conductance regulator channel gating. *J. Biol. Chem.* 276, 35660–35668.
- [100] Naren, A.P., Cormet-Boyaka, E., Fu, J., Villain, M., Blalock, J.E., Quick, M.W. and Kirk, K.L. (1999) CFTR chloride channel regulation by an interdomain interaction. *Science* 286, 544–548.
- [101] Li, C. and Naren, A.P. (2005) Macromolecular complexes of cystic fibrosis transmembrane conductance regulator and its interacting partners. *Pharmacol. Ther.* 108, 208–223.
- [102] Wang, S., Yue, H., Derin, R.B., Giggino, W.B. and Li, M. (2000) Accessory protein facilitated CFTR–CFTR interaction, a molecular mechanism to potentiate the chloride channel activity. *Cell* 103, 169–179.
- [103] Naren, A.P., Cobb, B., Li, C., Roy, K., Nelson, D., Heda, G.D., Liao, J., Kirk, K.L., Sorscher, E.J., Hanrahan, J. and Clancy, J.P. (2003) A macromolecular complex of beta 2 adrenergic receptor, CFTR, and ezrin/radixin/moesin-binding phosphoprotein 50 is regulated by PKA. *Proc. Natl. Acad. Sci. USA* 100, 342–346.
- [104] Li, J., Dai, Z., Jana, D., Callaway, D.J.E. and Bu, Z. (2005) Ezrin controls the macromolecular complexes formed between an adapter protein Na<sup>+</sup>/H<sup>+</sup> exchanger regulatory factor and the cystic fibrosis transmembrane conductance regulator. *J. Biol. Chem.* 280, 37634–37643.
- [105] Raghuram, V., Hormuth, H. and Foskett, J.K. (2003) A kinase-regulated mechanism controls CFTR channel gating by disrupting bivalent PDZ domain interactions. *Proc. Natl. Acad. Sci. USA* 100, 9620–9625.
- [106] Fu, J., Ji, H.L., Naren, A.P. and Kirk, K.L. (2001) A cluster of negative charges at the amino terminal tail of CFTR regulates ATP-dependent channel gating. *J. Physiol.* 536, 459–470.
- [107] Bakos, E., Evers, R., Calenda, G., Tusnády, G.E., Szakacs, G., Varadi, A. and Sarkadi, B. (2000) Characterization of the amino-terminal regions in the human multidrug resistance protein (MRP1). *J. Cell Sci.* 113, 4451–4461.
- [108] Babenko, A.P. and Bryan, J. (2002) SUR-dependent modulation of K<sub>ATP</sub> channels by an N-terminal K<sub>IR</sub> 6.2 peptide. Defining intersubunit gating interactions. *J. Biol. Chem.* 277, 43997–44004.
- [109] Bryan, J., Vila-Carriles, W.H., Zhao, G., Babenko, A.P. and Aguilar-Bryan, L. (2004) Toward linking structure with function in ATP-sensitive K<sup>+</sup> channels. *Diabetes* 53 (Suppl. 3), S104–S112.
- [110] Mikhailov, M.V., Mikhailova, E.A. and Ashcroft, S.J. (2001) Molecular structure of the glibenclamide binding site of the beta-cell K(ATP) channel. *FEBS Lett.* 499, 154–160.
- [111] Dhar-Chowdhury, P., Harrell, M.D., Han, S.Y., Jankowska, D., Parachuru, L., Morrissey, A., Srivastava, S., Liu, W., Malester, B., Yoshida, H. and Coetzee, W.A. (2005) The glycolytic enzymes, glyceraldehyde-3-phosphate dehydrogenase, triose-phosphate isomerase, and pyruvate kinase are components of the K<sub>ATP</sub> channel macromolecular complex and regulate its function. *J. Biol. Chem.* 280, 38464–38470.
- [112] Gao, M., Yamazaki, M., Loe, D.W., Westlake, C.J., Grant, C.E., Cole, S.P. and Deeley, R.G. (1998) Multidrug resistance protein. Identification of regions required for active transport of leukotriene C<sub>4</sub>. *J. Biol. Chem.* 273, 10733–10740.
- [113] Qian, Y.M., Qiu, W., Gao, M., Westlake, C.J., Cole, S.P. and Deeley, R.G. (2001) Characterization of binding of leukotriene C<sub>4</sub> by human multidrug resistance protein 1: evidence of differential interactions with NH<sub>2</sub>- and COOH-proximal halves of the protein. *J. Biol. Chem.* 276, 38636–38644.
- [114] Ren, X.Q., Furukawa, T., Aoki, S., Nakajima, T., Sumizawa, T., Haraguchi, M., Chen, Z.S., Kobayashi, M. and Akiyama, S. (2001) Glutathione-dependent binding of a photoaffinity analog of agosterol A to the C-terminal half of human multidrug resistance protein. *J. Biol. Chem.* 276, 23197–23206.
- [115] Haimeur, A., Conseil, G., Deeley, R.G. and Cole, S.P.C. (2004) The MRP-related and BCRP/ABCG2 multidrug resistance proteins: Biology, substrate specificity and regulation. *Curr. Drug Metab.* 5, 21–53.
- [116] Fath, M.J. and Kolter, R. (1993) ABC transporters: bacterial exporters. *Microbiol. Rev.* 57, 995–1017.
- [117] Holo, H., Nilssen, O. and Nes, I.F. (1991) Lactococcin A, a new bacteriocin from *Lactococcus lactis* subsp. *cremoris*: isolation and characterization of the protein and its gene. *J. Bacteriol.* 173, 3879–3887.
- [118] Hui, F.M. and Morrison, D.A. (1991) Genetic transformation in *Streptococcus pneumoniae*: nucleotide sequence analysis shows comA, a gene required for competence induction, to be a member of the bacterial ATP-dependent transport protein family. *J. Bacteriol.* 173, 372–381.
- [119] Juranka, P., Zhang, F., Kulpa, J., Endicott, J., Blight, M., Holland, I.B. and Ling, V. (1992) Characterization of the hemolysin transporter, HlyB, using an epitope insertion. *J. Biol. Chem.* 267, 3764–3770.
- [120] Dirix, G., Monsieurs, P., Marchal, K., Vanderleyden, J. and Michiels, J. (2004) Screening genomes of Gram-positive bacteria for double-glycine-motif-containing peptides. *Microbiology* 110, 617–620.
- [121] Havarstein, L.S., Holo, H. and Nes, I.F. (1994) The leader peptide of colicin V shares consensus sequences with leader peptides that are common among peptide bacteriocins produced by gram-positive bacteria. *Microbiology* 140, 2383–2389.
- [122] Havarstein, L.S., Diep, D.B. and Nes, I.F. (1995) A family of bacteriocin ABC transporters carry out proteolytic processing of their substrates concomitant with export. *Mol. Microbiol.* 16, 229–240.
- [123] Wu, K.-H. and Tai, P.C. (2004) Cys32 and His105 are the critical residues for the calcium-dependent cysteine proteolytic activity of CvaB, an ATP-binding cassette transporter. *J. Biol. Chem.* 279, 901–909.
- [124] Franke, C.M., Tiemersma, J., Venema, G. and Kok, J. (1999) Membrane topology of the lactococcal bacteriocin ATP-binding cassette transporter protein LcnC. Involvement of LcnC in lactococcal maturation. *J. Biol. Chem.* 274, 8484–8490.
- [125] Chervaux, C. and Holland, I.B. (1996) Random and directed mutagenesis to elucidate the functional importance of helix II and F-989 in the C-terminal secretion signal of *Escherichia coli* hemolysin. *J. Bacteriol.* 178, 1232–1236.

- [126] Benabdelhak, H., Kiontke, S., Horn, C., Ernst, R., Blight, M.A., Holland, I.B. and Schmitt, L. (2003) A specific interaction between the NBD of the ABC transporter HlyB and a C-terminal fragment of its transport substrate haemolysin A. *J. Mol. Biol.* 327, 1169–1179.
- [127] Holland, I.B., Cole, S.P.C., Kuchler, K. and Higgins, C.F. (2003) ABC Proteins; From Bacteria to Man, Elsevier Science Ltd, London.
- [128] Richarme, G. and Caldas, T.D. (1997) Chaperone properties of the bacterial periplasmic substrate-binding proteins. *J. Biol. Chem.* 272, 15607–15612.
- [129] Kokoeva, M.V., Storch, K-F., Klein, C. and Oesterhelt, D. (2002) A novel mode of sensory transduction in archaea: binding protein-mediated chemotaxis towards osmoprotectants and amino acids. *EMBO J.* 21, 2312–2322.
- [130] Driessen, A.J.M., Rosen, B.P. and Konings, W.N. (2000) Diversity of transport mechanisms: common structural principles. *Trends Biochem. Sci.* 25, 397–401.
- [131] Madden, D. (2002) The structure and function of glutamate receptor ion channels. *Nat. Rev. Neurosci.* 3, 91–101.
- [132] Gouaux, E. (2003) Structure and function of AMPA receptors. *J. Physiol.* 554.2, 249–253.
- [133] Kerppola, R.E. and Ames, G.F-L. (1992) Topology of the hydrophobic membrane-bound components of the histidine periplasmic permease. Comparison with other members of the family. *J. Biol. Chem.* 267, 2329–2336.
- [134] Pearce, S.R., Mimmack, M.L., Gallagher, M.P., Gileadi, U., Hyde, S.C. and Higgins, C.F. (1992) Membrane topology of the integral membrane components, OppB and OppC, of the oligopeptide permease of *Salmonella typhimurium*. *Mol. Microbiol.* 6, 47–57.
- [135] Dassa, E. and Muir, S. (1993) Membrane topology of MalG, an inner membrane protein from the maltose transport system of *Escherichia coli*. *Mol. Microbiol.* 7, 29–38.
- [136] Stewart, J.B. and Hermodson, M.A. (2003) Topology of RbsC, the membrane component of the *Escherichia coli* ribose transporter. *J. Bacteriol.* 185, 5234–5239.
- [137] Groeger, W. and Köster, W. (1998) Transmembrane topology of the two FhuB domains representing the hydrophobic components of bacterial ABC transporters involved in the uptake of siderophores, haem and vitamin B<sup>12</sup>. *Microbiology* 144, 2759–2769.
- [138] Van veen, H.W., Venema, K., Bolhuis, H., Oussenko, I., Kok, J., Poolman, B., Driessen, A.J.M. and Konings, W.N. (1996) Multidrug resistance mediated by a bacterial homolog of the human multidrug transporter MDR1. *Proc. Natl. Acad. Sci. USA* 93, 10668–10672.
- [139] Steinfeld, E., Orelle, C., Fantino, J-R., Dalmas, O., Rigaud, J-L., Denizot, F., Di Pietro, A. and Jault, J-M. (2004) Characterization of YvxC (BmrA), a multidrug ABC transporter constitutively expressed in *Bacillus subtilis*. *Biochemistry* 43, 7491–7502.
- [140] Cruz-Ramos, H., Cook, G.M., Wu, G., Cleeter, M.W. and Poole, R.K. (2004) Membrane topology and mutational analysis of *Escherichia coli* CydDC, an ABC type cysteine exporter required for cytochrome assembly. *Microbiology* 150, 3415–3427.
- [141] Wang, R., Seror, S.J., Blight, M., Pratt, J.M., Broome-Smith, J.K. and Holland, I.B. (1991) Analysis of the membrane organization of an *Escherichia coli* protein translocator, HlyB, a member of a large family of prokaryote and eukaryote surface transport proteins. *J. Mol. Biol.* 217, 441–454.
- [142] Gentschev, I. and Goebel, W. (1992) Topological and functional studies on HlyB of *Escherichia coli*. *Mol. Gen. Genet.* 232, 40–48.
- [143] Loo, T.W. and Clarke, D.M. (1999) Determining the structure and mechanism of the human multidrug resistance P-glycoprotein using cysteine-scanning mutagenesis and thiol-modification techniques. *Biochim. Biophys. Acta* 1461, 315–325.
- [144] Leslie, E.M., Deeley, R.G. and Cole, S.P.C. (2005) Multidrug resistance proteins: role of P-glycoprotein, MRP1, MRP2, and BCRP (ABCG2) in tissue defense. *Toxicol. Appl. Pharmacol.* 204, 216–237.
- [145] Conti, L.R., Radeke, C.M., Shyng, S-L. and Vandenberg, C.A. (2001) Transmembrane topology of the sulfonurea receptor SUR1. *J. Biol. Chem.* 276, 41270–41278.
- [146] Sheppard, D.N. and Welsh, M.J. (1999) Structure and function of the CFTR chloride channel. *Physiol. Rev.* 79 (1 Suppl.), S23–S45.
- [147] Geller, D., Taglicht, D., Edgar, R., Tam, A., Pines, O., Michaelis, S. and Bibi, E. (1996) Comparative topology studies in *Saccharomyces cerevisiae* and in *Escherichia coli*. The N-terminal half of the yeast ABC protein Ste6. *J. Biol. Chem.* 271, 13746–13753.
- [148] Park, Y. and Park, C. (1999) Topology of RbsC, a membrane component of the ribose transporter, belonging to the AraH superfamily. *J. Bacteriol.* 181, 1039–1042.