University of Groningen

(p-phenylenedimethylene)diammonium dichloride

Arkenbout, Anne H.; Meetsma, Auke; Palstra, Thomas T. M.

Published in:
Acta Crystallographica Section E-Structure Reports Online

DOI:
[10.1107/S1600536807002619](https://doi.org/10.1107/S1600536807002619)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2007

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Arkenbout, A. H., Meetsma, A., & Palstra, T. T. M. (2007). (p-phenylenedimethylene)diammonium dichloride. *Acta Crystallographica Section E-Structure Reports Online*, 63(2), O869-O870.
<https://doi.org/10.1107/S1600536807002619>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

(*p*-Phenylenedimethylene)diammonium dichloride**Anne H. Arkenbout, Auke Meetsma* and Thomas T. M. Palstra**

Solid State Chemistry Laboratory, Materials Science Center, University of Groningen, Nijenborgh 4, NL-9747 AG Groningen, The Netherlands

Correspondence e-mail: a.meetsma@rug.nl

Key indicators

Single-crystal X-ray study

 $T = 100\text{ K}$ Mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$ R factor = 0.038 wR factor = 0.092

Data-to-parameter ratio = 11.9

For details of how these key indicators were automatically derived from the article, see <http://journals.iucr.org/e>.

In the title compound, $\text{C}_8\text{H}_{14}\text{N}_2^{2+} \cdot 2\text{Cl}^-$, the cation possesses a crystallographically imposed center of symmetry. The crystal packing is stabilized by intermolecular $\text{N}-\text{H} \cdots \text{Cl}$ hydrogen bonds and π - π stacking interactions.

Received 8 January 2007

Accepted 17 January 2007

Comment

There is an interest in novel semiconductors that are more versatile than the current Si-based devices. Research has been carried out on organic semiconductors such as pentacene (Jurchescu *et al.*, 2004; de Boer *et al.*, 2004), which have the advantage that they are flexible and possibly inkjet printable. However, due to the weak van der Waals bonding, these materials have a limited carrier mobility. The properties of the organics can be improved by combining them with inorganics in hybrid materials (Mitzi, 1999). We present here the crystal structure of the title compound, (I), which we use as the starting point for the development of new hybrids.

In (I) (Fig. 1), all bond lengths and angles show normal values (Allen *et al.*, 1987). The cations lie on centers of symmetry and form stacks parallel to the a axis, with a short

Figure 1

The molecular structure of (I), showing the atomic labeling and 50% probability displacement ellipsoids [symmetry code: (a) $1 - x, -y, 1 - z$].

C2...C4^{iv} distance of 3.496 (2) Å [symmetry code: (iv) $x + 1, y, z$], suggesting π - π stacking interactions. Intermolecular N—H...Cl hydrogen bonds (Table 1) contribute to the stability of the crystal packing (Fig. 2).

Experimental

α, α' -Diamino-*p*-xylene (2 g, Fluka, purum > 98%) was dissolved in water (7 ml). Subsequently, saturated (37%) HCl (5 ml, Merck) was added with stirring and cooling on a water bath. The mixture was filtered and the resulting white powder was washed with water and dried in air. Small colorless crystals were obtained by recrystallization from a saturated aqueous solution at 343 K, which was slowly cooled to room temperature.

Crystal data

$C_8H_{14}N_2^{2+} \cdot 2Cl^-$	$V = 247.18 (8) \text{ \AA}^3$
$M_r = 209.12$	$Z = 1$
Triclinic, $P\bar{1}$	$D_x = 1.405 \text{ Mg m}^{-3}$
$a = 4.3496 (8) \text{ \AA}$	Mo $K\alpha$ radiation
$b = 5.809 (1) \text{ \AA}$	$\mu = 0.61 \text{ mm}^{-1}$
$c = 10.197 (2) \text{ \AA}$	$T = 100 (1) \text{ K}$
$\alpha = 101.836 (3)^\circ$	Platelet, colorless
$\beta = 99.727 (3)^\circ$	$0.17 \times 0.08 \times 0.03 \text{ mm}$
$\gamma = 93.800 (3)^\circ$	

Data collection

Bruker SMART APEX CCD area-detector diffractometer	1835 measured reflections
φ and ω scans	986 independent reflections
Absorption correction: multi-scan (SADABS; Sheldrick, 2006)	881 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.900, T_{\max} = 0.982$	$R_{\text{int}} = 0.046$
	$\theta_{\max} = 26.4^\circ$

Refinement

Refinement on F^2	$w = 1/[\sigma^2(F_o^2) + (0.0368P)^2 + 0.28P]$
$R[F^2 > 2\sigma(F^2)] = 0.038$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.092$	$(\Delta/\sigma)_{\max} < 0.001$
$S = 1.06$	$\Delta\rho_{\max} = 0.36 \text{ e \AA}^{-3}$
986 reflections	$\Delta\rho_{\min} = -0.25 \text{ e \AA}^{-3}$
83 parameters	
All H-atom parameters refined	

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N—H5...Cl ⁱ	0.92 (3)	2.32 (3)	3.190 (2)	158 (3)
N—H5'...Cl ⁱⁱⁱ	0.92 (3)	2.32 (3)	3.235 (2)	176 (2)
N—H5''...Cl ⁱⁱⁱ	0.93 (4)	2.28 (4)	3.199 (2)	175 (4)

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + 1, -y, -z + 1$; (iii) $x, y, z + 1$.

Figure 2

A portion of the crystal packing, showing the N—H...Cl hydrogen bonds as dashed lines.

All H atoms were located in a difference Fourier map and refined with isotropic displacement parameters [$O-H = 0.92 (3)–0.98 (3) \text{ \AA}$].

Data collection: SMART (Bruker, 2006); cell refinement: SAINT-Plus (Bruker, 2006); data reduction: SAINT-Plus; program(s) used to solve structure: DIRDIF99 (Beurskens *et al.*, 1999); program(s) used to refine structure: SHELXL97 (Sheldrick, 1997); molecular graphics: PLUTO (Meetsma, 2006) and PLATON (Spek, 2003); software used to prepare material for publication: PLATON.

The authors thank Oana D. Jurchescu for useful discussions. This project was performed with financial support from NanoNed.

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Beurskens, P. T., Beurskens, G., de Gelder, R., García-Granda, S., Gould, R. O., Israël, R. & Smits, J. M. M. (1999). *The DIRDIF99 Program System*. Technical Report of the Crystallography Laboratory, University of Nijmegen, The Netherlands.
- Boer, R. W. I. de, Gershenson, M. E., Morpurgo, A. F. & Podzorov, V. (2004). *Phys. Stat. Sol.* **201**, 1302–1331.
- Bruker, (2006). SMART, and SAINT-Plus. Bruker AXS Inc., Madison, Wisconsin, USA.
- Jurchescu, O. D., Baas, J. & Palstra, T. T. M. (2004). *Appl. Phys. Lett.* **84**, 3061–3063.
- Meetsma, A. (2006). PLUTO. Extended version. University of Groningen, The Netherlands. Unpublished.
- Mitzi, D. B. (1999). *Prog. Inorg. Chem.* **48**, 1–121.
- Sheldrick, G. M. (1997). SHELXL97. University of Göttingen, Germany.
- Sheldrick, G. M. (2006). SADABS. Version 2.10. University of Göttingen, Germany.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.