University of Groningen

ScbA from *Streptomyces coelicolor* A3(2) has homology to fatty acid synthases and is able to synthesize-butyrolactones

Hsiao, Nai-Hua; Soeding, Johannes; Linke, Dirk; Lange, Corinna; Hertweck, Christian; Wohlleben, Wolfgang; Takano, Eriko

Published in:
 Microbiology-Sgm

DOI:
[10.1099/mic.0.2006/004432-0](https://doi.org/10.1099/mic.0.2006/004432-0)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2007

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Hsiao, N-H., Soeding, J., Linke, D., Lange, C., Hertweck, C., Wohlleben, W., & Takano, E. (2007). ScbA from *Streptomyces coelicolor* A3(2) has homology to fatty acid synthases and is able to synthesize-butyrolactones. *Microbiology-Sgm*, 153, 1394-1404. <https://doi.org/10.1099/mic.0.2006/004432-0>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supplementary figures and mutant construct descriptions

ScbA from *Streptomyces coelicolor* A3(2) has homology to fatty acid synthases and is able to synthesise γ -butyrolactones, by N.-H. Hsiao, J. Söding, D. Linke, C. Lange, C. Hertweck, W. Wohlleben and E. Takano

Microbiology vol. **153**, part 5, pp. 1394-1404

MS data for M145::pSET152 (Fig. 1A, B) and M751::pIJ6147 (Fig. 2A, B) where SCB1 production was observed.

Fig. S1. A. (a) Base peak chromatogram of M145::pSET152. (b) Extracted ion chromatogram (SIM) for m/z 245 ($[244+H]^+$) of sample M145::pSET152. (c) Base peak chromatogram of SCB1 standard. (B) (a) Mass spectrum of the peak at retention time 14.5 min of sample M145::pSET152. (b) Mass spectrum of SCB1 standard. (c) MS/MS spectrum of the m/z 245 hydrogen adduct, peak at 14.5 min of sample M145::pSET152. (d) MS/MS spectrum of the m/z 245 hydrogen adduct, SCB1 standard.

Fig. S2. A. (a) Base peak chromatogram of sample M751::pIJ6147. (b) Extracted ion chromatogram (SIM) for m/z 245 ($[244+H]^+$) of sample M751::pIJ6147. (c) Base peak chromatogram of SCB1 standard. B. (a) Mass spectrum of the peak at retention time 14.5 min of sample M751::pIJ6147. (b) Mass spectrum of SCB1 standard.