University of Groningen

Synthesis and structures of scandium and lutetium benzyl complexes

Meyer, N.; Roesky, Peter W.; Bambirra, Sergio; Meetsma, Auke; Hessen, Bart; Saliu, Kuburat; Takats, Josef

Published in:
 Organometallics

DOI:
[10.1021/om701171y](https://doi.org/10.1021/om701171y)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2008

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Meyer, N., Roesky, P. W., Bambirra, S., Meetsma, A., Hessen, B., Saliu, K., & Takats, J. (2008). Synthesis and structures of scandium and lutetium benzyl complexes. *Organometallics*, 27(7), 1501-1505.
<https://doi.org/10.1021/om701171y>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Correction to Synthesis and Structures of Scandium and Lutetium Benzyl Complexes

Nils Meyer, Peter W. Roesky,* Sergio Bambirra, Auke Meetsma, Bart Hessen,* Kuburat Saliu, and Josef Takats*

Organometallics 2008, 27 (7), 1501–1505. DOI: 10.1021/om701171y

In the article, the broad signals observed in the ^1H NMR spectrum of $[\text{Lu}(\text{CH}_2\text{Ph})_3(\text{THF})_3]$, obtained in Edmonton, in $\text{THF}-d_8$ at RT and the two sets of signals at LT were attributed to some fluxional behavior. Subsequently, KS discovered¹ that the actual reason for this was the presence of the ate-complex, $\text{K}[\text{Lu}(\text{CH}_2\text{Ph})_4(\text{THF})_x]$, in the sample. The Edmonton synthesis was repeated several times, but in each case, it was accompanied by the formation of variable amounts of the ate-complex contamination, as determined by ^1H NMR spectroscopy in $\text{THF}-d_8$. Although crystals of $[\text{Lu}(\text{CH}_2\text{Ph})_3(\text{THF})_3]$ were obtained from such preparations, in our hands, pure neutral tribenzyl was best obtained by trituration/extraction of the crude product with toluene,

which gave $[\text{Lu}(\text{CH}_2\text{Ph})_3(\text{THF})_2]$ with two coordinated THF molecules (structure at hand) but with loss of yield. Figure 1 shows the ^1H NMR spectra of pure $[\text{Lu}(\text{CH}_2\text{Ph})_3(\text{THF})_3]$, the ate-complex, and mixture of the two in $\text{THF}-d_8$ at LT; the ^1H NMR spectra of the former two complexes are temperature-invariant except for minor changes in chemical shifts, whereas that of the mixture shows broad signals at RT due to intermolecular exchange.

JT regrets the error and wishes to emphasize that the problem with ate-complex formation was only observed in the Edmonton synthetic protocol and workup procedure. Alternative, large-scale synthesis of $[\text{Ln}(\text{CH}_2\text{Ph})_3(\text{THF})_3]$ ² and most recently including that of $[\text{Lu}(\text{CH}_2\text{Ph})_3(\text{THF})_3]$ has appeared.³

REFERENCES

- (1) Saliu, K. Scorpionate-Anchored Lanthanide Complexes. Ph.D. Thesis, University of Alberta, 2009.
- (2) Wooles, A. J.; Mills, D. P.; Lewis, W.; Blake, A. J.; Liddle, S. T. *Dalton Trans.* 2010, 39, 500–510.
- (3) Huang, W.; Upton, B. M.; Khan, S. I.; Diaconescu, P. L. *Organometallics* 2013, 32, 1379–1386 (see Supporting Information); 2275 (erratum).

Figure 1. ^1H NMR spectra of (a) mixture of $[\text{Lu}(\text{CH}_2\text{Ph})_3(\text{THF})_3]/\text{K}[\text{Lu}(\text{CH}_2\text{Ph})_4(\text{THF})_x]$ (shown primed), (b) $[\text{Lu}(\text{CH}_2\text{Ph})_3(\text{THF})_3]$, and (c) $\text{K}[\text{Lu}(\text{CH}_2\text{Ph})_4(\text{THF})_x]$ in $\text{THF}-d_8$ at -80°C .

Published: May 28, 2013