University of Groningen

Seasonal Redistribution of Immune Function in a Migrant Shorebird

Buehler, Deborah M.; Piersma, Theunis; Matson, Kevin D.; Tieleman, B. Irene

Published in:
American Naturalist

DOI:
[10.1086/592865](https://doi.org/10.1086/592865)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2008

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Buehler, D. M., Piersma, T., Matson, K. D., & Tieleman, B. I. (2008). Seasonal Redistribution of Immune Function in a Migrant Shorebird: Annual-Cycle Effects Override Adjustments to Thermal Regime. *American Naturalist*, 172(6), 783-796. <https://doi.org/10.1086/592865>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Seasonal Redistribution of Immune Function in a Migrant Shorebird: Annual-Cycle Effects Override Adjustments to Thermal Regime

Deborah M. Buehler,^{1,*} Theunis Piersma,^{1,2,†} Kevin Matson,^{1,‡} and B. Irene Tieleman^{1,§}

1. Animal Ecology Group, Centre for Ecological and Evolutionary Studies, University of Groningen, P.O. Box 14, 9750 AA, Haren, The Netherlands;

2. Department of Marine Ecology and Evolution, Royal Netherlands Institute for Sea Research (NIOZ), P.O. Box 59, 1790 AB Den Burg, Texel, The Netherlands

Submitted March 5, 2008; Accepted June 18, 2008;
Electronically published October 31, 2008

Online enhancement: appendix.

ABSTRACT: Throughout the annual cycle, demands on competing physiological systems change, and animals must allocate resources to maximize fitness. Immune function is one such system and is important for survival. Yet detailed empirical data tracking immune function over the entire annual cycle are lacking for most wild animals. We measured constitutive immune indices once a month for a year on captive red knots (*Calidris canutus*). We also examined temperature as an environmental contributor to immune variation by manipulating ambient temperature to vary energy expenditure. To identify relationships among immune indices, we performed principal-component analysis. We found significant repeatability in immune indices over the annual cycle and covariation of immune indices within and among individuals. This covariation suggests immune strategies as individual traits among individuals and the use of different immune strategies during different annual-cycle stages within individuals. Over the annual cycle, both higher-cost phagocyte-based immunity and lower-cost lymphocyte-based immunity were high during mass change, but there was a clear shift toward lower-cost lymphocyte-based immunity during peak molt. Experimental manipulation of temperature had little effect on annual variation in immune function. This suggests that other environmental

factors, such as food availability and disease, should also be examined in the future.

Keywords: annual cycle, immune function, immune strategies, temperature, red knot, *Calidris canutus*.

Throughout the annual cycle, demands on competing physiological systems change, and animals must allocate resources to maximize fitness (King 1974). Immune function represents one such physiological system, and it contributes to host survival by limiting infection and performing self-maintenance duties such as clearing apoptotic cells. Given its importance, one might expect animals to maintain strong immune defense throughout the year. However, variability in immune defense is widespread (reviewed by Martin et al. [2008]), and hypotheses about why this variation occurs include endogenous annual cycles, changes in disease threat, and immunoenhancement in anticipation of disease threat based on changes in environmental cues (Nelson et al. 2002). Variation may also occur because immune defense comes with costs as well as benefits. Building an immune system is developmentally costly, and maintaining and using the system have energetic and immunopathology costs (Råberg et al. 1998; Schmid-Hempel and Ebert 2003; Klasing 2004). These costs have led to predictions about trade-offs between immune function and other activities (e.g., reproduction, migration, molt, thermoregulation; Sheldon and Verhulst 1996; Lochmiller and Deerenberg 2000; Norris and Evans 2000). However, evaluating these hypotheses is difficult without detailed empirical data (Hasselquist 2007).

Empirical data tracking immune variation over the annual cycle exist for humans and small mammals (Nelson et al. 2002) but are lacking for other animals. Avian studies, for example, have included comparisons between pairs of life-cycle stages (e.g., breeding vs. nonbreeding or molt vs. nonmolt; see Martin et al. 2008 for a review) but have not detailed variation over the entire annual cycle. Furthermore, baseline data collected under controlled environ-

* Corresponding author; e-mail: d.m.buehler@rug.nl.

† E-mail: theunis@nioz.nl.

‡ E-mail: k.d.matson@rug.nl.

§ E-mail: b.i.tieleman@rug.nl.

mental, nutritional, and disease risk conditions are necessary to interpret data taken across seasons in the wild. In small mammals, such baseline data, coupled with data from the wild, show that complex physiology-environment interactions in the wild can mask underlying seasonal anticipatory patterns (Nelson et al. 2002).

In addition to baseline data, controlled studies that manipulate environmental factors are needed to tease apart the relative importance of contributors to immune variation. Many environmental factors change over the annual cycle (e.g., temperature, disease risk, photoperiod, and food availability), and two, ambient temperature and photoperiod, have been linked to annual variation in immune function in mammals (Nelson et al. 2002). We focus on ambient temperature, which may also influence immune function in birds, especially species wintering in temperate regions where temperature greatly affects energy expenditure (Wiersma and Piersma 1994). If immune function is traded off when birds need to invest more in thermoregulation, then we predict that birds living in colder temperatures should show reduced immune function or a shift away from higher-cost immunity (*sensu* Lee 2006, i.e., less phagocytosis/inflammation).

Another environmental contributor to annual variation in immune function is disease risk. Seasonal fluctuations in diseases have been detected in humans, mammals, and birds (Nelson et al. 2002) and are likely to affect investment in immune defense, but in the wild these patterns are extremely complex (Altizer et al. 2006). Even in captivity, disease risk is not easily manipulated; however, in some species it can be somewhat controlled by cleaning regimes (Buehler et al. 2008). If disease risk in captivity is relatively stable, then patterns of immune function observed in captivity may be interpreted in the context of anticipatory cycles, in species where annual patterns of disease risk can be predicted.

Red knots (*Calidris canutus islandica*) are medium-sized (100–200-g) long-distance migrant shorebirds with well-studied annual cycles (Piersma 2007). In this subspecies, disease risks are predicted to be highest during migration (Buehler and Piersma 2008), when the birds pass through a variety of environments where they might encounter novel pathogens (Møller and Erritzøe 1998). During breeding in the Arctic, knots are widely dispersed, decreasing the risk of disease transmission. Little is known about disease risk during fall migration, molt, and wintering; however, it is likely lower than during spring migration (Buehler and Piersma 2008). Fall migration is less synchronous than spring migration, and birds are aggregated in smaller flocks (Battley et al. 2004). Given appropriate photoperiod cues, captive red knots exhibit premigratory fattening, mass loss, and molt cycles comparable to those of free-living birds (Piersma et al. 1995; Jenni-

Eiermann et al. 2002). Furthermore, because captive knots do not migrate, anticipatory cycles corresponding to disease risk, which might be masked by extreme physical activity in migrating birds, may also be detected.

The immune system can be divided along an innate (nonspecific)-acquired (specific) axis and a constitutive (noninduced)-induced axis (Schmid-Hempel and Ebert 2003). Immune defense may shift between or even within these axes to provide optimal immune strategies in different situations (Schmid-Hempel and Ebert 2003; Lee 2006; Martin et al. 2007). We chose constitutive immunity to examine resource allocation over the annual cycle because it is effective at controlling multiple pathogen types and responds immediately to threats, making it an evolutionarily relevant first line of defense. Furthermore, mediators of constitutive immunity must be maintained even when not in use, generating costs that may be important in physiological trade-offs during the annual cycle (Schmid-Hempel and Ebert 2003; Martin et al. 2008). Finally, because a response is not induced and immunological memory is not stimulated, repeated measures of individuals throughout the annual cycle can be made. We quantified three categories of constitutive immune indices covering a range of protective functions. The first is microbial killing ability, which measures the functional capacity of blood to limit microbial infection (Tieleman et al. 2005; Millet et al. 2007). The second consists of leukocyte concentrations, which provide information on circulating immune cells, and differential leukocyte counts, which are useful in multivariate analysis in terms of their relationship to functional measures of immunity, such as microbe killing. Heterophils and eosinophils mediate innate immunity against novel pathogens and are important phagocytes, monocytes link innate and acquired defense, and lymphocytes mediate pathogen-specific antibody and cell-mediated responses of the acquired immune system (Campbell 1995). The third category of index is levels of complement and natural antibodies (Matson et al. 2005), which provide a first line of defense against spreading infections via cell lysis and link innate and acquired immunity (Ochsenbein and Zinkernagel 2000).

In this study, we describe how immune function changes over the annual cycle of red knots and how ambient temperature influences this pattern. To do so, we sampled captive birds once a month over a complete annual cycle. We manipulated thermoregulatory costs, using treatments of cold, warm, and variable temperatures (the latter tracking outdoor conditions). We first establish the repeatability of immune indices, then examine how different indices covary, and finally analyze changes over the annual cycle and the effects of thermoregulatory costs.

Methods

Birds, Aviaries, and Experimental Treatments

In August and September 2004 ($n = 26$) and February 2005 ($n = 4$), red knots *Calidris canutus islandica* were captured in the Dutch Wadden Sea (53°31'N, 6°23'E) and brought into captivity at the Royal Netherlands Institute for Sea Research (NIOZ). At capture, the birds were ringed, weighed, and aged (all were older than 2 years; Prater et al. 1977), and we later determined sex using molecular techniques (Baker et al. 1999). We randomly assigned the birds (total $n = 30$, 21 females, 9 males) to three treatment groups in identical indoor aviaries (4.5 m $\times$ 1.5 m $\times$ 2.3 m high). The birds had free access to food (mud-snails *Hydrobia ulvae*), fresh water for drinking, and before the start of measurements, all birds had at least 1 month to acclimate to captivity. Photoperiod was set to track the seasonal changes in day length in the northern Netherlands for all cages. To verify that all study birds were healthy, we performed weekly assessments of body mass, feather condition, and the skin of the feet, elbows (carpals), and keel for signs of local inflammation (e.g., bumblefoot).

The variable-treatment birds ($n = 12$) were divided equally between two aviaries ventilated with outdoor air and maintained at outdoor temperature. During the experiment, variable-treatment birds experienced a mean temperature of $15.4^\circ \pm 4.8^\circ\text{C}$ (maximum of 25.5°C on June 24, 2005; minimum of 4.6°C on March 6, 2006). This treatment approximated temperatures experienced by free-living *C. c. islandica*, which spend 10 months a year in the temperate climate of western Europe (Piersma 2007). Warm-treatment birds ($n = 12$) were also divided between two aviaries and were maintained at $24.7^\circ \pm 1.8^\circ\text{C}$ (i.e., within the zone of thermoneutrality; Wiersma and Piersma 1994; Piersma et al. 1995). Cold-treatment birds ($n = 6$) were kept in a single aviary maintained at $4.9^\circ \pm 1.2^\circ\text{C}$. These temperature differences led to a 14% higher mass-specific basal metabolic rate in the cold than in the warm birds in February and March 2005 (Vézina et al. 2006). All groups were similar in terms of sex ratio and morphometrics (Vézina et al. 2006).

During the experiment, three birds died (all female). Two cold-treatment birds that died in April and August 2005 were replaced sequentially in June (with a female captured in February 2005) and September (with a male captured in August 2005). Both were given at least a month to acclimate to captivity before entering the experiment. One warm-treatment bird that died in February 2006 was not replaced because there was insufficient time to capture and acclimate a new bird before the end of the experiment.

Blood Sampling

Birds were sampled monthly, with sampling order randomized by cage and individual. We always collected blood in the morning between 10:30 and 11:00 a.m. (mean $\pm$ SD = $10:44 \pm 5$ min). Each day, we sampled two birds within 20 min of entering an aviary (mean $\pm$ SD = 6.89 ± 4.48 min). We thoroughly sterilized the area around the brachial vein with 70% ethanol and then collected about 600 μL of blood into presterilized, heparinized capillary tubes. Immediately after sampling, we made blood smears and transported the remaining blood in sterile boxes to the laboratory for further processing within an hour of sampling.

Body Mass and Molt

Throughout the experiment, all birds were weighed and scored for molt and plumage color once a week. We scored wing molt by ranking the left primary feathers on a scale of 0–5 (0 for old feathers, 5 for fully grown new feathers, 1–4 for growing feathers, depending on completeness; Ginn and Melville 1983). We scored body molt by examining the breast area and using the breast molt index (BMI: 0 for no feathers growing, 1 for <25% feathers growing, 2 for 25%–50% feathers growing, 3 for >50% feathers growing; Piersma and Jukema 1993). Because we were interested in molt as a whole, we calculated total molt (TM) as the sum of BMI and the number of growing primaries (feathers in wing molt categories 1–4).

Measuring Immune Function

Microbial Killing Abilities. We followed the basic procedure of Millet et al. (2007) and Tieleman et al. (2005) and performed the assay in a sterile working environment (CleanAir CA/R3 Flow Cabinet, CleanAir Techniek, Woerden, The Netherlands). To minimize the effects of different antigen-exposure histories, we used three ubiquitous microbial strains: *Escherichia coli* (ATCC 8739), *Candida albicans* (ATCC 10231), and *Staphylococcus aureus* (ATCC 6538; see the appendix in the online edition of the *American Naturalist* for further details). First, we reconstituted lyophilized microbial pellets, following the manufacturer's instructions (MicroBioLogics, St. Cloud, MN). Then we diluted freshly collected blood in CO_2 -independent medium (18045-054, Invitrogen, Breda, The Netherlands). After subdividing the diluted blood samples for different incubation times (*E. coli*: dilution 1 : 10, incubation for 10 or 20 min; *C. albicans*: dilution 1 : 10, incubation for 30 or 60 min; *S. aureus*: dilution 1 : 4, incubation for 60 or 120 min), we added 20 μL of each microbial suspension to each subsample. These mixtures (totaling 220 μL) were

incubated at 41°C, and 75 μL was spread evenly onto agar plates in duplicate. Inverted plates were incubated at 36°C overnight; colonies were counted the next day.

For all strains, we calculated the proportion of microorganisms killed as 1 minus the number of colonies on blood plates (described above) relative to the number of colonies on inoculate control plates (200 μL of medium mixed with 20 μL of microorganism without incubation). These inoculate plates reflect the initial situation, when the blood starts to act on the microorganism, thus best mimicking the situation of a bird responding to a pathogen.

Incubation times for *C. albicans* and *S. aureus* were long enough to allow microbial growth in some cases (i.e., negative killing; see the appendix). To take this into account, we prepared extra inoculate plates, which were incubated the same amount of time as plates containing microorganism and blood. We included counts from these plates in our statistical models as incubated controls (see “Statistics”).

Circulating Cellular Immunity. At the end of each month, blood smears were randomized and counted blind to treatment by a single observer (D.M.B.). After staining (Giemsa Stain, Sigma-Aldrich, Zwijndrecht, The Netherlands), the smears were examined at 1,000 $\times$ magnification with oil immersion, and the first 100 leukocytes were counted and classified as heterophils, eosinophils, lymphocytes, or monocytes. We were unable to distinguish B and T cells on our blood smears; thus, both are included in the lymphocyte count. Basophils were extremely rare (<0.5%) and were therefore not included in the counts. The number of thrombocytes seen while counting the first 100 leukocytes was also recorded as an estimate of the relative number of thrombocytes per leukocyte. In combination with the blood smears, we obtained leukocyte concentrations using the indirect eosinophil Unopette method (Campbell 1995), following the manufacturer’s instructions (5877; Becton Dickinson, Franklin Lakes, NJ).

Complement and Natural Antibodies. We performed the assay as described by Matson et al. (2005). We pipetted 25 μL of plasma into the first and second rows of columns B–G in 96-well plates. Chicken serum (Gibco 16110-082, Breda, The Netherlands) was used as a positive standard in columns A and H. Using Dulbecco’s phosphate-buffered solution (PBS; Mauck et al. 2005), we serially diluted the plasma and controls from row 2 to row 11 and left the twelfth row as a negative control (PBS only). We then added 25 μL of 1% of rabbit red blood cell suspension to all wells and incubated the plates at 37°C for 90 min. After incubation, plates were tilted at a 45° angle and scanned (Epson Perfection 4990 scanner) for agglutination after 20

min and lysis after 90 min. The scans were randomized with respect to sample origin, plate, and location within the plate and scored blindly for lysis and agglutination by a single observer (D.M.B.). Hemolysis, indicating complement and other lytic proteins, was identified by the absence of intact cells and the presence of free hemoglobin. Hemagglutination, indicating natural antibody activity, was identified by clumped red blood cells. Scores reflect the last plasma dilution in the dilution series (i.e., rows 2–11) exhibiting lysis or agglutination (see fig. 1 of Matson et al. 2005 for an example).

Statistics

We used one-sample Kolmogorov-Smirnov tests and visual examination of histograms to examine response variables and model residuals for normality. Although we calculated microbial killing as the proportion of colonies killed, the data were better described by normal than by binomial distributions (appendix). Leukocyte data were right skewed and $\log_{10}$ transformed. After transformation, all variables and the residuals of the models were normally distributed. Thus, we performed univariate analyses using a linear mixed model with treatment and month as fixed factors and cage and bird as random factors. This approach allowed us to incorporate the nested and repeated-measures design of our experiment (bird within cage within treatment, individual birds sampled monthly) and to include all individuals, even those with missing values (Quinn and Keough 2002). In all comparisons, we ran the models including and excluding an effect of sex. Sex was never significant and never changed the outcome of the model; thus, the statistics presented are from models excluding sex. We also included mass and time from cage entry to blood sampling as covariates in all models. For microbial killing capacities, we included the number of colonies inoculated, the ratio of incubated control and inoculation, and the microorganism suspension age (in days) as covariates. Covariates were sequentially removed from the models when not statistically significant at the 0.05 level. Effect sizes were calculated as generalized η^2 (η_G^2 ; Olejnik and Algina 2003), and repeatability was calculated following Lessells and Boag’s (1987) use of general linear models with fixed effects and covariates.

To gain insight into relationships between different measures of immune function, we performed principal-component analyses (PCAs). We included only one time point for each killing measure (*E. coli* after 10 min, *C. albicans* after 60 min, and *S. aureus* after 120 min), and we excluded total leukocyte concentration because it is the sum of the differential concentrations. Eosinophil concentrations were also excluded because of a large number of 0 values. For the leukocyte concentrations we used the transformed

data (although transformed and untransformed data produced the same result). To account for the nested and repeated-measures structure of our data, we performed the PCAs on correlation matrixes, with the data partitioned into within-individual (between-month) and among-individual levels following the procedure described by Matson et al. (2006b). We used varimax rotation to maximize contrasts of variable loadings between factors (NB, with this rotation, component scores calculated from the rotated loadings can be correlated; Jolliffe 1995). We also tested the saliency criteria for these loadings (Cliff and Hamburger 1967) and saved scores for components with eigenvalues >1 for further analysis. SPSS, version 14.0 (SPSS 2005), was used for univariate comparisons, and we created the correlation matrixes and ran the PCAs using STATISTICA 7 (StatSoft 2004).

Results

Patterns over the Annual Cycle

Body Mass and Molt. Body mass and molt data confirm previous findings that red knots in captivity retain fattening and molt schedules (Jenni-Eiermann et al. 2002). Birds fattened in early May, reached peak mass in late May, lost mass in June, and restabilized mass in July (fig. 1A; table 1). Molt data show that prealternate molt (breast, back, and head feathers) climaxed during April and May. Pre-basic molt (all body, wing, and tail feathers) peaked from July until September, but low levels of wing molt extended until January (fig. 1B; table 1).

Immune Function. Repeatability and variability of individual immune measures. Immune function varied throughout the annual cycle at both the among- and within-individual levels. Among-individual differences explained 5%–47% of total variation, depending on the index, and were significantly repeatable in all measures over the annual cycle (table 2). At the within-individual level, all immune indices

Figure 1: Trends in (A) body mass, (B) molt, and (C–J) constitutive immunity throughout the annual cycle. In cases where significant treatment $\times$ month interactions were detected, open circles, filled circles, and triangles indicate cold, variable, and warm treatments, respectively. For all other variables, the general trend is presented. For E–H, statistical analysis was performed on transformed values, but the raw data are shown for ease of interpretation. Symbols represent means, and error bars show 1 SE. Because of repeated measures, error bars can be used only to infer statistical differences between treatments in A, C, F, and H, not month-to-month differences. The annual-cycle stages for free-living (Buehler and Piersma 2008) and captive (Jenni-Eiermann et al. 2002) birds are given above the plots. Numbers under the months correspond to the labels in figure 2C, and boxed points in month 13 represent variable-treatment birds only.

Table 1: Categorization of months

Month ^a	Mean body mass $\pm$ SE (g)	Mass description	Mean total molt $\pm$ SE ^b	Total molt description	Category ^c
March	123.1 $\pm$ 1.8	Stable	.10 $\pm$ .17	Latent	Overwintering (B)
April	124.4 $\pm$ 1.3	Mass gain	1.20 $\pm$ .18	Body molt	Migration (E)
May	144.9 $\pm$ 1.3	Peak mass	.61 $\pm$ .12	Latent	Migration (F)
June	121.2 $\pm$ 1.0	Mass loss	.35 $\pm$ .10	Latent	Migration (G)
July	119.8 $\pm$ 1.2	Lowest and rebound	2.74 $\pm$ .11	Wing and molt increasing	Oversummering (H)
August	127.4 $\pm$ 1.1	Stable	4.01 $\pm$ .10	Peak wing and body molt	Prebasic molt (A)
September	129.6 $\pm$ 1.2	Stable	3.31 $\pm$ .12	Wing and body molt decreasing	Prebasic molt (A)
October	127.6 $\pm$ 1.2	Stable	1.97 $\pm$ .11	Wing molt decreasing	Overwintering (B)
November	129.7 $\pm$ 1.1	Stable	1.59 $\pm$ .10	Wing molt decreasing	Overwintering (B)
December	132.7 $\pm$ 1.4	Stable	1.22 $\pm$ .13	Wing molt decreasing	Overwintering (B)
January	133.3 $\pm$ 1.2	Stable	.85 $\pm$ .11	Latent	Overwintering (B)
February	132.9 $\pm$ 1.0	Stable	.38 $\pm$ .10	Latent	Overwintering (B)

^a Months correspond to those in figure 1.^b For calculation, see "Body Mass and Molt."^c Categories and letters are from table 1 of Jenni-Eiermann et al. (2002).

except hemagglutination varied significantly over months (table 3).

Covariation of immune variables among and within individuals. Among individuals, PCA revealed three components with eigenvalues >1 accounting for 70% of the total variability in the data (table 4, pt. A). *Staphylococcus aureus* killing, *Candida albicans* killing, heterophil concentrations, and hemolysis were positively correlated with PC1; *Escherichia coli* killing, hemagglutination, and monocyte concentrations were positively correlated with PC2

(fig. 2A; monocytes also salient on PC3); and lymphocytes and thrombocytes were positively correlated with PC3. The PC1 and PC2 scores were significantly correlated (Pearson $r = 0.763$, $P < .001$; after the outlier was omitted, $r = 0.655$, $P < .001$; fig. 2B), as were PC2 and PC3 scores ($r = 0.273$, $P = .035$). The PC1 and PC3 scores were not correlated ($r = 0.173$, $P = .186$).

Within individuals (among months), PCA also revealed three components with eigenvalues >1 accounting for 78% of the total variability in the data (table 4, pt. B). Lym-

Table 2: Individual repeatability of immune function in red knots over the annual cycle

Response	<i>r</i>	SE	df	<i>F</i>	<i>P</i>
Microbial killing (proportion killed):					
<i>Staphylococcus aureus</i> after 60 min	.15	.043	22, 235	3.26	<.001
<i>S. aureus</i> after 120 min	.216	.053	22, 235	4.72	<.001
<i>Candida albicans</i> after 30 min	.101	.033	22, 255	2.43	<.001
<i>C. albicans</i> after 60 min	.097	.032	22, 258	2.68	<.001
<i>Escherichia coli</i> after 10 min	.400	.070	22, 258	9.10	<.001
<i>E. coli</i> after 20 min	.469	.072	22, 258	11.64	<.001
White blood cell parameters (per μ L):					
Total leukocytes	.168	.046	22, 264	3.44	<.001
Heterophils	.362	.068	22, 286	7.69	<.001
Eosinophils	.061	.024	22, 286	1.79	.020
Lymphocytes	.196	.05	22, 264	3.94	<.001
Monocytes	.135	.040	22, 286	2.85	<.001
Thrombocytes	.049	.021	22, 286	1.61	.043
Plasma parameters ($-\log_2$):					
Hemolysis	.291	.063	22, 286	5.92	<.001
Hemagglutination	.107	.034	22, 286	2.441	<.001

Note: After Lessells and Boag (1987). Repeatability was calculated by using models with fixed effects and covariates from individuals for which we had data for all 12 months ($n = 27$; 11 months for microbial killing). Because covariates differed for different immune indices, error degrees of freedom differ.

Table 3: Immune function in red knots over the annual cycle

Response	Among individuals										Within individuals			
	Treatment, fixed					Cage within treatment, random					Bird within cage within treatment, random			
	df	F	P	η^2_G		df	F	P	η^2_G		df	F	P	η^2_G
Microbial killing (proportion killed):														
<i>Staphylococcus aureus</i> after 60 min	2, 2	.71	.58	<.01		2, 25	.44	.65	<.01		26, 278	3.31	<.01**	.18
<i>S. aureus</i> after 120 min	2, 2	.89	.53	.01		2, 26	.49	.62	.01		26, 258	4.14	<.01**	.18
<i>Candida albicans</i> after 30 min	2, 2	6.50	.14	.02		2, 25	.24	.78	<.01		26, 278	2.82	<.01**	.19
<i>C. albicans</i> after 60 min	2, 2	2.28	.29	<.01		2, 25	.13	.88	<.01		26, 278	2.66	<.01**	.18
<i>Escherichia coli</i> after 10 min	2, 2	.49	.72	<.01		2, 26	.13	.88	<.01		26, 279	9.31	<.01**	.13
<i>E. coli</i> after 20 min	2, 2	.32	.76	<.01		2, 26	.33	.72	<.01		26, 281	11.80	<.01**	.22
White blood cell parameters (per μL):														
Total leukocytes	2, 2	.19	.84	<.01		2, 25	2.07	.15	.03		26, 285	3.03	<.01**	.17
Heterophils	2, 2	.73	.93	<.01		2, 26	2.94	.7	.07		26, 307	6.55	<.01**	.28
Eosinophils	2, 2	.82	.55	.01		2, 25	1.15	.33	.01		26, 219	1.60	.04*	.12
Lymphocytes	2, 2	.40	.72	<.01		2, 25	1.07	.36	.02		26, 285	3.50	<.01**	.19
Monocytes	2, 2	.08	.92	<.01		2, 25	4.79	.02*	.06		26, 305	2.35	<.01**	.15
Thrombocytes	2, 2	.70	.59	.02		2, 24	4.87	.02*	.04		26, 307	1.47	.07	.09
Plasma parameters ($-\log_2$):														
Hemolysis	2, 2	7.50	.14	.07		2, 25	.35	.71	.01		26, 310	5.57	<.01**	.2
Hemagglutination	2, 2	4.40	.13	.01		2, 25	.08	.92	.01		26, 311	2.05	.01**	.17

Note: Effect sizes are generalized η^2 (η^2_G ; Olejnik and Algina 2003). Degrees of freedom vary among indices because in some birds we did not obtain the full set of immune indices from every bird in every month. Statistics for interactions are in the "Immune Function" subsection of "Effects of Temperature Treatments."

* $P < .05$.

** $P < .01$.

Table 4: Principal-component loadings after varimax rotation among and within individuals

	PC1	PC2	PC3
A. Among individuals:			
Response:			
<i>Staphylococcus aureus</i> killing (proportion killed)	.870	.067	-.178
<i>Candida albicans</i> killing (proportion killed)	.907	.152	.102
<i>Escherichia coli</i> killing (proportion killed)	.149	.872	-.104
Heterophils (per μ L)	.835	.242	.001
Lymphocytes (per μ L)	.178	-.311	.808
Monocytes (per μ L)	.186	.610	.523
Thrombocytes (per μ L)	-.243	.239	.734
Hemolysis ($-\log_2$)	.542	.337	.099
Hemagglutination ($-\log_2$)	.415	.654	.022
Totals:			
Variance per component (%)	32.1	21.3	17.0
Cumulative variance (%)	32.1	53.4	70.4
B. Within individuals:			
Response:			
<i>S. aureus</i> killing (proportion killed)	-.429	.780	.113
<i>C. albicans</i> killing (proportion killed)	-.351	.810	-.018
<i>E. coli</i> killing (proportion killed)	.290	.178	.776
Heterophils (per μ L)	.040	.730	.458
Lymphocytes (per μ L)	.836	-.401	.192
Monocytes (per μ L)	.852	.019	.252
Thrombocytes (per μ L)	.788	-.088	-.051
Hemolysis ($-\log_2$)	.496	.714	-.133
Hemagglutination ($-\log_2$)	.044	.033	-.952
Totals:			
Variance per component (%)	29.8	27.9	20.6
Cumulative variance (%)	29.8	57.7	78.3

Note: Boldface indicates the highest loading for a measure across the principal components, and underlined loadings meet the saliency criteria for that principal component.

phocytes, monocytes, and thrombocytes were positively correlated with PC1; *S. aureus* killing, *C. albicans* killing, heterophil concentrations, and hemolysis were positively correlated with PC2; and *E. coli* killing was positively correlated and hemagglutination was negatively correlated with PC3.

Changes in immune function over the annual cycle. Changes in immune indices corresponded to physiological changes that knots experience over a captive annual cycle (fig. 1). *Staphylococcus aureus* killing, *C. albicans* killing, and heterophil concentrations peaked during mass change and dropped during peak molt (fig. 1C–1E). Conversely, lymphocyte and monocyte concentrations showed a small increase during mass change but peaked during peak molt (fig. 1F, 1G). Total leukocyte concentrations (the sum of all other leukocyte types) closely matched the pattern in lymphocytes because lymphocytes made up, on average, 74% of the total leukocyte count (fig. 1H).

Escherichia coli killing and hemolysis also changed over the annual cycle; however, these changes did not occur during mass change or molt (fig. 1I, 1J). *Escherichia coli*

killing dropped between September and October and then rebounded modestly by April 2006 (fig. 1I, boxed point). In contrast, hemolysis increased from December through February and then trended back toward original levels by April 2006 (fig. 1J, boxed point).

We plotted monthly mean scores from PC1 and PC2 to further examine annual patterns among immune measures (fig. 2C). These scores cluster into three physiologically relevant periods: (1) mass change (gain, loss, and reestablishment associated with migration), (2) peak wing and prebasic molt, and (3) winter (see Jenni-Eiermann et al. 2002 and table 1 for categorization). During mass change PC2 scores are high, whereas during peak molt PC2 scores are low and PC1 scores are high. In month 5, an over-summering transition month between mass change and peak molt for captive red knots, PC2 scores are intermediate.

Effects of Temperature Treatments

Body Mass and Molt. Body mass was significantly affected by temperature treatment. Cold birds were consistently

Figure 2: Relationships among immune indices, based on principal-component analysis among individuals. Axis labels correspond to the alphabetical groups in table 5. In A, vectors show the PC1 and PC2 loadings of each measure (dashed vectors are best explained by PC3). Vector length indicates the strength of the relationship, and the angle between two vectors gives the degree of correlation (adjacent = highly correlated, orthogonal (90°) = uncorrelated, and opposite (180°) = negatively correlated). In B, the relationship between PC1 and PC2 scores among individuals is shown, and data points (mean $\pm$ SE) represent

heavier than warm birds; variable birds were intermediate (outside the fattening period; Tukey post hoc: cold > variable > warm; $P < .001$). A significant week $\times$ treatment interaction ($F = 3.26$, $df = 104, 1,379$, $P < .001$) resulted from variable birds gaining more premigratory mass than cold birds or warm birds (fig. 1A; see Vézina et al. 2007). Molt was not affected by treatment (molt: $\chi^2 = 2.33$, $df = 2$, $P = .31$).

Immune Function. Annual-cycle effects (reflected by month in table 3) dominated over the effect of our experimental manipulations of thermoregulatory costs (treatment in table 3). Significant month $\times$ treatment interactions were found only for *S. aureus* killing after 120 min ($F = 1.83$, $df = 20, 258$, $P = .02$, $\eta_G^2 = 0.09$), lymphocyte concentrations ($F = 1.86$, $df = 20, 285$, $P = .01$, $\eta_G^2 = 0.10$), and total leukocyte concentrations ($F = 1.98$, $df = 20, 285$, $P < .01$, $\eta_G^2 = 0.13$). With *S. aureus* killing, the interaction, which was driven by significantly lower killing ability in the variable treatment in April ($F = 30.97$, $df = 2, 3$, $P = .008$), is clearly less important than the overall temporal trends (fig. 1C). For lymphocyte and total leukocyte concentrations, cold birds peaked in June and July during lowest mass; warm and variable birds peaked in August and September during peak molt (fig. 1F, 1H); however, Tukey post hoc tests revealed no significant treatment effect in any month. No other immune measures showed significant treatment effects (table 3) or significant treatment $\times$ month interactions (all $P > .10$).

Discussion

This study explored immune function over the annual cycle of red knots and examined how ambient temperature influenced this pattern. Our results established the repeatability of immune indices at the level of individual birds and showed that immune indices covaried similarly among and within birds over time, suggesting functional immune strategies. We discuss these strategies in terms of their possible costs and protective benefits at the among- and within-individual levels. We then discuss how these strategies shift during different stages of the annual cycle in terms of physiological trade-offs and predicted disease risk in the wild. Finally, we discuss the relative lack of temperature effect on these patterns.

individual birds. In C, relationships between PC1 and PC2 scores within individuals indicate redistribution of immune strategies over the annual cycle. Points represent months (mean $\pm$ SE), and numbers correspond to the months in figure 1. Annual-cycle stages are based on Jenni-Eiermann et al. (2002) and table 1.

Relationships among Immune Indices

We examined relationships among nine immune indices with a range of protective benefits and hypothesized energetic and immunopathology costs (table 5). The PCAs identified three axes (PCs) consisting of the same indices; however, the ordering of the PCs differed within and among individuals. For clarity, we define these axes as groups A–C (table 5) and refer to them as such throughout the discussion.

Group A comprised *Staphylococcus aureus* killing, *Candida albicans* killing, heterophils, and hemolysis (table 4). *Staphylococcus aureus* and *C. albicans* killing are primarily the result of phagocytosis (Millet et al. 2007), and heterophils are important phagocytic cells in birds and are the avian equivalent of the mammalian neutrophils (Campbell 1995). Neutrophil phagocytosis produces harmful reactive oxygen and nitrogen species (Spletstoesser and Schuff-Werner 2002), and energy requirements may be heightened by their high turnover rates (Janeway et al. 2004). Complement, which mediates lysis, is linked to inflammation (high immunopathology costs) and must be tightly regulated to prevent host-tissue damage (Janeway et al. 2004). Thus, we associate group A with more costly immunity in terms of collateral damage to the host (immunopathology cost) and energy requirements when a host encounters a pathogen and the constitutive immunity we measured is put to use (table 5; Klasing 2004; Buehler et al. 2008). Functionally, these indices are associated with protection from extracellular bacteria and yeast infections. Thus, this strategy (group A) may be favored under conditions where diversity of these pathogens is high and encounters with new strains are frequent (Schmid-Hempel and Ebert 2003).

Group B comprised *E. coli* killing and natural antibodies (table 4). Killing of the *E. coli* strain we used relies on soluble components in blood plasma (Matson et al. 2006a; Millet et al. 2007) rather than more costly phagocytosis (Spletstoesser and Schuff-Werner 2002). Natural antibodies link innate and acquired immune systems (Ochsenbein and Zinkernagel 2000). These molecules exhibit weak affinities and low specificities but are important for initiating acquired antibody responses, which are highly specific and unlikely to be self-reactive. Thus, relative to group A, we interpreted group B as moderate in terms of immune cost (table 5). Functionally, *E. coli* killing represents the ability to control or prevent infection by some gram-negative bacteria, and natural antibodies are a constitutive component of the acquired immune system, providing a first line of defense against pathogen attack (Ochsenbein et al. 1999). Interestingly, hemolysis did not correlate with group B. Complement and other lytic proteins are also soluble and have correlated with natural

antibodies in other analyses (Matson et al. 2006b; Buehler et al. 2008). The inclusion of monocytes in group B at the among-individual level is also puzzling; however, because monocytes meet the saliency criteria for group C among individuals and clearly fall into group C within individuals (table 4, pt. B), we discuss them with group C.

Group C comprised lymphocytes, monocytes, and thrombocytes (table 4). Lymphocytes form the basis of cell-mediated (T-cell) and antibody-mediated (B-cell) acquired immunity (Janeway et al. 2004). Antibody-mediated immunity, in particular, is considered low cost because of its high efficiency and high specificity, which lower energy and immunopathology costs (Klasing 2004). Monocytes mature into long-lived macrophages and antigen-presenting cells important for initiating acquired immune responses, and thrombocytes are important for blood coagulation (Janeway et al. 2004). Thus, relative to groups A and B, we interpreted group C as lower in cost because of its association with acquired immunity (table 5). Functionally, lymphocyte- and monocyte-based immunity might be favored in conditions where energy demands are high or where disease risk is associated with reoccurring infections rather than with novel pathogens (Schmid-Hempel and Ebert 2003).

Immune Function as an Individual Trait

Our immune indices covaried similarly among and within birds over time, suggesting functional immune strategies at multiple levels of organization. This idea is supported by a study describing species-specific immune strategies in mammals (Martin et al. 2007) and a study describing similar covariation among constitutive immune indices in captive and free-living red knots (Buehler et al. 2008). We found that the proportion of variability explained by each strategy differed among and within individuals (table 4). Higher-cost indices (table 5) explained more variability among birds, suggesting that individuals are best distinguished by more costly immunity. Furthermore, the positive correlation between group A and B indices among birds (fig. 2B) indicates that individuals maintain either high or low levels of more costly immune strategies. It is also notable that *E. coli* killing and natural antibody levels are positively correlated among individuals (table 4, pt. A) but negatively correlated within individuals (table 4, pt. B). This indicates that, as in the relationship between group A and B strategies, individuals maintain either high or low levels of soluble constitutive defenses but that across the year there is an antagonistic relationship between *E. coli* killing and natural antibody levels.

Table 5: Summary and clarification of the nine independent indices measured and hypothesized costs and classifications for the groupings identified by principal-component analysis

Specific index	Description and function	Strategy	Hypothesized cost	Among bird	Within bird
Group A		Constitutive innate immunity (phagocytosis and inflammation)	High: high energy costs due to short-lived cells and high immunopathology costs due to inflammation	PC1	PC2
<i>Staphylococcus aureus</i> killing	Bacteria defense, mostly phagocytosis ^a				
<i>Candida albicans</i> killing	Yeast defense, mostly phagocytosis ^a				
Heterophils	Short-lived phagocytes associated with inflammation ^{b,c}				
Hemolysis	Complement mediated lysis during inflammation ^d				
Group B		Constitutive innate immunity (soluble factors)	Moderate: lower energy and immunopathology costs for killing by soluble factors rather than phagocytosis	PC2	PC3
<i>Escherichia coli</i> killing	Bacteria defense, little phagocytosis ^a				
Hemagglutination	Natural antibody mediated clumping of invaders ^d				
Group C		Constitutive potential for induced acquired immunity	Low: low energy costs due to longer-lived cells and low immunopathology costs due to less host damage	PC3	PC1
Lymphocytes	Basis of specific induced immunity ^b				
Monocytes	Long-lived phagocytes and antigen presenting cells for specific immunity ^b				
Thrombocytes	Important for blood clotting ^b				

^a Millet et al. 2007.

^b Campbell 1995; Janeway et al. 2004.

^c Spletstoesser and Schuff-Werner 2002.

^d Matson et al. 2005.

*Redistribution of Immune Function in
Different Annual-Cycle Stages*

Over the annual cycle, limited resources must be allocated to immune defense, predicting an optimal portfolio of defenses (Schmid-Hempel and Ebert 2003). Our results suggest that different strategies may be used during different periods of the year (fig. 2C).

The period of mass change in captive birds is associated with fattening and spring migration in the wild. Because fattening and migration are energy-demanding activities, trade-offs with immune function may be predicted when resources are limited. However, our results indicate that group A indices, which are associated with higher energy and immunopathology costs, were high during the period of mass change. This may be due to the fact that our birds had ad lib. access to food and were not experiencing resource limitation. Studies manipulating food availability are under way to examine this possibility (Buehler et al., forthcoming), and studies on free-living birds will be necessary to test whether trade-offs occur under resource-limited conditions in the wild. From a disease perspective, fattening and spring migration are considered high risk (Buehler and Piersma 2008). During migration, birds pass through a variety of environments where they might encounter novel pathogens (Møller and Erritzøe 1998), and during stopovers they feed in dense flocks, facilitating disease transmission (Altizer et al. 2006). Thus, during this period, the protective benefits of group A indices might outweigh their costs.

The period of body mass stabilization in captive birds is associated with oversummering or breeding in the wild. During this period, we found an abrupt drop in group A indices (month 5 in figs. 1C–1E, 2B), which may represent a transition period between mass change and peak molt in captive birds. From a disease perspective, in the wild, once the knots establish breeding territories, they are widely dispersed. Thus, transmission risk and the change of encountering novel pathogens may be decreased (Buehler and Piersma 2008).

Peak molt occurs at approximately the same time in captive and free-living knots (fig. 1). During molt, feathers must be synthesized by keratinocytes (Haake and Sawyer 1986), and experiments have demonstrated that hydrogen peroxides, such as those generated during respiratory burst and phagocyte ingestion (Splettstoesser and Schuff-Werner 2002), inhibit keratinocyte proliferation (O'Toole et al. 1996). Furthermore, growing feathers are a potential cause of dermal inflammation (Silverin et al. 1999), which can be costly in terms of immunopathology (Råberg et al. 1998). Thus, immune strategies that counter infection with less phagocytosis should be favored during molt. Our results support this idea, because we found a clear shift from

group A to group C indices during peak molt (fig. 2C), indicating a decrease in phagocytosis-based defenses and a greater role for lymphocytes and monocytes.

For the annual cycle as a whole, our data suggest two main strategies. Group A indices represent defenses that are necessary during periods of high disease risk but are costly enough to warrant downregulation when disease risks are lower and the costs of these defenses outweigh their benefits. Conversely, group C indices represent relatively low cost immunological functions that may be necessary year-round. To fully test the ideas presented here, studies on free-living birds and induced immune responses will also be required.

*No Trade-Off between Constitutive Immunity
and Thermoregulatory Costs*

Contrary to predictions that ambient temperature might affect immune function in knots, this study suggests that temperature in itself is relatively unimportant to overall annual variation. Higher thermoregulatory costs alone do not seem to necessitate trade-offs between constitutive immune function and thermoregulation. Nevertheless, cold birds did respond physiologically to low temperatures. They were heavier than warm birds (fig. 1) and exhibited 14% higher mass-specific basal metabolic rate than warm birds in February and March 2005 (Vézina et al. 2006). Exposure to cold for a year also led to larger spleen size in cold birds (100% heavier than that of warm birds, fresh mass; F. Vézina, unpublished data). The spleen is an important organ for the presentation and recognition of antigens in induced acquired immunity (Janeway et al. 2004; but see Smith and Hunt 2004), indicating that thermoregulatory costs may affect aspects of immunity that were not measured.

Another explanation for the limited effects of temperature treatments on measured immune indices may be that the ad lib. feeding regime allowed cold-treatment birds to compensate for increased energy expenditure via higher food intake (Vézina et al. 2006). Energy expenditure is only one component of the energy budget, and future experiments that simultaneously manipulate energy intake are needed to examine the effect of limited resources on immune function. Furthermore, future experiments manipulating other environmental factors that vary over the annual cycle (e.g., photoperiod and disease risk) will help resolve the proximate causes of variation in immune function in wild birds.

Acknowledgments

We thank M. Brugge, A. Gustowska, K. Jalvingh, M. Petit, and F. Vézina for help with captive knots; C. Pool of the

Animal Experiment Committee of the Royal Netherlands Academy for Arts and Sciences for overseeing experimental procedures (protocol NIOZ.05.01); and M. Brugge, A. Dekinga, B. Spaans, and the crew of the research vessel *Navicula* for catching birds and fishing *Hydrobia*. We thank D. Barjaktarov, A. Castillo, S. Haan, M. Lange, and E. Marks for lab assistance and J. van Bleijswijk for lab space. We thank D. Visser for excellent help with finalizing the figures. This article was greatly improved by comments from A. Hegemann, B. Helm, F. Vézina, and several anonymous reviewers. Funding was provided by the Natural Science and Engineering Council of Canada (NSERC PGSB-267701-2003), the University of Groningen, and Schure-Beijerinck-Popping Fonds to D.M.B., the Netherlands Organization for Scientific Research and the University of Groningen to T.P. and B.I.T., and the Royal Netherlands Institute of Sea Research to T.P.

Literature Cited

- Altizer, S., A. Dobson, P. Hosseini, P. Hudson, M. Pascual, and P. Rohani. 2006. Seasonality and the dynamics of infectious diseases. *Ecology Letters* 9:467–484.
- Baker, A. J., T. Piersma, and A. D. Greenslade. 1999. Molecular vs. phenotypic sexing in red knots. *Condor* 101:887–893.
- Battley, P. F., T. Piersma, D. I. Rogers, A. Dekinga, B. Spaans, and J. A. van Gils. 2004. Do body condition and plumage during fuelling predict northwards departure dates of great knots *Calidris tenuirostris* from north-west Australia? *Ibis* 146:46–60.
- Buehler, D. M., and T. Piersma. 2008. Travelling on a budget: predictions and ecological evidence for bottlenecks in the annual cycle of long-distance migrants. *Philosophical Transactions of the Royal Society B: Biological Sciences* 363:247–266.
- Buehler, D. M., T. Piersma, and B. I. Tieleman. 2008. Captive and free-living red knots *Calidris canutus* exhibit differences in non-induced immunity that suggest different immune strategies in different environments. *Journal of Avian Biology* 39:560–566.
- Buehler, D. M., F. Encinas-Viso, M. Petit, F. Vézina, B. I. Tieleman, and T. Piersma. Forthcoming. Limited access to food and physiological trade-offs in a long-distance migrant shorebird. II. Constitutive immune function and acute phase response. *Physiological and Biochemical Zoology*.
- Campbell, T. W. 1995. *Avian hematology and cytology*. Iowa State University Press, Ames.
- Cliff, N., and C. D. Hamburger. 1967. Study of sampling errors in factor analysis by means of artificial experiments. *Psychological Bulletin* 68:430.
- Ginn, H. B., and D. S. Melville. 1983. Molt in birds. *BTO Guide* 19. British Trust for Ornithology, Tring.
- Haake, A. R., and R. H. Sawyer. 1986. Differences in the histogenesis and keratin expression of avian extraembryonic ectoderm and endoderm recombined with dermis. *Developmental Biology* 113:295–304.
- Hasselquist, D. 2007. Comparative immunoecology in birds: hypotheses and tests. *Journal of Ornithology* 148(suppl.):S571–S582.
- Janeway, C. A., P. Travers, M. Walport, and M. Shlomchik. 2004. *Immunobiology: the immune system in health and disease*. Garland, New York.
- Jenni-Eiermann, S., L. Jenni, and T. Piersma. 2002. Plasma metabolites reflect seasonally changing metabolic processes in a long-distance migrant shorebird (*Calidris canutus*). *Zoology* 105:239–246.
- Jolliffe, I. T. 1995. Rotation of principal components: choice of normalization constraints. *Journal of Applied Statistics* 22:29–35.
- King, J. R. 1974. Seasonal allocation of time and energy resources in birds. Pages 4–85 in R. A. J. Paynter, ed. *Avian energetics*. Nuttall Ornithological Club, Cambridge, MA.
- Klasing, K. C. 2004. The costs of immunity. *Acta Zoologica Sinica* 50:961–969.
- Lee, K. A. 2006. Linking immune defenses and life history at the levels of the individual and the species. *Integrative and Comparative Biology* 46:1000–1015.
- Lessells, C. M., and P. T. Boag. 1987. Unrepeatable repeatabilities: a common mistake. *Auk* 104:116–121.
- Lochmiller, R. L., and C. Deerenberg. 2000. Trade-offs in evolutionary immunology: just what is the cost of immunity? *Oikos* 88:87–98.
- Martin, L. B., Z. M. Weil, and R. J. Nelson. 2007. Immune defense and reproductive pace of life in *Peromyscus* mice. *Ecology* 88:2516–2528.
- . 2008. Seasonal changes in vertebrate immune activity: mediation by physiological trade-offs. *Philosophical Transactions of the Royal Society B: Biological Sciences* 363:321–339.
- Matson, K. D., R. E. Ricklefs, and K. C. Klasing. 2005. A hemolysis-hemagglutination assay for characterizing constitutive innate humoral immunity in wild and domestic birds. *Developmental and Comparative Immunology* 29:275–286.
- Matson, K. D., B. I. Tieleman, and K. C. Klasing. 2006a. Capture stress and the bactericidal competence of blood and plasma in five species of tropical birds. *Physiological and Biochemical Zoology* 79:556–564.
- Matson, K. D., A. A. Cohen, K. C. Klasing, R. E. Ricklefs, and A. Scheuerlein. 2006b. No simple answers for ecological immunology: relationships among immune indices at the individual level break down at the species level in waterfowl. *Proceedings of the Royal Society B: Biological Sciences* 273:815–822.
- Mauck, R. A., K. D. Matson, J. Philipsborn, and R. E. Ricklefs. 2005. Increase in the constitutive innate humoral immune system in Leach's storm-petrel (*Oceanodroma leucorhoa*) chicks is negatively correlated with growth rate. *Functional Ecology* 19:1001–1007.
- Millet, S., J. Bennett, K. A. Lee, M. Hau, and K. C. Klasing. 2007. Quantifying and comparing constitutive immunity across avian species. *Developmental and Comparative Immunology* 31:188–201.
- Møller, A. P., and J. Erritzøe. 1998. Host immune function and sexual selection in birds. *Journal of Evolutionary Biology* 11:703–719.
- Nelson, R. J., G. E. Demas, S. L. Klein, and L. J. Kriegsfeld. 2002. Seasonal patterns of stress, immune function and disease. Cambridge University Press, Cambridge.
- Norris, K., and M. R. Evans. 2000. Ecological immunology: life history trade-offs and immune defense in birds. *Behavioral Ecology* 11:19–26.
- Ochsenbein, A. F., and R. M. Zinkernagel. 2000. Natural antibodies and complement link innate and acquired immunity. *Immunology Today* 21:624–630.
- Ochsenbein, A. F., T. Fehr, C. Lutz, M. Suter, F. Brombacher, H. Hengartner, and R. M. Zinkernagel. 1999. Control of early viral and bacterial distribution and disease by natural antibodies. *Science* 286:2156–2159.

- Olejnik, S., and J. Algina. 2003. Generalized eta and omega squared statistics: measures of effect size for some common research designs. *Psychological Methods* 8:434–447.
- O'Toole, E. A., M. Goel, and D. T. Woodley. 1996. Hydrogen peroxide inhibits human keratinocyte migration. *Dermatologic Surgery* 22: 525–529.
- Piersma, T. 2007. Using the power of comparison to explain habitat use and migration strategies of shorebirds worldwide. *Journal of Ornithology* 148(suppl.):S45–S59.
- Piersma, T., and J. Jukema. 1993. Red breasts as honest signals of migratory quality in a long-distance migrant, the bar-tailed godwit. *Condor* 95:163–177.
- Piersma, T., N. Cadée, and S. Daan. 1995. Seasonality in basal metabolic rate and thermal conductance in a long-distance migrant shorebird, the knot (*Calidris canutus*). *Journal of Comparative Physiology B* 165:37–45.
- Prater, A. J., J. H. Marchant, and J. Vuorinen. 1977. Guide to the identification and aging of Holarctic waders. BTO Guide 17. British Trust for Ornithology, Tring.
- Quinn, G. P., and M. J. Keough. 2002. Experimental design and data analysis for biologists. Cambridge University Press, Cambridge.
- Råberg, L., M. Grahm, D. Hasselquist, and E. Svensson. 1998. On the adaptive significance of stress-induced immunosuppression. *Proceedings of the Royal Society B: Biological Sciences* 265:1637–1641.
- Schmid-Hempel, P., and D. Ebert. 2003. On the evolutionary ecology of specific immune defence. *Trends in Ecology & Evolution* 18: 27–32.
- Sheldon, B. C., and S. Verhulst. 1996. Ecological immunology: costly parasite defences and trade-offs in evolutionary ecology. *Trends in Ecology & Evolution* 11:317–321.
- Silverin, B., R. Fänge, P.-A. Viebke, and J. Westin. 1999. Seasonal changes in mass and histology of the spleen in willow tits *Parus montanus*. *Journal of Avian Biology* 30:255–262.
- Smith, K. G., and J. L. Hunt. 2004. On the use of spleen mass as a measure of avian immune system strength. *Oecologia (Berlin)* 138: 28–31.
- Spletstoeser, W. D., and P. Schuff-Werner. 2002. Oxidative stress in phagocytes: “the enemy within.” *Microscopy Research and Technique* 57:441–455.
- SPSS. 2005. SPSS. Version 14.0. SPSS, Chicago.
- StatSoft. 2004. STATISTICA. Version 7. StatSoft, Tulsa, OK.
- Tieleman, B. I., J. B. Williams, R. E. Ricklefs, and K. C. Klasing. 2005. Constitutive innate immunity is a component of the pace-of-life syndrome in tropical birds. *Proceedings of the Royal Society B: Biological Sciences* 272:1715–1720.
- Vézina, F., K. M. Jalvingh, A. Dekinga, and T. Piersma. 2006. Acclimation to different thermal conditions in a northerly wintering shorebird is driven by body mass-related changes in organ size. *Journal of Experimental Biology* 209:3141–3154.
- . 2007. Thermogenic side effects to migratory predisposition in shorebirds. *American Journal of Physiology* 292:R1287–R1297.
- Wiersma, P., and T. Piersma. 1994. Effects of microhabitat, flocking, climate and migratory goal on energy expenditure in the annual cycle of knots. *Condor* 96:257–279.

Associate Editor: Greg Demas
Editor: Monica A. Geber

Red knot (*Calidris canutus*) feeding at Delaware Bay (photograph by Deborah Buehler).