University of Groningen

Conformational Disorder and Ultrafast Exciton Relaxation in PPV-family Conjugated Polymers

Dykstra, Tienieke E.; Hennebicq, Emmanuelle; Beljonne, David; Gierschner, Johannes; Claudio, Gil; Bittner, Eric R.; Knoester, Jasper; Scholes, Gregory D.

Published in:
Journal of Physical Chemistry B

DOI:
[10.1021/jp807249b](https://doi.org/10.1021/jp807249b)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2009

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Dykstra, T. E., Hennebicq, E., Beljonne, D., Gierschner, J., Claudio, G., Bittner, E. R., Knoester, J., & Scholes, G. D. (2009). Conformational Disorder and Ultrafast Exciton Relaxation in PPV-family Conjugated Polymers. *Journal of Physical Chemistry B*, 113(3), 656-667. <https://doi.org/10.1021/jp807249b>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Conformational Disorder and Ultrafast Exciton Relaxation Dynamics in PPV-family Conjugated Polymers

Tieneke E. Dykstra¹, Emmanuelle Hennebicq², David Beljonne², Johannes Gierschner², Gil Claudio³, Eric R. Bittner⁴, Jasper Knoester⁵ and Gregory D. Scholes^{1}*

1 Lash-Miller Chemical Laboratories, Institute for Optical Sciences and Centre for Quantum Information and Quantum Control, University of Toronto, 80 St. George Street, Toronto, Ontario, M5S 3H6 Canada

2 Laboratory for Chemistry of Novel Materials, Center for Research on Molecular Electronics and Photonics, University of Mons-Hainaut, Place du Parc 20, B-7000 Mons Belgium

3 Max Planck Institute for Polymer Research Postfach 3148 D-55021 Mainz, Germany

4 Department of Chemistry and Center for Materials Chemistry, University of Houston, Houston, Texas 77204

5 Institute for Theoretical Physics and Zernike Institute for Advanced Materials, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands

Additional Supporting Information

The three archives contain information on the structure and calculations for the three PPV polymer chains described in the paper. There are 4 files for each of the three chains.

Detailed description of the data:

coords.dat

Column one : Atom number

Column two: Atom type (C or H).

Column three: Chromophore number (which atoms belong to oligomer 1, 2, and so forth). All the atoms for every chromophore are listed contiguously, that is, once the number changes from n to $n+1$ (from 1 to 2, for example), there will be no other atom in the list that will belong to chromophore n .

Column four: Monomer number. One phenyl group is considered one monomer and one vinyl group is also considered as one monomer

Columns five, six, seven: xyz-coordinates of the atom.

Structure of other data files:

Chromophores in PPV chain with size smaller or equal to 2 units (one vinylene or phenylene group being considered as one unit)-- isolated phenylene, isolated vinylene , phenylenevinylene group -- were not included in the calculations. The excitations energies, transition dipole moments, and coupling factors that are given in the different data files only concern the so-called “effective” chromophores with size larger than 2 (phenylene or vinylene) units.

res.dat files:

Line 1: total number of chromophores; number of effective chromophores (Neff) with size larger than two units, the number of excited states computed for all chromophores=30

Following lines:

Column two: Index of the chromophores,

Column three: number of atoms for this chromophore

Column four: the number excited states with energy lower than 4 eV.

The rest of the files deal with only the Neff effective chromophores.

coup.dat

Columns one and two: labels of chromophores of interest

Columns three and four: the index of the excited states considered for the two chromophores

Column five: the electronic coupling in cm⁻¹.

Note: coupling is given between the first excited state of chromophore in column 1 with each of the n=30 excited states of chromophore in column 2. There are Neff blocks of 30 lines in this file, Neff being the number of effective chromophores

ene.dat

Columns one and two: For each effective chromophore the excitation energy of the first 30 singlet excited states (in eV and cm⁻¹)

Columns three, four, five: xyz components of the corresponding transition dipole moments (in Debye).

There are thus Neff blocks of 30 lines in this file.