University of Groningen

Calcanectomy, an alternative amputation? Two case reports

Geertzen, Jan H. B.; Jutte, Paul; Rompen, Christiaan; Salvans, Merse

Published in:
Prosthetics and Orthotics International

DOI:
[10.1080/03093640802419163](https://doi.org/10.1080/03093640802419163)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2009

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Geertzen, J. H. B., Jutte, P., Rompen, C., & Salvans, M. (2009). Calcanectomy, an alternative amputation? Two case reports. *Prosthetics and Orthotics International*, 33(1), 78-81. [908802328].
<https://doi.org/10.1080/03093640802419163>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

This article was downloaded by: [University of Groningen]

On: 24 October 2009

Access details: Access Details: [subscription number 906397486]

Publisher Informa Healthcare

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK

Prosthetics and Orthotics International

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t714595820>

Calcanectomy, an alternative amputation? Two case reports

Jan H. B. Geertzen ^a; Paul Jutte ^b; Christiaan Rompen ^b; Merse Salvans ^b

^a Center for Rehabilitation and Graduate School for Health Research, ^b Department of Orthopedic Surgery, University Medical Center Groningen, Groningen, The Netherlands

Online Publication Date: 01 March 2009

To cite this Article Geertzen, Jan H. B., Jutte, Paul, Rompen, Christiaan and Salvans, Merse(2009)'Calcanectomy, an alternative amputation? Two case reports',Prosthetics and Orthotics International,33:1,78 — 81

To link to this Article: DOI: 10.1080/03093640802419163

URL: <http://dx.doi.org/10.1080/03093640802419163>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Calcanectomy, an alternative amputation? Two case reports

JAN H. B. GEERTZEN¹, PAUL JUTTE², CHRISTIAAN ROMPEN², &
MERSE SALVANS²

¹Center for Rehabilitation and Graduate School for Health Research, ²Department of Orthopedic Surgery, University Medical Center Groningen, Groningen, The Netherlands, and ³Rehabilitacio, Hospital Universitari vall d'Hebron, Barcelona, Spain

Abstract

A limb amputation is a traumatic experience for the amputee but it is also a challenge for the recipient to get used to a new situation and reach her/his greatest level of independence. Two patients are presented who had undergone a total calcanectomy. In the first case, a woman with spina bifida aperta L4-L5, calcanectomy was performed to excise a non-healing ulcer and osteomyelitis of the right heel. In the second case, a woman of 72 years of age, calcanectomy was the treatment of the isolated central chondrosarcoma. Both patients were somewhat disabled prior to calcanectomy, one because of spina bifida aperta and the other because of her age. After calcanectomy both patients were able to remain independent for several years. Calcanectomy is an uncommon amputation that may provide better quality of life than other more proximal amputations such as Syme or transtibial amputation. Furthermore more proximal levels of amputation are associated with increased mortality and morbidity. When indicated, calcanectomy should be considered as an alternative type of amputation which provides an effective chance to maintain the patient's independency.

Keywords: *Orthopaedic surgery, calcanectomy*

Introduction

In industrial countries transtibial amputations represent 47% and foot amputations (Syme to transmetatarsal level) represent 6% of all lower limb amputations.¹ The reason for amputation is vascular diseases in 94% of the cases. Trauma and oncology each represent 3% of the reasons to amputate.¹

A transtibial amputation has disadvantages compared to a (partial) foot amputation, such as the loss of length or the higher level of energy consumption. In some cases a calcanectomy should be considered. Calcanectomy was first described in 1931 to treat osteomyelitis of the calcaneus.² The indications for a calcanectomy are osteomyelitis of the calcaneus, intractable ulceration of the heel, deformity of the calcaneus after a severe intra-articular fracture and primary tumours of the calcaneus.³ Data concerning epidemiology of calcanectomy is lacking.

Total resection of the calcaneus is a relatively unknown procedure with a good functional outcome.^{3–6} Sometimes an allograft replacement after total calcanectomy is performed in order to improve the functional outcome.⁷

Benefits of calcanectomy include maintenance of ambulatory status and quality of life, additionally calcanectomy does not alter the next step in the treatment algorithm, which is Syme amputation or transtibial amputation.^{3,7}

The purpose of this report is to present the surgical and functional result of two total calcanectomy procedures as alternatives to transtibial amputation for a non-healing fistula and a chondrosarcoma, both heel sited. Both patients gave written informed consent. The local ethics committee does not require ethical approval for case studies.

Case report

Case 1

A 54-year-old woman born with spina bifida aperta (L4-L5) which was closed a few days after birth presented an ulcer of her right heel. Despite her spina bifida she could walk with a rollator, wearing orthopaedic shoes, for more than 30 min. In 1996 the ulcer was excised from her right heel, after which a fistula existed for more than six years which was not treatable with antibiotics and several plastic surgical reconstructions. Osteomyelitis of the calcaneus developed and transtibial amputation was initially proposed by the orthopaedic surgeon. From 1996 to 2002 the patient was mainly wheelchair-bound. The Spina Bifida multidisciplinary working group, University Medical Center Groningen (UMCG) discussed the suggestion of a transtibial amputation. The possibility of walking was much reduced by paresis of several lower limb muscles. A calcanectomy was suggested, agreed upon and performed in 2002. The recovery period was several months due to wound complications. Since the calcanectomy there were no ulcer recurrences. Currently she is walking again, in and around the house, using a rollator and orthopaedic shoes with an insole for periods of 15 min. She is very satisfied with her current ability to walk which probably would have been lost after a transtibial amputation.

Case 2

A 72-year-old woman came to the hospital in 2001 with a lateral swelling of her right heel. Radiographs showed a lytic lesion with a diameter of 4 cm in the calcaneus with cortical erosion. Computerized tomography suggested soft tissue invasion. Bone scintigraphy revealed a solitary hot spot in the right calcaneus. Chest X-rays were normal. A bone biopsy was performed confirming the suspicion of a chondrosarcoma, histological grade I-II. The first surgical option of the surgeon was transtibial amputation but the orthopaedic surgeon at UMCG who saw the patient for a second opinion proposed a calcanectomy. This procedure was agreed upon by the regional Bone Tumour Board. The calcanectomy was performed and recovery was uneventful. The patient was able to walk using a custom-made orthopaedic shoe and no walking support was needed. During the first two years after the operation she was able to walk for one hour. Three years after calcanectomy bone scintigraphy and X-rays showed no signs of a local recurrence or metastases (Figure 1). Maximal walking time had gradually reduced to 30 min. She experienced some pain due to progressive degeneration of the talonavicular joint of the involved foot.

Currently, six years after the calcanectomy the walking time is about 20 min due to comorbidity. She continues to be very satisfied and wears orthopaedic shoes with a special adapted insole to compensate the leg length difference due to calcanectomy. She walks with a crutch in and around the house and uses a rollator if she walks for more than 15 min. She can walk short distances barefoot in home. The outcome in terms walking ability would have been worse than it is currently if the surgeon had chosen transtibial amputation.

Figure 1. Status after total calcaneectomy in Case 2.

Discussion

Calcaneectomy is presented as a limb salvage procedure that may provide a better quality of life than transtibial amputation. The few reports concerning calcaneectomy are mainly about diabetic or dysvascular patients with untreatable wounds of the feet.

The foot is a rare site for primary and secondary bone neoplasm (<1% of all tumours of the locomotor system).⁸ If the tumour is solely located in the calcaneus a calcaneectomy is an option to be considered.

The more proximal levels of amputation than calcaneectomy are associated with increased mortality, morbidity and increased energy demands.^{3,9} The survival rate after lower limb amputation is poorer with increasing age.⁹

In some cases a partial calcaneectomy can be performed instead of a total calcaneectomy to preserve some of the connecting ligaments resulting in a more stable and functional foot.³ Calcaneectomy ensures walking ability (barefoot walking; toileting in the night) by preserving most of the leg length. An insole in the (customized) shoe may be needed to keep same leg length and provide independent function.^{3,6,10} The total contact inlay must be built up with proper lifting of the heel region to compensate the leg discrepancy, a sufficient buffer heel and with support of the talus. The orthopaedic shoe must have sufficient depth and height to enclose this insole.

After calcaneectomy normal weight-bearing is possible with or without an orthopaedic shoe for small distance walking but for long distances orthopaedic shoes (with insoles) are necessary to prevent wounds.⁶ Although patients present a slight weakness in plantar flexion due to the sub-optimal lever arm after reinsertion of the triceps surae in the dorsal talar region, it is still enough to allow walking barefoot for a short distance.^{5,6} Most patients have no or mild pain after calcaneectomy.^{6,10} Complications are cosmetic deformity (Figure 2), ulceration of the skin, talonavicular subluxation and other late developing

Figure 2. Cosmetic appearance after total calcanectomy in Case 2.

additional deformities.^{6,7} In the cases presented, the outcome is compromised by neurological deficits in one case and morbidity and age in the other.

Both cases presented limitations in walking distances, but both have retained their independency without the need for a prosthesis. With the right indication calcanectomy should be considered as an alternative for Syme or transtibial amputations.

Declaration of interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

References

1. Rommers GM. 2008. Epidemiology of amputations of the lower extremity. In: Geertzen JHB, Rietman JS, editors. *Amputation and prosthetics lower extremity*. Utrecht, The Netherlands: Lemma [in Dutch].
2. Gaenslen FJ. 1931. Split-heel approach in osteomyelitis of os calcis. *J Bone Joint Surg* 13:759–772.
3. Valerie L, Serin S, Kurkoawa CH. 2002. Calcanectomy: A surgical treatment for pathologies affecting the calcaneus. *National Foot Ankle Rev* 8:47–52.
4. Crandall RC, Wagner FW. 1981. Partial and total calcanectomy: A review of 31 consecutive cases over a ten-year period. *J Bone Joint Surg* 63:152–155.
5. Dhillon MS, Singh B, Gill SS, Walker R, Nagi ON. 1993. Management of giant cell tumor of the tarsal bones: A report of nine cases and a review of the literature. *Foot Ankle* 14:5:265–272.
6. Smith DG, Stuck RM, Ketner L, Sage RM, Pinzur MS. 1991. Partial calcanectomy for the treatment of large ulcerations of the heel and calcaneal osteomyelitis. An amputation of the back of the foot. *J Bone Joint Surg* 74A:4:571–576.
7. Muscolo DL, Ayerza MA, Apontetinao LA. 2000. Long-term results of allograft replacement after total calcanectomy. A report of two cases. *J Bone Joint Surg* 82:109–112.
8. Casadei R, Ferraro A, Ferruzzi A, Biagini R, Ruggieri P. 1991. Bone tumors of the foot: Epidemiology and diagnosis. *Chirurgia degli organi di Movimento LXXVI*:76:47–62.
9. Mayfield JA, Reiber GE, Maynard Ch, Czerniecki JM, Caps MT, Sangeorzan BJ. 2001. Survival following lower-limb amputation in a veteran population. *J Rehabil Res Dev* 38:3:341–345.
10. Bollinger M, Thordarson DB. 2004. Partial calcanectomy: An alternative to below the knee amputation. *Exp Clin Endocrinol Diab* 112:9:526–530.