University of Groningen

Interstellar Ices as Witnesses of Star Formation: Selective Deuteration of Water and Organic Molecules Unveiled

Cazaux, S.; Caselli, P.; Spaans, M.

Published in:
Astrophysical Journal Letters

DOI:
[10.1088/2041-8205/741/2/L34](https://doi.org/10.1088/2041-8205/741/2/L34)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2011

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Cazaux, S., Caselli, P., & Spaans, M. (2011). Interstellar Ices as Witnesses of Star Formation: Selective Deuteration of Water and Organic Molecules Unveiled. *Astrophysical Journal Letters*, 741(2), [34].
<https://doi.org/10.1088/2041-8205/741/2/L34>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

INTERSTELLAR ICES AS WITNESSES OF STAR FORMATION: SELECTIVE DEUTERATION OF WATER AND ORGANIC MOLECULES UNVEILED

S. CAZAUX¹, P. CASELLI², AND M. SPAANS¹

¹ Kapteyn Astronomical Institute, P.O. Box 800, 9700AV Groningen, The Netherlands; cazaux@astro.rug.nl

² School of Physics and Astronomy, University of Leeds, LS2 9JT, Leeds, UK
 Received 2011 July 22; accepted 2011 October 3; published 2011 October 20

ABSTRACT

Observations of star-forming environments revealed that the abundances of some deuterated interstellar molecules are markedly larger than the cosmic D/H ratio of 10^{-5} . Possible reasons for this pointed to grain surface chemistry. However, organic molecules and water, which are both ice constituents, do not enjoy the same deuteration. For example, deuterated formaldehyde is very abundant in comets and star-forming regions, while deuterated water rarely is. In this paper, we explain this selective deuteration by following the formation of ices (using the rate equation method) in translucent clouds, as well as their evolution as the cloud collapses to form a star. Ices start with the deposition of gas-phase CO and O onto dust grains. While reaction of oxygen with atoms (H or D) or molecules (H_2) yields H_2O (HDO), CO only reacts with atoms (H and D) to form H_2CO (HDCO, D_2CO). As a result, the deuteration of formaldehyde is sensitive to the gas D/H ratio as the cloud undergoes gravitational collapse, while the deuteration of water strongly depends on the dust temperature at the time of ice formation. These results reproduce well the deuterium fractionation of formaldehyde observed in comets and star-forming regions and can explain the wide spread of deuterium fractionation of water observed in these environments.

Key words: dust, extinction – ISM: abundances – ISM: molecules – stars: abundances – stars: formation

Online-only material: color figures

1. INTRODUCTION

Stars form through the collapse of interstellar clouds which are composed of gas and dust. As the cloud collapses, its density increases, its temperature decreases, and it becomes shielded from external UV radiation that would otherwise dissociate chemical species present in the medium. In these regions, dust grains grow thick icy mantles from the deposition of atomic and molecular species onto their surfaces. This has been confirmed by observations of starless cores (Tafalla et al. 2006), which show that most gas-phase species suffer a significant drop toward the core center. The missing gas species constitute the icy mantles that cover dust grains. As a star forms and heats up its environment, a portion of the icy mantles are released into the gas phase. This phase of star formation, called the hot core (massive stars) or hot corino (low mass stars) phase, exhibits a very complex chemistry rich in oxygen- and nitrogen-bearing molecules (Cazaux et al. 2003), but also shows species such as H_2CO and methanol, which are building blocks for more complex organic molecules essential for the formation of life. Also, while deuterium is known to be 10^5 times less abundant than hydrogen (Linsky 2003), many of these environments present a chemistry rich in highly deuterated species. In particular, deuterated methanol has been found in molecular clouds (Parise et al. 2004) and in low-mass protostars, with a maximum of $\sim 30\%$ for CH_2DOH/CH_3OH (Parise et al. 2002) and of $\sim 3\%$ for CD_3OH/CH_3OH (Parise et al. 2004), while no detection has been reported in comets (Crovisier et al. 2004). Formaldehyde presents a high deuterium fractionation as well (see Table 1); HDCO/ H_2CO is of a few percent toward protostellar cores (Roberts & Millar 2007; Bacmann et al. 2002) and toward hot cores (Turner 1990) and of a tenth of a percent toward hot corinos (Parise et al. 2006). Its doubly deuterated form D_2CO/H_2CO can reach up to 5% in hot corinos (Parise et al. 2006). Species that have been observed with three deuterium atoms (ND_3 ; van der Tak et al. 2002) and CD_3OH (Parise et al.

2004) are expected to be 10^{15} times less abundant than their hydrogen form, with respect to elemental abundances, yet their enhancement can be of 12 orders of magnitude (Ceccarelli et al. 2007).

This high degree of deuteration is attributed to grain surface chemistry. However, while water co-exists with methanol and formaldehyde in ices, it rarely has the same deuteration. In general, the degree of deuteration of organic material is very high compared to that measured in water (Ehrenfreund et al. 2002). The deuterium fractionation of water (Table 1) can be about $HDO/H_2O \sim 0.01\%$ toward massive hot cores (Gensheimer et al. 1996) and $\sim 0.02\% - 0.05\%$ in comets and asteroids (Altwegg & Bockelée-Morvan 2003). These values are similar to standard mean ocean water, which reports a fraction of HDO/H_2O of 0.015% in our oceans. However, some detections showed a high HDO/H_2O ratio of a few percent toward the hot corino NGC1333-IRAS2A (Liu et al. 2011), and IRAS16283-2422 (A. Coutens et al., in preparation; Parise et al. 2005). In this study, we address the selective deuteration of water compared to formaldehyde. We used H_2CO as a representative of organic molecules (H_2CO is a building block of most of the complex organic molecules found in space), but our study also applies to other species formed with CO, such as CH_3OH . First, we determine the chemical gas phase content of the region where ices form. Then, we use a rate equation method that follows the accretion and reactions of gas-phase species that yield ices. Finally, we simulate how the ice compositions change as they undergo gravitational collapse to form stars. These latest results are compared with observations of star-forming environments as well as cometary material.

2. ORIGIN OF THE ICES

Ices are formed in translucent clouds (clouds where H_2 dominates over HI and carbon is converted from CII to CI to CO), in regions that are shielded from UV radiation.

Figure 1. Left panel: atomic (H and D) and molecular (H_2 and HD) hydrogen and deuterium abundances with respect to total hydrogen ($nHI+nH_2$) in a cloud with a density of 10 cm^{-3} (dashed), 100 cm^{-3} (solid), and 10^3 cm^{-3} (dotted) as a function of the visual extinction (mag). Right panel: variation of the D/H ratio as a function of the visual extinction (mag) for densities varying from 10 cm^{-3} to 10^8 cm^{-3} . The line shows $A_V \sim 3$, where the ices initiate.

Table 1
Deuterium Fractionation of Formaldehyde and Water Observed in Different Environments

Environment	Hot Corinos	Hot Cores	Compact Ridges	Comets
Densities (cm^{-3})	$\sim 10^8$ ^a	10^9 – 10^{10} ^b	10^6 ^c	10^{10} ^d
HDCO/ H_2 CO	0.07–0.22 ^e	0.01–0.03 ^e	0.09–0.26 ^f	≤ 0.05 ^g 0.28 ^h
D ₂ CO/ H_2 CO	0.01–0.03 ^e	≤ 0.01 ^e	0.016–0.03 ^f	
HDO/ H_2 O	$\leq 6 \times 10^{-4}$ ⁱ ≥ 0.01 ^l 0.007–0.027 ^m	6×10^{-5} – 5×10^{-4} ^j		3×10^{-4} – 4×10^{-4} ^k

Notes.

^a Ceccarelli (2005).

^b Roberts & Millar (2007).

^c Charnley (1997).

^d Witte et al. (2009).

^e Roberts & Millar (2007), Parise et al. (2006).

^f Turner (1990).

^g Crovisier et al. (2004) Hale-Bopp.

^h Kuan et al. (2008) C/2002 T7.

ⁱ Liu et al. (2011) NGC 1333-IRAS2A.

^j Gensheimer et al. (1996).

^k Meier et al. (1998), Villanueva et al. (2009).

^l Jørgensen & van Dishoeck (2010) NGC 1333-IRAS4B.

^m Herschel results IRAS16293–2422 (A. Coutens et al., in preparation).

At extinctions $A_V \geq 3$ mag, species from the gas phase accrete onto dust grains and initiate the formation of ices (Whittet et al. 2001; Chiar et al. 2011). Therefore, the chemical content of the gas phase at this extinction is crucial for determining which species accrete onto dust and what the composition of ices is. Gas-phase chemistry of dense cores (Roberts et al. 2003) shows that a high D/H ratio can be reached because of the inclusion of ions such as HD_2^+ and D_3^+ in gas-phase chemistry, which produce deuterons upon dissociative recombination. This gas-phase model is able to reproduce a high deuteration of formaldehyde in dense cores. Also, the degree of deuteration of molecular ions is sensitive to the ortho:para H_2 ratio and hence to the chemical and thermal history of the precursor molecular cloud (Flower et al. 2006).

In this study, we concentrate on the ice formation driven by the accretion of gas-phase species. To this effect, we determine

the gas-phase composition (in hydrogen and deuterium) of a medium submitted to UV radiation from neighboring stars, by using a model that couples gas phase and grain surface chemistry (Cazaux & Spaans 2009). This model follows the gas-phase densities of molecular and atomic hydrogen as well as deuterium, as the medium becomes shielded to the external UV field (Draine & Bertoldi 1996). The results reported in Figure 1 show the chemical composition in hydrogen and deuterium as a function of the extinction and therefore provide the initial conditions that are used in the next section (Table 2) for the formation of ices (at $A_V \sim 3$). Note that neither oxygen nor CO or ice chemistry are treated here, our goal being to assess the hydrogen and deuterium content of a medium in the region where ices are forming. As the hydrogen becomes molecular, deuterium can remain atomic, which leads to an enhanced D/H ratio at low extinctions. Then, when the medium is shielded,

Table 2
Initial Conditions of Translucent (at $A_V \sim 3$) and Collapsing Clouds Used in Our Simulations

Env.	n_H	$n_{H I}$	n_{H_2}	n_{DI}	n_{OI}^a	n_{CO}^a	T_{dust}^b	T_{gas}^b
Translucent	10^3	0.5	5×10^2	3×10^{-3}	0.15	0.15	12	20
	10^3	0.5	5×10^2	3×10^{-3}	0.15	0.15	15	30
	10^3	0.5	5×10^2	3×10^{-3}	0.15	0.15	17	70
Collapsing	10^5	0.5	5×10^4	2.5×10^{-2}	0.001	0.001	12	12

Notes. n densities in cm^{-3} , n_H total hydrogen density ($n_{H I} + n_{H_2}$), and T temperatures in K.

^a Hollenbach et al. (2009).

^b Cuppen & Herbst (2007).

the chemistry led by ions drives a D/H ratio that can reach 0.3 (Roberts et al. 2003). In Figure 1, the H/H₂ and D/HD fronts are presented, as well as the D/H ratio as a function of the extinction for an UV radiation field similar to the one in our local interstellar medium (defined as $G_0 = 1$; Habing 1968). The D/H ratio increases with density for extinctions higher than 1 mag. In this study, we consider the formation of ices in translucent clouds with initial gas-phase conditions listed in Table 2.

3. DUST FAVORS DEUTERATION

In our grain surface chemistry model, we consider the formation of ices by accreting species from the gas phase, and let the icy mantles grow until the CO and oxygen from the gas phase are strongly depleted onto dust. Jeans instability, turbulence, and/or triggering due to winds from nearby young stars will then lead to local density increase and gravitational collapse of a cloud fragment. During the collapse, the molecules remaining in the gas can still interfere with the ices and change their composition. Our model is therefore separated into two parts: (1) the slow formation of ices in diffuse environments and (2) the gravitational collapse of dense icy regions of the cloud beginning star formation.

3.1. Translucent Clouds

3.1.1. Gas-phase Composition

Ices form in shielded regions ($A_V \sim 3$ mag), with CO and oxygen equally abundant in the gas phase (Hollenbach et al. 2009) and scale as $1.5 \times 10^{-4} n_H$, where n_H is the total hydrogen density. We assume that ices are first formed in translucent clouds, with initial gas-phase conditions listed in Table 2. As time passes, oxygen and CO will mostly be depleted onto dust, impoverishing the gas phase. Their densities can be calculated as

$$\frac{dn_O}{dt} = 1.5 \times 10^{-4} n_H - O_{dust} \times 4n_{dust} \sigma n_{site}, \quad (1)$$

$$\frac{dn_{CO}}{dt} = 1.5 \times 10^{-4} n_H - CO_{dust} \times 4n_{dust} \sigma n_{site}, \quad (2)$$

where O_{dust} and CO_{dust} are the amount of oxygen and CO on the dust, in monolayers, $n_{sites} \sim 10^{15} \text{ cm}^{-2}$ is the density of sites, n_{dust} is the density of dust (cm^{-3}), and σ is their cross-section (cm^2). Here we consider a grain size distribution that takes into account very small grain particles (Weingartner & Draine 2001) and derive $n_{dust} \sigma / n_H = 2.8 \times 10^{-21} \text{ cm}^2$. With this, we compute the densities of gas-phase CO and oxygen with time and also follow in parallel the chemistry occurring on the dust surfaces.

3.1.2. Grain Surface Chemistry

The gas-phase composition of the environments where ices originate, discussed above, is reported in Table 2. We follow the population of species on the dust by using the rate equation method. Species present in the gas accrete onto dust at a rate:

$$R_{acc_i} = \frac{n_i v_i S}{n_{site}} \text{ monolayer s}^{-1}, \quad (3)$$

where n_i and v_i are, respectively, the densities and velocities of the species i , and S is the sticking coefficient of the species with the dust (here we consider $S = 1$). Since our rate equation model follows the populations of the species of the surface (with one monolayer = 100% coverage), this rate is expressed in monolayer s^{-1} . The binding energies between the chemical species and the grain surface are assumed to be weak (Van der Waals or physisorption), which is typical for icy surfaces. These energies are taken from previous studies (Cuppen & Herbst 2007), with the exception of oxygen, which we take as 1390 K (Bergeron et al. 2008) and H₂, which we assume as 520 K (Dulieu et al. 2005). The binding energies of CO and hydrogenated/deuterated forms are all considered to be 1200 K (Allouche et al. 1998).

The species i present on the surface may return to the gas phase if they evaporate with a rate:

$$R_{evap(i)} = v_i \times \exp\left(-\frac{E_i}{k_B T}\right), \quad (4)$$

where v_i is the oscillation factor of the species i (which is of 10^{12} s^{-1} in physisorbed sites) and E_i is the binding energy of the species i , as reported in Cazaux et al. (2010) and discussed above.

On the surface, atoms and molecules can travel on the dust with a mobility (Cazaux & Tielens 2004; Cazaux et al. 2010):

$$R_i = v_i \times \exp\left(-\frac{E_{pp}}{k_B T}\right), \quad (5)$$

where E_{pp} is the energy of the barrier between two physisorbed sites (considered as 2/3 of the binding energies E_i).

Species on the surface can be photodissociated by UV photons (Cuppen & Herbst 2007) and can meet another species to form a product that will either stay on the surface or be released on the gas phase (Cazaux et al. 2010). In this study, we follow the coverage of the different species in order to determine the ices' composition.

We have constructed a chemical network that considers the hydrogenation and deuteration, but also the H–D and D–H exchanges for H₂CO, HDCO and D₂CO. The most important reactions of the chemical network involving O and CO are

(A color version of this figure is available in the online journal.)

presented in Figure 2 with the associated barriers for the reactions to occur, E_b . The characteristic time for a reaction can be calculated as $\sim 10^{12} \times \exp(-0.406 \times \sqrt{\mu_{\text{red}} \times E_b})$ (oscillation factor times the probability of tunneling through a barrier of 1 Å), with μ_{red} being the reduced mass for the reaction. This reaction occurs if its characteristic time is smaller than the time for the species to leave the site.

The chemistry that involves oxygen has been described in a previous work (Cazaux et al. 2010). For the chemistry of CO, the different barriers for the reactions considered in our model have been derived by several authors (Fuchs et al. 2009; Hidaka et al. 2007; Hidaka et al. 2009). The reactions $\text{H}_2 + \text{CO} \rightarrow \text{H}_2\text{CO}$ as well as $\text{HCO} + \text{H}_2 \rightarrow \text{H}_2\text{CO} + \text{H}$ have very high barriers of 38,400 K and 9000 K (Manion et al. 2008), respectively. Also, the reactions involving D_2 are negligible (Hidaka et al. 2009). Therefore, we do not include reactions involving H_2 , HD, and D_2 in the CO chemistry network.

Recent studies (Ratajczak et al. 2009, 2011) examined exchanges between H and D in ices composed of water and deuterated methanol as these ices warm up. D/H exchanges concern the hydroxyl functional group of methanol and therefore occur between CD_3OD and H_2O that lead to CD_3OH and HDO. In the ices, the amount of CD_3OD is very low compared to the amount of water. These H/D exchanges can be responsible for a very small enhancement of HDO and are therefore not considered in this study.

3.2. Collapse and Star Formation

In the second phase of our model, the dust grains covered by ices are present in denser environments that undergo gravitational collapse. We assume the density of the medium to be $n_{\text{H}} \sim 10^5 \text{ cm}^{-3}$ and our model cloud to collapse at the free-fall rate. The simulations follow the gas density evolution as

$$\frac{dn_{\text{H}}}{dt} = \frac{n_{\text{H}}}{t_{\text{ff}}}, \quad (6)$$

where $t_{\text{ff}} = \sqrt{3\pi/32G\rho} \sim 10^5 \text{ yr}$ is the free-fall time, with G being the gravitational constant and ρ the mass density. Because CO and oxygen are mostly depleted onto the dust, only H, D, H_2 , and HD can accrete onto the icy surfaces with an atomic D/H ratio $\sim 5 \times 10^{-2}$. The initial gas-phase composition of this step of our simulation is reported in Table 2. As the cloud collapses, and the medium becomes denser, the D/H ratio increases as

shown in Figure 1. Molecular hydrogen, which is by far the most abundant molecule in the medium, accretes onto the last layer of ice. Once one layer of H_2 covers the ice (around $n_{\text{H}} \sim 10^6 \text{ cm}^{-3}$), we assume that no further accretion is possible since the binding energies of species with H_2 are very low, and therefore species bounce back into the gas phase. The ice constituents are followed during the collapse of the cloud, as a function of the density of the medium.

4. RESULTS

We have computed the abundance of water and formaldehyde as well as their deuterated forms in interstellar ices.

The oxygen accreting onto the dust transforms into water ice. Figure 3(a) shows the formation of H_2O and HDO on the dust surface as a function of time. The deuteration of water is very sensitive to the dust temperature. At low T_{dust} (12 K), both H_2 and H are present on the ices, and oxygen can react with both H and H_2 . Because H_2 is much more abundant than H, water forms through the reaction $\text{O} + \text{H}_2$ while the formation of HDO involves atomic D. Therefore, the ratio $\text{HDO}/\text{H}_2\text{O}$ (Figure 3(b)) scales with D/ H_2 . At $T_{\text{dust}} = 15 \text{ K}$, short timescales allow an important amount of H atoms to be ice constituents, and oxygen reacts with atomic H to form water. As time proceeds, H atoms disappear by reacting with species present in the ices and H_2 becomes largely dominant. Oxygen atoms coming from the gas phase then react predominantly with H_2 molecules. Therefore, the ratio $\text{HDO}/\text{H}_2\text{O}$ scales with D/H for short timescales and with D/ H_2 for long timescales. At $T_{\text{dust}} = 17 \text{ K}$, H_2 molecules evaporate too fast to allow the reaction $\text{O} + \text{H}_2$ to occur. Water is formed through successive hydrogenation of oxygen, and the ratio $\text{HDO}/\text{H}_2\text{O}$ scales with D/H.

The CO accreting onto the dust transforms into formaldehyde. Figure 3(c) shows the build up of H_2CO , HDCO, and D_2CO ices. The formation of formaldehyde is made through the reactions $\text{CO} + \text{H} \rightarrow \text{HCO}$ and $\text{HCO} + \text{H} \rightarrow \text{H}_2\text{CO}$. The CO ices build up slowly, and CO present on the dust associates with an incoming H atom, or with a D atom. Therefore, $\text{HDCO}/\text{H}_2\text{CO}$ and $\text{D}_2\text{CO}/\text{H}_2\text{CO}$ (Figure 3(d)) scale with D/H and $(\text{D}/\text{H})^2$, respectively.

Once the medium becomes denser, the cloud made of gas and dust (covered by ices) can undergo gravitational collapse and form a star. This star heats up its environment and the content of the ices is released into the gas phase (hot core/corino). At later stages, the star is surrounded by a disk,

Figure 3. H_2O , HDO, H_2CO , HDCO, and D_2CO content on the dust during (1) the formation of ices in a translucent cloud and (2) the gravitational collapse of a cloud to form a star. Lines represent $T_{\text{dust}} = 17\text{ K}$ (solid), 15 K (dotted), and 12 K (dashed). Top: surface coverage in monolayers (%) as a function of time for (a) H_2O and HDO (b) H_2CO , HDCO, and D_2CO . Middle: (c) deuterium fractionation of water and (d) formaldehyde. Bottom panels: model vs. observations of (e) $\text{HDO}/\text{H}_2\text{O}$ (f) $\text{HDCO}/\text{H}_2\text{CO}$ (filled symbols), (f) $\text{D}_2\text{CO}/\text{H}_2\text{CO}$ (open symbols) toward several astrophysical objects as a function of total hydrogen density (Table 1).

(A color version of this figure is available in the online journal.)

and the content of the ices can be present in the warmer parts, or can be locked into cometary material. In these different phases of star formation, deuterated water and formaldehyde have been observed.

Deuterium fractionation of formaldehyde and water has been observed toward several astrophysical objects, as reported in Table 1. While HD₂CO/H₂CO and D₂CO/H₂CO ratios range between 0.01–0.28 and 0.01–0.04, respectively, the HDO/H₂O ratio is much more spread, ranging from a few 10^{−4} to 0.02. In Figures 3(e) and (f), we compare the results of our model to these observations.

Figure 3(f) presents the evolution of the deuterated forms of formaldehyde as the cloud collapses. As the ices are present in a denser environment with a higher D/H ratio, the fraction of deuterated formaldehyde increases by one order of magnitude. Our model reproduces the high degree of deuteration of formaldehyde as the cloud collapses. Also, the fractionation of formaldehyde is not sensitive to the temperature at which the ices formed in the translucent clouds.

For water, on the other hand, Figure 3(e) shows that the ratio HDO/H₂O is constant as the cloud collapses. Once H₂O and HDO are formed, a very important barrier (9600 K) has to be overcome to break these molecules. Therefore, the deuterium fractionation is set during the formation of ices, in the translucent clouds, and deuterated water retains the memory of the ice formation. The HDO/H₂O ratio is extremely sensitive to the dust temperature at which the ices formed in the translucent clouds. At low temperatures, water is formed with O+H₂, while at higher temperatures H₂ evaporate, and water is made through O+H. As a result, our model can explain the spread in the HDO/H₂O ratio observed in different astrophysical objects.

The mantles species that are released in the gas phase could be transformed in other species through gas-phase reactions. Goumans (2011) showed that gas-phase reactions of H₂CO with H(D) would lead to HCO + H₂(HD), while H + D₂CO would lead to CHD₂O. In this case, the deuterium fractionation of formaldehyde could be increased by further reactions in the gas phase. However, Charnley (1997) shows that the HDO/H₂O ratio stays constant after ices evaporate, while HD₂CO/H₂CO could be increased by a factor five. Therefore, the deuterium fractionation found in this study, which is the result of the chemistry occurring in the ices, should not be dramatically changed by the gas-phase chemistry that follows ice evaporation.

5. CONCLUSIONS

We used the rate equation method to simulate the formation of water and formaldehyde, as well as their isotopologues on dust grain surfaces. We model the formation of ices in translucent clouds and incorporate these ices in denser environments that undergo gravitational collapse to form stars.

We find that HD₂CO/H₂CO and D₂CO/H₂CO depend on the gas D/H ratio which increases as the medium collapses and becomes denser.

For water, on the other hand, HDO/H₂O strongly depends on the dust temperatures at which ices form since water formation involves H₂ at low T_{dust} (H₂ very abundant on dust surface) and H at higher T_{dust} (H₂ evaporate). Consequently, the degree of water deuteration is directly related to the physical conditions (i.e., dust temperature) at which ices form in the quiescent cloud just before it collapses to form stars. H₂CO is the simplest

representative of organic species formed via hydrogenation of CO onto dust surfaces. Our results can then be generalized to other species (such as CH₃OH), which also depends on surface hydrogenation of CO. This explains the selective deuteration of water and organic molecules observed in the ISM, similar to that found in comets, IDPs, and chondrites (Ehrenfreund et al. 2002, Figure 9).

The authors thank the anonymous referee for her/his constructive comments. S.C. is supported by the Netherlands Organization for Scientific Research (NWO).

REFERENCES

- Allouche, A., Verlaque, P., & Pourcin, J. 1998, *J. Phys. Chem. B*, 102, 89
 Altwegg, K., & Bockelée-Morvan, D. 2003, *Space Sci. Rev.*, 106, 139
 Bacmann, A., Lefloch, B., Ceccarelli, C., et al. 2002, *A&A*, 389, L6
 Bergeron, H., Rougeau, N., Sidis, V., et al. 2008, *J. Phys. Chem. A*, 112, 11921
 Cazaux, S., Cobut, V., Marseille, M., Spaans, M., & Caselli, P. 2010, *A&A*, 522, A74
 Cazaux, S., & Spaans, M. 2009, *A&A*, 496, 365
 Cazaux, S., & Tielens, A. G. G. M. 2004, *ApJ*, 604, 222
 Cazaux, S., Tielens, A. G. G. M., Ceccarelli, C., et al. 2003, *ApJ*, 593, L51
 Ceccarelli, C. 2005, in IAU Symp. 231, *Astrochemistry: Recent Successes and Current Challenges*, ed. D. C. Lis, G. A. Blake, & E. Herbst (Cambridge: Cambridge Univ. Press), 1
 Ceccarelli, C., Caselli, P., Herbst, E., Tielens, A. G. G. M., & Caux, E. 2007, in *Protostars and Planets V*, ed. B. Reipurth, D. Jewitt, & K. Keil (Tucson, AZ: Univ. Arizona Press), 47
 Charnley, S. B. 1997, *ApJ*, 481, 396
 Chiar, J. E., Pendleton, Y. J., Allamandola, L. J., et al. 2011, *ApJ*, 731, 9
 Crovisier, J., Bockelée-Morvan, D., Colom, P., et al. 2004, *A&A*, 418, 1141
 Cuppen, H. M., & Herbst, E. 2007, *ApJ*, 668, 294
 Draine, B. T., & Bertoldi, F. 1996, *ApJ*, 468, 269
 Dulieu, F., Amiaud, L., Baouche, S., et al. 2005, *Chem. Phys. Lett.*, 404, 187
 Ehrenfreund, P., Irvine, W., Becker, L., et al. 2002, *Rep. Prog. Phys.*, 65, 1427
 Flower, D. R., Pineau Des Forêts, G., & Walmsley, C. M. 2006, *A&A*, 449, 621
 Fuchs, G. W., Cuppen, H. M., Ioppolo, S., et al. 2009, *A&A*, 505, 629
 Gensheimer, P. D., Mauersberger, R., & Wilson, T. L. 1996, *A&A*, 314, 281
 Goumans, T. P. M. 2011, *MNRAS*, 413, 2615
 Habing, H. J. 1968, *Bull. Astron. Inst. Netherlands*, 19, 421
 Hidaka, H., Kouchi, A., & Watanabe, N. 2007, *J. Chem. Phys.*, 126, 204707
 Hidaka, H., Watanabe, M., Kouchi, A., & Watanabe, N. 2009, *ApJ*, 702, 291
 Hollenbach, D., Kaufman, M. J., Bergin, E. A., & Melnick, G. J. 2009, *ApJ*, 690, 1497
 Jørgensen, J. K., & van Dishoeck, E. F. 2010, *ApJ*, 725, L172
 Kuan, Y.-J., Chuang, H.-E., Charnley, S., & Huang, H.-C. 2008, in *COSPAR Scientific Assembly*, 37, HD₂CO in Comet C/2002 T7 (LINEAR), 1640
 Linsky, J. L. 2003, *Space Sci. Rev.*, 106, 49
 Liu, F., Parise, B., Kristensen, L., et al. 2011, *A&A*, 527, A19
 Manion, J. A., Huie, R. E., Levin, R. D., et al. 2008, NIST Chemical Kinetics Database, NIST Standard Reference Database 17, Version 7.0 (Web Version), Release 1.4.3, Data version 2008.12 (Gaithersburg, MD: National Institute of Standards and Technology), <http://kinetics.nist.gov/>
 Meier, R., Owen, T. C., Matthews, H. E., et al. 1998, *Science*, 279, 842
 Parise, B., Castets, A., Herbst, E., et al. 2004, *A&A*, 416, 159
 Parise, B., Caux, E., Castets, A., et al. 2005, *A&A*, 431, 547
 Parise, B., Ceccarelli, C., Tielens, A. G. G. M., et al. 2002, *A&A*, 393, L49
 Parise, B., Ceccarelli, C., Tielens, A. G. G. M., et al. 2006, *A&A*, 453, 949
 Ratajczak, A., Quirico, E., Faure, A., Schmitt, B., & Ceccarelli, C. 2009, *A&A*, 496, L21
 Ratajczak, A., Taquet, V., Kahane, C., et al. 2011, *A&A*, 528, L13
 Roberts, H., Herbst, E., & Millar, T. J. 2003, *ApJ*, 591, L41
 Roberts, H., & Millar, T. J. 2007, *A&A*, 471, 849
 Tafalla, M., Kumar, M. S. N., & Bachiller, R. 2006, *A&A*, 456, 179
 Turner, B. E. 1990, *ApJ*, 362, L29
 van der Tak, F. F. S., Schilke, P., Müller, H. S. P., et al. 2002, *A&A*, 388, L53
 Villanueva, G. L., Mumma, M. J., Bonev, B. P., et al. 2009, *ApJ*, 690, L5
 Weingartner, J. C., & Draine, B. T. 2001, *ApJ*, 548, 296
 Whittet, D. C. B., Gerakines, P. A., Hough, J. H., & Shenoy, S. S. 2001, *ApJ*, 547, 872
 Woitke, P., Kamp, I., & Thi, W. 2009, *A&A*, 501, 383