University of Groningen

Climate change leads to decreasing bird migration distances

Visser, Marcel E.; Perdeck, Albert C.; van Balen, Johan H.; Both, Christiaan

Published in:
Global Change Biology

DOI:
[10.1111/j.1365-2486.2009.01865.x](https://doi.org/10.1111/j.1365-2486.2009.01865.x)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2009

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
Visser, M. E., Perdeck, A. C., van Balen, J. H., & Both, C. (2009). Climate change leads to decreasing bird migration distances. *Global Change Biology*, 15(8), 1859-1865. <https://doi.org/10.1111/j.1365-2486.2009.01865.x>

Copyright
Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taveme" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taveme-amendment>.

Take-down policy
If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supplementary Material - Table

Linear regression of $\log_{10}$ recovery distance (km) in winter of recovery for 24 species (ordered according Euring number), ringed in the Netherlands from 1932 to 2004 during the breeding season (April-July) and recovered during winters 1932/33-2004/05 (Nov-Feb). Included are the species characteristics: habitat, diet, $\log_{10}$ weight, $\log_{10}$ distance in 1984 (see Methods) and trend (population estimate for the Netherlands in 2000 / estimate in 1977). N is the number of recoveries, the estimate is log distance versus year slope (with SE) and the P-value the significance of the slope when species are tested singly in a linear regression.

Latin name	Common name	Habitat	Diet	Log(weight)	Log(distance)	Trend	N	Estimate	SE	P-value
<i>Phalacrocorax carbo</i>	Cormorant	open wet	vertebrate	3.398	2.58	6.33	144	0.001	0.002	0.37
<i>Ardea cinerea</i>	Grey Heron	open wet	vertebrate	3.176	1.57	1.15	1969	-0.003	0.002	0.07
<i>Cygnus olor</i>	Mute Swan	open wet	invert, plant	4.000	1.42	2.4	135	-0.007	0.007	0.33
<i>Somateria mollissima</i>	Eider	open wet	invert, plant	3.352	1.44	2.25	613	-0.004	0.002	0.07
<i>Circus aeruginosus</i>	Marsh Harrier	open wet	vertebrate	2.778	2.25	1.76	52	0.005	0.005	0.31
<i>Circus cyaneus</i>	Hen Harrier	open dry	vertebrate	2.653	1.93	0.83	62	-0.028	0.007	<0.0001
<i>Accipiter nisus</i>	Sparrow Hawk	woodland	vertebrate	2.301	1.42	3.75	818	-0.001	0.002	0.77
<i>Buteo buteo</i>	Buzzard	woodland	vertebrate	2.94	1.59	4.00	911	-0.005	0.002	0.03
<i>Falco tinnunculus</i>	Kestrel	open dry	vertebrate	2.255	1.58	1.09	3146	-0.004	0.001	<0.0001
<i>Fulica atra</i>	Coot	open wet	invert, plant	2.875	1.27	2.38	165	-0.009	0.005	0.10
<i>Haematopus ostralegus</i>	Oystercatcher	open wet	invert, plant	2.763	1.76	2.13	840	-0.006	0.002	0.0015
<i>Recurvirostra avosetta</i>	Avocet	open wet	invert, plant	2.477	2.63	1.60	62	-0.008	0.004	0.06
<i>Vanellus vanellus</i>	Lapwing	open dry	invert, plant	2.301	2.73	2.08	311	-0.009	0.003	0.0006
<i>Larus ridibundus</i>	Black-headed Gull	open wet	invert, plant	2.455	2.22	0.68	1130	-0.007	0.001	<0.0001
<i>Larus canus</i>	Common Gull	open dry	invert, plant	2.602	1.87	0.87	115	-0.017	0.006	0.006
<i>Larus argentatus</i>	Herring Gull	open wet	invert, plant	2.973	1.73	1.23	1394	0.002	0.001	0.07
<i>Tyto alba</i>	Barn Owl	open dry	vertebrate	2.477	1.59	2.61	2571	-0.014	0.001	<0.0001
<i>Athene noctua</i>	Little Owl	open dry	vertebrate	2.279	1.10	0.86	120	-0.006	0.005	0.27
<i>Strix aluco</i>	Tawny Owl	woodland	vertebrate	2.690	1.08	1.82	265	-0.003	0.003	0.47
<i>Asio otus</i>	Long-eared Owl	woodland	vertebrate	2.447	1.41	1.22	283	-0.009	0.003	0.001
<i>Turdus merula</i>	Blackbird	woodland	invert, plant	2.041	1.58	1.50	83	-0.018	0.004	<0.0001
<i>Turdus philomelos</i>	Song Thrush	woodland	invert, plant	1.903	2.46	1.08	59	-0.004	0.003	0.25
<i>Parus major</i>	Great Tit	woodland	invert, plant	1.267	1.12	2.56	226	-0.004	0.002	0.05
<i>Sturnus vulgaris</i>	Starling	open dry	invert, plant	1.903	2.00	0.97	207	-0.007	0.002	0.0001