University of Groningen

Photoswitchable molecular wires

Milder, Maaïke T. W.; Areephong, Jetsuda; Feringa, Ben L.; Browne, Wesley R.; Herek, Jennifer L.

Published in:
Chemical Physics Letters

DOI:
[10.1016/j.cplett.2009.08.013](https://doi.org/10.1016/j.cplett.2009.08.013)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2009

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Milder, M. T. W., Areephong, J., Feringa, B. L., Browne, W. R., & Herek, J. L. (2009). Photoswitchable molecular wires: From a sexithiophene to a dithienylethene and back. *Chemical Physics Letters*, 479(1-3), 137-139. <https://doi.org/10.1016/j.cplett.2009.08.013>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supplementary material

Fig. S1. Absorption spectra of **1cc** at 298 K (line) and 120 K (dashed).

Fig. S2. Transient absorption spectra of the **1cc** in isopentane $T=125$ K. Pump 620 nm, 33 fs, 30 μ W. Panel A shows the spectral evolution of the closed form at 1 ps (black squares), 2 ps (red circles), 5 ps (green triangles), and 10 ps (blue stars). The steady state absorption spectrum is presented scaled and in grey shading. The symbols represent the spectra from global analysis of the data. Panel B shows the temporal evolution of the transient signals at 650 nm (black squares), 750 nm (red triangles). As a comparison panel B shows the time evolution of the open form at 515 nm (green squares). The solid lines are global fits of the data.