University of Groningen

BVMO-catalysed dynamic kinetic resolution of racemic benzyl ketones in the presence of anion exchange resins

Rodriguez, Cristina; de Gonzalo, Gonzalo; Rioz-Martinez, Ana; Pazmino, Daniel E. Torres; Fraaije, Marco W.; Gotor, Vicente

Published in:
Organic & Biomolecular Chemistry

DOI:
[10.1039/b922693a](https://doi.org/10.1039/b922693a)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2010

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Rodriguez, C., de Gonzalo, G., Rioz-Martinez, A., Pazmino, D. E. T., Fraaije, M. W., & Gotor, V. (2010). BVMO-catalysed dynamic kinetic resolution of racemic benzyl ketones in the presence of anion exchange resins. *Organic & Biomolecular Chemistry*, 8(5), 1121-1125. <https://doi.org/10.1039/b922693a>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

BVMO-catalysed dynamic kinetic resolution of racemic benzylketones in presence of anion exchange resins

Cristina Rodríguez,^a Gonzalo de Gonzalo,^a Ana Rioz-Martínez,^a Daniel E. Torres Pazmiño,^b Marco W. Fraaije^b and Vicente Gotor^{*,a}

^a Departamento de Química Orgánica e Inorgánica, Instituto de Biotecnología de Asturias, Universidad de Oviedo, c/ Julián Clavería 8, 33006, Oviedo, Spain.

e-mail: vgs@fq.uniovi.es

^b Laboratory of Biochemistry, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands.

Supporting Information

Contents

1. Characterization data of compounds.....	S2
2. Experimental data.....	S4
3. NMR Spectra.....	S6

1. Characterization data of compounds.

(S)-1-Phenylethyl acetate (1b, colourless oil). (38.2 mg, 69% yield); δ_{H} (300.13 MHz, CDCl_3 , Me_4Si) 1.56 (3H, d, J 6.6 Hz), 2.10 (3H, s), 5.91 (1H, q, J 6.6 Hz), 7.28-7.39 (5H, m); δ_{C} (75.5 MHz, CDCl_3 , Me_4Si) 20.9, 21.7, 71.8, 125.6, 125.7, 127.4, 128.0, 128.1, 141.2, 169.8; m/z (APCI^+) 165 ($\text{M}+\text{H}^+$, 100%). Determination of the *ee* by GC analysis: RT β DEXse, 70°C (5 min), 1°C/min, 120°C (5 min), t_{R} (R) 50.5 min; t_{R} (S) 42.9 min.

(S)-1-(3-Methylphenyl)ethyl acetate (2b, colourless oil). (32.9 mg, 60% yield); δ_{H} (300.13 MHz, CDCl_3 , Me_4Si) 1.53 (3H, d, J 6.6 Hz), 2.08 (3H, s), 2.36 (3H, s), 5.85 (1H, q, J 6.6 Hz), 7.09-7.25 (4H, m); δ_{C} (75.5 MHz, CDCl_3 , Me_4Si) 21.3, 21.4, 22.1, 72.3, 123.0, 126.7, 128.3, 128.6, 138.1, 141.5, 170.3; m/z (EI^+) 178 (M^+ , 28%), 136 (100%), 117 (91%). Determination of the *ee* by GC analysis: Chiralsil DexCB, 70°C, 1°C/min, 120°C (5 min), t_{R} (R) 32.7 min; t_{R} (S) 29.0 min.

(S)-1-(3-Trifluoromethylphenyl)ethyl acetate (3b, colourless oil). (40.3 mg, 75% yield); δ_{H} (300.13 MHz, CDCl_3 , Me_4Si) 1.55 (3H, d, J 6.8), 2.10 (3H, s), 5.91 (1H, q, J 6.6), 7.47- 7.61 (4H, m); δ_{C} (75.5 MHz, CDCl_3 , Me_4Si) 21.2, 22.2, 71.5, 122.7 (d, J 3.76 Hz), 124.0 (d, J 270.6 Hz), 124.6 (d, J 3.6 Hz), 128.9, 129.5, 130.9 (d, J 32.0 Hz), 142.7, 170.1; m/z (EI^+) 232 (M^+ , 11%), 190 (90%). Determination of the *ee* by GC analysis: RT β DEXse, 100°C (20 min), 2°C/min, 150°C (5 min), t_{R} (R) 22.8 min; t_{R} (S) 18.3 min.

(S)-1-(4-Chlorophenyl)ethyl acetate (4b, yellow pale oil). (32.6 mg, 60% yield); δ_{H} (300.13 MHz, CDCl_3 , Me_4Si) 1.53 (3H, d, J 6.6 Hz), 2.09 (3H, s), 5.86 (1H, q, J 6.6 Hz), 7.28- 7.36 (4H, m); δ_{C} (75.5 MHz, CDCl_3 , Me_4Si) 21.2, 22.1, 71.5, 127.4, 127.5, 128.6, 128.7, 133.5, 140.1, 170.1; m/z (ESI^+) 221 ($\text{M}+\text{Na}^+$, 100%). Determination of the *ee* by GC analysis: RT β DEXse, 110°C (10 min), 1°C/min, 140°C (5 min), t_{R} (R) 38.5 min; t_{R} (S) 34.6 min.

(S)-1-Phenylethyl propionate (5b, colourless oil). (32.4 mg, 59% yield); δ_{H} (300.13 MHz, CDCl_3 , Me_4Si) 1.06 (3H, t, J 7.4 Hz), 1.45 (3H, d, J 6.6 Hz), 2.22-2.31 (2H, m), 5.81 (1H, q, J 6.6 Hz), 7.17-7.27 (5H, m); δ_{C} (75.5 MHz, CDCl_3 , Me_4Si) 9.0, 22.2, 27.8, 72.0, 126.0, 126.1, 127.7, 128.2, 128.3, 141.8, 173.6; m/z (APCI^+) 179 ($\text{M}+\text{H}^+$, 100%).

Determination of the *ee* by GC analysis: Rt β DEXse, 70 °C (5 min), 3°C/ min, 200 °C (5 min): t_R (*R*) 29.3 min; t_R (*S*) 28.9 min.

(*S*)-1-Phenylpropyl acetate (6b, colourless oil): (34.6 mg, 63% yield); δ_H (300.13 MHz, CDCl₃, Me₄Si) 0.88 (3H, t, *J* 7.4 Hz), 1.81-1.95 (2H, m), 2.08 (3H, s), 5.67 (1H, t, *J* 6.9 Hz), 7.28-7.37 (5H, m); δ_C (75.5 MHz, CDCl₃, Me₄Si) 9.8, 21.2, 29.2, 77.3, 126.5, 126.6, 127.7, 128.3, 128.4, 140.4, 170.4; m/z (APCI⁺) 179 (M+H⁺, 100%).
Determination of the *ee* by GC analysis: Rt β DEXse, 110 °C isotherm: t_R (*R*) 26.6 min; t_R (*S*) 23.7 min.

(*S*)-1-Phenylbutyl propionate (7b, yellow pale oil). (20.6 mg, 38% yield); δ_H (300.13 MHz, CDCl₃, Me₄Si) 0.85 (3H, t, *J* 7.2 Hz), 1.07 (3H, t, *J* 7.4 Hz), 1.15-1.34 (2H, m), 1.61-1.73 (1H, m), 1.77-1.90 (1H, m), 2.24-2.33 (2H, m), 5.69 (1H, t, *J* 7.0 Hz), 7.20-7.28 (5H, m); δ_C (75.5 MHz, CDCl₃, Me₄Si) 9.1, 13.7, 18.7, 27.8, 38.5, 75.6, 126.4, 126.5, 127.7, 128.2, 128.3, 140.9, 173.7; m/z (ESI⁺) 229 (M⁺+Na, 100%).
Determination of the *ee* by GC analysis: Hydrodex β -TBOAc, 50 °C (10 min), 1°C/min, 120°C (5 min): t_R (*R*) 78.4 min; t_R (*S*) 78.8 min.

2. Experimental data.

Racemisation experiments were performed by dissolving optically active 3-phenylpentan-2-one (*R*)-**1a** (10 mM) isolated from the preparative kinetic resolution of ($\pm$)-**1a** performed at pH 8.0 and 20°C with HAPMO, in Tris/HCl buffer containing the anion exchange resins at the selected conditions (pH and temperature). The solution was shaken for different reaction times and aliquots were taken, extracted with ethyl acetate, dried onto Na₂SO₄ and analyzed by GC in order to determine the enantiomeric excesses. Results are summarized in Figure 1.

Figure 1. Racemisation of ketone (*R*)-**1a** (10 mM) when dissolved in different reaction media: Tris/HCl pH 8.0 (◇), Tris/HCl pH 10.0 (□) and Tris/HCl pH 10.0 containing: Amberlite IRA-440C (Δ), Lewatit MP62 (■) or Dowex MWA-1 (○).

Racemisation is supposed to be a pseudo-first order reaction. Relative initial rate constants (k_{rac}) were determined from the enantiomeric excesses of ketone **1a** according to Equation 1:

$$ee(t) = ee(0)e^{-2(k_{\text{rac}})t}$$

where $ee(t)$ is the enantiomeric excess of **1a** at the time established, $ee(0)$ the initial optical purity of **1a**, k_{rac} is the relative initial racemisation constant and t is the time of measurement.

Racemisation studies were also performed with optically pure (*S*)-1-phenylethyl acetate (*S*)-**1b**. This compound was dissolved in Tris/HCl buffer at different reaction conditions (pH and anion exchange resins). The solution was shaken for different reaction times

and aliquots were taken, extracted with ethyl acetate, dried onto Na₂SO₄ and analyzed by GC in order to determine the enantiomeric excesses. No change in optical purity of (*S*)-**1b** was observed after 120 h at all the conditions tested.

3. NMR Spectra

(±)-1b

(±)-1b

(±)-1b

(±)-4b

(±)-4b

(±)-4b

(±)-5b

(±)-5b

(±)-5b

(±)-6b

(±)-6b

(±)-6b

(±)-7b

(±)-7b