University of Groningen

An efficient yttrium catalysed version of the "Aufbaureaktion" for the synthesis of terminal functionalised polyethylene

Kretschmer, Winfried P.; Bauer, Tobias; Hessen, Bart; Kempe, Rhett

Published in:
Dalton Transactions

DOI:
[10.1039/c002499f](https://doi.org/10.1039/c002499f)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2010

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Kretschmer, W. P., Bauer, T., Hessen, B., & Kempe, R. (2010). An efficient yttrium catalysed version of the "Aufbaureaktion" for the synthesis of terminal functionalised polyethylene. *Dalton Transactions*, 39(29), 6847-6852. <https://doi.org/10.1039/c002499f>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supplementary material

An efficient yttrium catalysed version of the “Aufbaureaktion” for the synthesis of terminal functionalised PE

Winfried P. Kretschmer,^{*a,b} Tobias Bauer,^b Bart Hessen^a and Rhett Kempe^{*b}

^a Stratingh Institute for Chemistry, Centre for Catalytic Olefin Polymerisation, University of Groningen, Nijenborgh 4 NL-9747 AG Groningen, The Netherlands. Fax: (+49) 921 55 2157; Tel: (+49) 921 55 2580; E-mail: Winfried.Kretschmer@uni-bayreuth.de

^b Lehrstuhl Anorganische Chemie II, Universität Bayreuth, 95440 Bayreuth, Germany. Fax: (+49) 921 55 2157; Tel: (+49) 921 55 2540; E-mail: Kempe@uni-bayreuth.de

Figure 1. ^1H -NMR spectrum ($\text{C}_2\text{D}_2\text{Cl}_4$, 393K) of PE yielded after polymerization with TIBA chain transfer reagent and oxidative work up ($M_w = 2160$).

Figure 2. ^1H -NMR spectrum ($\text{C}_2\text{D}_2\text{Cl}_4$, 393K) of PE yielded after polymerization with TIBA chain transfer reagent and oxidative work up ($M_w = 1460$).

Figure 3. ^1H -NMR spectrum ($\text{C}_2\text{D}_2\text{Cl}_4$, 393K) of bromo terminated polyethylene.*
unidentified impurity

Figure 4. ^1H -NMR spectrum ($\text{C}_2\text{D}_2\text{Cl}_4$, 393K) of chloro sulfite terminated polyethylene.

Figure 5. ^1H -NMR spectrum ($\text{C}_2\text{D}_2\text{Cl}_4$, 393K) of α -bromo-acetate terminated polyethylene.

Figure 6. ^1H -NMR spectrum ($\text{C}_2\text{D}_2\text{Cl}_4$, 393K) of α -bromo-isobuturate terminated polyethylene.

Figure 7. ^1H -NMR spectrum ($\text{C}_2\text{D}_2\text{Cl}_4$, 393K) of 1-(1-PE-oxycarbonyl-methyl)-3-methyl-imidazoliumbromid.

Figure 8. ^1H -NMR spectrum ($\text{C}_2\text{D}_2\text{Cl}_4$, 393K) of 1-(1-PE-oxycarbonyl-methyl)-3-(2,4,6-trimethyl-phenyl)-imidazoliumbromid.

Figure 9. ^1H -NMR spectrum ($\text{C}_2\text{D}_2\text{Cl}_4$, 393K) of 1-(1-PE-oxycarbonyl-methyl)-3-(2,6-di-*iso*-propyl-phenyl)-imidazoliumbromid.