University of Groningen

Exploiting donor-acceptor interactions in aqueous dynamic combinatorial libraries

Au-Yeung, Ho Yu; Cougnon, Fabien B. L.; Otto, Sijbren; Pantos, G. Dan; Sanders, Jeremy K. M.; Pantoş, G. Dan

Published in:
Chemical Science

DOI:
[10.1039/c0sc00307g](https://doi.org/10.1039/c0sc00307g)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2010

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Au-Yeung, H. Y., Cougnon, F. B. L., Otto, S., Pantos, G. D., Sanders, J. K. M., & Pantoş, G. D. (2010). Exploiting donor-acceptor interactions in aqueous dynamic combinatorial libraries: exploratory studies of simple systems. *Chemical Science*, 1(5), 567-574. <https://doi.org/10.1039/c0sc00307g>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supporting information for:

Exploiting Donor–Acceptor Interactions in Aqueous Dynamic Combinatorial Libraries: exploratory studies of simple systems

Ho Yu Au-Yeung,^a Fabien B. L. Cougnon,^a Sijbren Otto,^b G. Dan Pantoş^a and Jeremy K. M. Sanders^a

^a University Chemical Laboratory, University of Cambridge, Lensfield Road, CB2 1EW, UK

^b Centre for Systems Chemistry, Stratingh Institute, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands.

Email: gdp26@cam.ac.uk, jkms@cam.ac.uk

Fax: 44 (0)1223 336017

Synthesis	S2
DCL setup and HPLC/LCMS methods	S9
HPLC analysis of DCLs	S11
MS spectra	S15
NMR spectra	S21
References	S23

Synthesis of **2**:

To a Schlenk flask charged with **C** (0.28 g, 0.26 mmol) was added degassed TFA (3 ml, 40 mmol). The solution was stirred under N₂ at room temperature for 1.5 hour and added SiEt₃H (0.2 ml, 1.3 mmol). Stirring was continued for 30 minutes. Volatiles were removed *in vacuo*. The solid left was washed with Et₂O (20 ml) and collected by filtration. Yield: 0.13 g, 87%. M.p.: > 250 °C. ¹H NMR (400 MHz, DMSO-*d*₆, 300 K) δ (ppm): 12.93 (br, 2 H, COOH), 8.73 (s, 4 H, NDI), 8.68 (d, *J* = 7.9 Hz, 2 H, NH), 4.82 (d, *J* = 16 Hz, 2 H, Gly), 4.77 (d, *J* = 16 Hz, 2 H, Gly), 4.48–4.43 (m, 2 H, α-Cys), 2.91–2.84 (m, 2 H, β-Cys), 2.81–2.73 (m, 2 H, β-Cys), 2.40 (t, *J* = 8.4 Hz, 2 H, SH). ¹³C{¹H} NMR (100.60 MHz, DMSO-*d*₆, 300 K) δ (ppm): 171.7, 162.8, 152.4, 131.2, 129.4, 126.6, 54.9, 42.9, 26.0. HRMS (ESI+) calcd. for C₂₄H₂₁N₄O₁₀S₂ [M+H]⁺ (*m/z*): 589.0699, found: 589.0718.

• Synthesis of NDI building block **3**

NDI dithiol building block **3** was synthesised according to the following scheme:

Trityl protected glutathione was synthesised according to the literature.⁶

COOH), 8.75 (s, 4 H, NDI), 8.23 (t, $J = 6$ Hz, 2 H, NH-Gly), 7.90 (d, $J = 8$ Hz, 2 H, NH-Cys), 5.58 (dd, $J = 9$ Hz, 5 Hz, 2 H, α -Glu), 4.35 (dd, $J = 13$ Hz, 7 Hz, 2 H, α -Cys), 3.73 (dd, $J = 11$ Hz, 6 Hz, 4 H, Gly), 2.80–2.63 (m, 4 H), 2.42–2.24 (m, 10 H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100.62 MHz, DMSO- d_6 , 300 K): 173.4, 171.8, 171.3, 170.8, 170.4, 162.8, 131.6, 65.3, 56.7, 54.8, 32.0, 26.7, 24.4, 15.5. HRMS (ESI+) calcd. for $\text{C}_{34}\text{H}_{35}\text{N}_6\text{O}_{16}\text{S}_2$ $[\text{M}+\text{H}]^+$ (m/z): 847.1545, found: 847.1571.

• Synthesis of DN building block 4

DN dithiol building block **4** was synthesised according to the following scheme:

Synthesis of DN ester E:

To a solution of 1,5-dihydroxynaphthalene (1.61 g, 10 mmol) in acetone (200 ml) was added finely ground K_2CO_3 (7 g, 50 mmol) and methyl bromoacetate (2.2 ml, 22 mmol). The mixture was heated to reflux for 8 hours. The resulting solution was filtered, Et_2O (100 ml) was added, and washed successively with 1 M HCl (50 ml), water (50 ml) and brine (50 ml). Evaporation of solvents from the organic layer gave the product as a yellow crystalline solid which was used in the next step without further purification. Yield: 3.06 g, quant. ^1H NMR (400 MHz, CDCl_3 , 300 K) δ (ppm): 7.98 (d, $J = 8.4$ Hz, 2 H, DN), 7.36 (t, $J = 7.6$ Hz, 2 H, DN), 6.76 (d, $J = 7.2$ Hz, 2 H, DN), 4.81 (s, 4 H, CH_2), 3.82 (s, 6 H, CH_3).

Synthesis of trityl protected dithiol H:

To a mixture of **G** (0.47 g, 1 mmol) and *S*-trityl-L-cysteine (0.80 g, 2.2 mmol) in DMF (50 ml) was added Et₃N (0.3 ml, 2.2 mmol). The solution was stirred under N₂ for 8 hours at room temperature. Solvent was removed and the residue re-dissolved in acetone (10 ml). The acetone solution was added dropwise to a vigorously stirred solution of 1 M HCl (200 ml). The yellowish-brown solid was collected by filtration and dried. Yield: 0.79 g, 82%. M.p.: 207–209 °C. ¹H NMR (400 MHz, DMSO-*d*₆, 300 K) δ (ppm): 12.91 (br, 2 H, COOH), 8.43 (d, *J* = 8.4 Hz, 2 H, NH), 7.91 (d, *J* = 8.4 Hz, 2 H, DN), 7.34–7.22 (m, 32 H, Trt, DN), 6.97 (d, *J* = 7.6 Hz, 2 H, DN), 4.73 (s, 4 H, OCH₂), 4.33 (dt, *J* = 5.2 Hz, 8.4 Hz, 2 H, α), 2.70 (dd, *J* = 8.4 Hz, 12.0 Hz, 2 H, β); ¹H NMR (400 MHz, acetone-*d*₆, 300 K) δ (ppm): 7.98 (d, *J* = 8.0 Hz, 2 H, NH), 7.74 (d, *J* = 8.0 Hz, 2 H, DN), 7.40–7.21 (m, 32 H, Trt, DN), 7.05 (d, *J* = 7.6 Hz, 2 H, DN), 4.75 (s, 4 H, OCH₂), 4.63 (ddd, *J* = 8.0 Hz, 6.8 Hz, 5.2 Hz, 2 H, α), 2.84 (dd, *J* = 12 Hz, 6.8 Hz, 2 H, β), 2.72 (dd, *J* = 12 Hz, 5.2 Hz, 2 H, β). ¹³C{¹H} NMR (100.62 MHz, DMSO-*d*₆, 300 K) δ (ppm): 171.3, 167.4, 153.0, 144.2, 129.1, 128.1, 126.8, 125.8, 125.3, 114.8, 106.7, 67.2, 66.2, 50.8, 33.0. HRMS (ESI⁺) calcd. for C₅₈H₅₀N₂O₈S₂Na [M+Na]⁺ (*m/z*): 989.2906, found: 989.2911.

Synthesis of DN dithiol 4:

To a Schlenk tube charged with **H** (0.48 g, 0.5 mmol) was added degassed TFA (5 ml, 65 mmol) under N₂. The solution was stirred at room temperature for 1.5 hour, and triethylsilane added (0.4 ml, 2.5 mmol). Immediate precipitation was observed, and the mixture was stirred for an additional 30 minutes. Volatiles were removed *in vacuo*. The resulting yellow solid was washed with Et₂O (50 ml). Yield: 0.23 g, 94%. M.p.: 175–179 °C (decomposed). ¹H NMR (400 MHz, DMSO-*d*₆, 300 K) δ (ppm): 13.01 (br, 2 H, COOH), 8.37 (d, *J* = 7.6 Hz, 2 H, NH), 7.91 (d, *J* = 8.4 Hz, 2 H, DN), 7.42 (t, *J* = 8.0 Hz, 2 H, DN), 7.00 (d, *J* = 8.0 Hz, 2 H, DN), 4.78 (s, 4 H, OCH₂), 4.55 (dt, *J* = 7.2 Hz, 4.8 Hz, 2 H, α), 3.02–2.95 (m, 2 H, β), 2.93–2.86 (m, 2 H, β), 2.43 (t, *J* = 8.4 Hz, 2 H, SH). ¹³C{¹H} NMR (100.62 MHz, DMSO-*d*₆, 300 K) δ (ppm): 171.3, 167.6, 153.0, 125.8, 125.4, 114.7, 106.6, 67.2, 53.9, 25.4. HRMS (ESI⁺) calcd. for C₂₀H₂₃N₂O₈S₂ [M+H]⁺ (*m/z*): 483.0896, found: 483.0907.

DCL setup and HPLC/LCMS methods

A typical analytical DCL was prepared in 1 ml scale by dissolving the building blocks in 10 mM NaOH to give a 5 mM solution, followed by titration with 100 mM NaOH to pH 8. Templates or salt (NaNO_3) were added either as a concentrated solution or directly in solid form where appropriate. The library solutions were stirred in close-capped vials for at least 5 days before HPLC or LCMS analysis. Preparative DCLs were prepared using the same method in 10 ml scale.

HPLC grade MeOH (Fisher or Romil) and formic acid (FA) (Fluka or Romil) were used without further purification. MilliQ water was obtained from a Millipore water purification system. HPLC analyses were carried out on Hewlett Packard 1050 or 1100 systems coupled to diode array or multiple wavelength UV/Vis detectors. Data was processed using HP ChemStation software. In general, analytical separations were achieved by injecting 5 μl aliquots of 5 mM library solution onto a reverse phase Symmetry C_8 column (150 x 4.6 cm, 3 μm particle size). Separations were achieved by running the column either with THF/water, acetonitrile/water, or MeOH/water as eluent at a flow rate of 1 ml/min at 32.5 °C. UV/Vis absorbance was monitored at 292 nm (for 1,5-DN containing DCLs) and 383 nm (for NDI containing DCLs).

Preparative separations were performed on a HP 1050 system coupled to a single variable wavelength UV detector. Samples were injected onto a reverse phase SymmetryPrep C_{18} column (300 x 7.8 mm, 7 μm particle size) by using a Gilson 234 auto-injector. Solvent systems used were the same as that for the analytical separation with elution profiles modified from the corresponding analytical methods. Fractions were manually collected and combined. Solvents were removed from the combined fractions by a rotary evaporator and the isolated sample stored in a fridge.

LCMS was carried out on an Agilent 1100 LC/MSD trap XCT system. Data was processed using ChemStation and LC/MSD trap software. Influx of analyte solution was maintained at a flow rate of 50 $\mu\text{l}/\text{min}$ and ionised with an electrospray source (ESI). ESI-MS spectra (negative ion) were acquired in ultra scan mode with drying temperature of 350 °C, nebuliser pressure of 18 psi, drying gas flow of 5 l/min. Capillary voltage was set to 4000 V. An ICC target of 20,000 ions or accumulation time of 100 ms, and a target mass of 1000 was set for the trap. Unless otherwise stated, dissociation amplitude of 1 V was used for MS/MS studied.

Analytical HPLC methods for DCLs of NDIs

Analytical HPLC method for DCL containing only NDI 1 (5 mM total):

Time/min	Water (0.1% FA)	THF
0	60%	40%
40	60%	40%

Analytical HPLC method of the other NDI DCLs (5 mM total):

Time/min	Water (0.1% FA)	Acetonitrile (0.1% FA)
0	90%	10%
30	30%	70%
40	30%	70%

• Analytical HPLC methods for DCLs of DNs

Analytical HPLC method for DCLs containing either 4 or 5 (5 mM total):

Time/min	Water (0.1% FA)	MeOH (0.1% FA)
0	40%	60%
9	40%	60%
20	0%	100%
30	0%	100%

Analytical method for DCL containing both 4 and 5 (1:1, 5 mM total):

Time/min	Water (0.1% FA)	MeOH (0.1% FA)
0	45%	55%
6	45%	55%
7	40%	60%
14	40%	60%
20	20%	80%
30	20%	80%

• Preparative HPLC methods

Preparative HPLC methods:

DCL	NDI 1	DN 4 or 5
Injection volume	200 µl	300 µl
Flow rate	2.5 ml/min	3 ml/min
Temperature	45 °C	32.5 °C
Elution	45% THF/water (0.1% FA, isocratic elution)	65% MeOH/water (0.1% FA, isocratic elution)
Run time	35 minutes	20 minutes

HPLC analysis of DCLs

- HPLC analysis of DCLs of NDI building block 2

Figure S1. HPLC analysis of 5 mM DCL of **2** (5 mM). No change was observed upon addition of template (2.5 mM) or NaNO₃ (1M).

- HPLC analysis of DCLs of NDI building block 3

Figure S2. HPLC analysis of 5 mM DCL of **3** (5 mM). No change was observed upon addition of template (2.5 mM) or NaNO₃ (1M).

- HPLC analysis of DCLs of NDI building blocks 1 and 2

Figure S3. HPLC analysis of 5 mM DCL of **1** and **2** (1:1 molar ratio, 5 mM total).

Table S1. Amplification factors of library members of DCL of **1** and **2** (1:1 molar ratio, 5 mM total, 2.5 mM guest). In the presence of the other templates, no change in the library distribution was observed.

Guest	6	7	9	11
G1	0.8	0.8	0.8	1.2
G2	0.1	0.2	0.4	1.9

• HPLC analysis of DCLs of NDI building blocks 1 and 3

Figure S4. HPLC analysis of 5 mM DCL of **1** and **3** (1:1 molar ratio, 5 mM total).

Table S2 Amplification factors of library members of DCL of **1** and **3** (1:1 molar ratio, 5 mM total, 2.5 mM guest). In the presence of the other templates, no change in the library distribution was observed.

Guest	6	10	I
G1	0.6	0.9	12.5
G2	0.5	0.1	13.0

• HPLC analysis of DCLs of NDI building blocks 2 and 3

Figure S5. HPLC analysis of 5 mM DCL of **2** and **3** (1:1 molar ratio, 5 mM total).

Table S3 Amplification factors of library members of DCL of **2** and **3** (1:1 molar ratio, 5 mM total, 2.5 mM guest). In the presence of the other templates, no change in the library distribution was observed.

Guest	9	10	J
G2	0.7	0.8	1.6

HPLC analysis of DCLs of NDI building blocks 1, 2 and 3

Figure S6. HPLC analysis of 5 mM DCL of **1**, **2** and **3** (1:1:1 molar ratio, 5 mM total).

Table S4. Amplification factors of library members of DCL of **1**, **2** and **3** (1:1:1 molar ratio, 5 mM total, 2.5 mM guest).

Guest	6	7	9	10	11
G1	0.9	0.7	0.8	1	1.2
G2	0.2	0.2	0.1	1	2.3

• HPLC analysis of DCLs of DN building blocks

Figure S7. HPLC analysis of 5 mM DCL of **5** in (a) the absence of salt, (b) the presence of 1 M NaNO₃.

Figure S8. HPLC analysis of 5 mM DCL of **4** and **5** (1:1 molar ratio, 5 mM total) in (a) the absence of salt, (b) the presence of 1 M NaNO₃.

Table S5. Amplification factors of library members of DCL of **4** (5 mM, 2.5 mM guest).

Guest	Monomer 12	Dimer 13	Trimer 14
G5	0.9	1.3	1.1
G6	1.0	1.1	1.0
G7	1.0	0.9	1.0

Table S6. Amplification factors of library members of DCL of **4** and **5** (1:1 molar ratio, 5 mM, 2.5 mM guest).

Guest	Monomer		Dimer			Trimer		
	12	15	13	16	18	14	19	20
G6	0.8	0.9	0.9	1.1	1.4	1.2	1.0	1.1
G8	0.9	0.6	0.8	1.4	1.5	2.0	2.0	1.5

MS spectra

• MS spectra of NDI macrocycles

Figure S9. ESI-MS spectra of NDI macrocycles (a) **6**, (b1) minor isomer of **7** at 19 min, (b2) major isomer of **7** at 23 min, and (c) **8** (as dianion). Expansions of the molecular ion are shown as insets.

Figure S10. ESI-MS spectra of the NDI intermediates observed during equilibration (a) the dithiol building block **1**, and (b) the linear dimer **6'**. Expansions of the molecular ions are shown as insets.

Figure S11. ESI-MS spectra of NDI macrocycles (a) **9**, and (b) **10**. Expansions of the molecular ion are shown as insets.

Figure S12. ESI-MS spectra of NDI macrocycles (a) **11**, (b) **I**, and (c) **J**. Expansions of the molecular ion are shown as insets.

• MS spectra of DN macrocycles

Figure S13. ESI-MS spectra of the DN macrocycles (a) **12**, (b) **13**, and (c) **14**. Expansions of the molecular ions are shown as insets.

Figure S14. ESI-MS spectra of the DN macrocycles (a) **15**, (b) **16**, and (c) **17**. Expansions of the molecular ions are shown as insets.

Figure S15. ESI-MS spectra of the mixed DN macrocycles (a) **18**, (b) **19**, and (c) **20**. Expansions of the molecular ions are shown as insets.

Figure S16. (a) ESI-MS and (b) MS/MS spectra of the DN tetramer **21**. Expansion of the molecular ion is shown as inset.

NMR spectra

For NMR characterisation of NDI **1** and DN **5**, please refer to our previous communications.^{2, 3}

For NMR characterisation of NDI macrocycles **6–8**, please refer to our previous communication.²

References

1. P. Pengo, G. D. Pantoş, S. Otto and J. K. M. Sanders, *J. Org. Chem.*, 2006, **71**, 7063.
2. H. Y. Au-Yeung, G. D. Pantoş and J. K. M. Sanders, *Chem. Commun.*, 2009, 419.
3. H. Y. Au-Yeung, G. D. Pantoş and J. K. M. Sanders, *Proc. Natl. Acad. Sci. USA*, 2009, **106**, 10466.
4. D. G. Hamilton, J. E. Davies, L. Prodi and J. K. M. Sanders, *Chem. Eur. J.*, 1998, **4**, 608.
5. H. Y. Au-Yeung, G. D. Pantoş and J. K. M. Sanders, *Angew. Chem., Int. Ed.*, 2010, DOI: 10.1002/anie.201000807..
6. D. Crunch, V. Krishnamurthy, F. Brebion, M. Karatholuvhu, V. Subramanian and T. K. Hutton, *J. Am. Chem. Soc.*, 2007, **129**, 10282.